

The Specific Vulnerabilities of Cancer Cells to the Cold Atmospheric Plasma- Stimulated Solutions

Dayun Yan^{1*}, Haitao Cui¹, Wei Zhu¹, Niki Nourmohammadi², Julian Milberg³, Lijie G. Zhang¹, Jonathan H. Sherman⁴, Michael Keidar^{1*}.

¹Department of Mechanical and Aerospace Engineering, The George Washington University, Science & Engineering Hall, 800 22nd Street, NW, Room 3550, Washington, DC 20052, USA

²Department of Epidemiology, Columbia University, Mailman School of Public Health, 722 West 168th Street, New York, NY 10032, USA

³Department of Biomedical Engineering, University of Miami, 1251 Memorial Drive McArthur Engineering Building, Coral Gables, FL, 33146-0621, USA

⁴Neurological Surgery, The George Washington University, Foggy Bottom South Pavilion, 22nd Street, NW, 7th Floor, Washington, DC 20037, USA

*Corresponding authors: Dr. Dayun Yan: ydy2012@gwmail.gwu.edu,

Dr. Michael Keidar: keidar@gwu.edu.




Fig. S1. The toxicity of NO₂⁻ on MDA-MB-231 cells. Results are presented as the mean ± s.d. of three independently repeated experiments performed in sextuplicate.