

Cell Reports, Volume 19

Supplemental Information

Definition of a High-Confidence

Mitochondrial Proteome at Quantitative Scale

Marcel Morgenstern, Sebastian B. Stiller, Philipp Lübbert, Christian D. Peikert, Stefan Dannenmaier, Friedel Drepper, Uri Weill, Philipp Höß, Reinhild Feuerstein, Michael Gebert, Maria Bohnert, Martin van der Laan, Maya Schuldiner, Conny Schütze, Silke Oeljeklaus, Nikolaus Pfanner, Nils Wiedemann, and Bettina Warscheid




Figure S1

Figure S1. Illustration of experimental strategies employed for the determination of the mitochondrial proteome and the detailed characterization of individual mitochondrial proteins by quantitative MS

Related to Figure 1

(A) Crude and gradient-purified mitochondrial fractions, prepared from differentially light (Arg0/Lys0) and heavy (Arg10/Lys8) SILAC-labeled yeast cells, were mixed in equal ratio and subsequently analyzed by LC-MS either directly without (w/o) fractionation or following various peptide and protein fractionation techniques such as high pH reversed-phase chromatography (pH 10 RP), strong cation exchange chromatography (SCX), and SDS-PAGE using Bis-Tris gradient and Tricine gels. For enzymatic protein digestion, trypsin (Tryp.), LysC, GluC, chymotrypsin (Chym.), and AspN were used as indicated. The experiment, referred to as 'pure/crude experiment', was performed in four independent biological replicates (including a label-switch). SILAC-based protein quantification and subsequent data analyses facilitated the definition of a mitochondrial core proteome.

(B) For a proteome-wide absolute quantification of yeast proteins grown on different carbon sources, a triple SILAC approach was applied. Cells were cultivated in the presence of glucose, galactose, or glycerol and either labeled with light (Arg0/Lys0), medium-heavy (Arg6/Lys4), or heavy (Arg10/Lys8) amino acids. Whole cell lysates were mixed in equal ratios, proteins were digested with trypsin (Tryp.) and peptides were fractionated by pH 10 RP followed by quantitative LC-MS analysis. The experiment, referred to as 'absolute quantification experiment', was performed in three independent biological replicates including a label-switch. Absolute protein quantification was based on the MS intensities determined by MaxQuant for light, medium-heavy, and heavy SILAC-labeled proteins following the proteomic ruler strategy (Wiśniewski et al., 2014).

(C) To globally profile submitochondrial protein localizations, protease accessibility assays were performed. Gradient-purified mitochondria isolated from differentially SILAC-labeled cells were either treated with proteases (i.e., trypsin and proteinase K) only (S1), digitonin and proteases (S2), or Triton X-100 (TX-100) and proteases (S3). Samples S1 - S3 were mixed with equal amounts of untreated mitochondria (M) serving as reference. Proteins were separated by SDS-PAGE, in-gel digested with trypsin, and analyzed by quantitative MS. The experiment, referred to as 'submitochondrial profiling experiment', was performed in three independent biological replicates including a label-switch.

(A - C) Further information given for each experimental strategy: duration of each LC-MS run and the number (#) of fractions obtained for the indicated sample processing method or LC-MS runs per replicate.




Figure S2

Figure S2. Evaluation of the sample processing regime, protein identification and quantification of pure/crude experiments

Related to Figure 2

(A) Number of protein groups identified in individual replicates following different sample fractionation protocols (I - IX; see also Fig. S1A, Tables S2B and S2C) and in total. Proteins were analyzed by LC-MS either without fractionation (I) or following gel electrophoresis using TrisTricine (II) and BisTris (III) gels, high pH reversed-phase chromatography of tryptic peptides (IV), or strong cation exchange chromatography (SCX) of peptides derived from tryptic (V), LysC (VI), GluC (VII), chymotryptic (VIII), and AspN (IX) digests. Red, mitochondrial proteins according to GO cellular component annotations; light grey, other proteins; Rep, replicate.

(B) Overlap of protein groups identified in tryptic, LysC, GluC, chymotryptic, and AspN digests (left) and impact of the protease(s) chosen for proteolytic digestion on sequence coverage and the number of protein groups identified (right). All samples were analyzed by SCX (V - IX in [A]).

(C) Cartoons illustrating the sequence coverage of individual components of the TIM23 complex identified in pure/crude experiments by SCX following proteolytic digestion with trypsin only (V in A; left) or a multi-protease digestion approach using trypsin, LysC, GluC, chymotrypsin, and AspN (V - IX in A; right). IMS, intermembrane space.

(D) Overlap of proteins identified in pure/crude experiments in this study and in previous studies targeting the mitochondrial proteome of *S. cerevisiae*. Numbers in parentheses indicate the number of proteins identified in this study, by (Ohlmeier et al., 2004) and (Prokisch et al., 2004), and the number of proteins present in the ePROMITO list comprising mitochondrial proteins reported by (Reinders et al., 2006) (global mitochondrial proteome), (Zahedi et al., 2006) (mitochondrial outer membrane proteome), and (Vögtle et al., 2012) (mitochondrial intermembrane space proteome). See also Table S2B.

(E) Overview of proteins reported to be mitochondrial in the studies shown in (D) that were not identified in this study in pure/crude experiments. Proteins were classified according to the category of annotation as mitochondrial protein in the SGD, i.e. manually curated, high-throughput (lacking firm evidence for a mitochondrial localization), computational prediction, or none. In case the annotation was based on more than one category, the protein was assigned to the category with the highest reliability (manually curated > high-throughput > computationally predicted).

(F) Multiscatter plot showing the reproducibility of protein quantification between four independent replicates. Protein abundance ratios (pure/crude) calculated for individual replicates were log₂-transformed and plotted against each other. Values in the upper left corner indicate the Pearson correlation coefficient between replicates. Rep, replicate.

(G) Overlap of proteins quantified in four individual replicates.

(H) Histogram and density curve visualizing the distribution of protein abundance changes between pure and crude mitochondria. Shown are the mean of \log_2 -transformed pure/crude ratios ($n = 4$) of proteins quantified in all replicates with at least two fractionation methods. The data show a bimodal distribution, as indicated by the density curves (solid blue and black dotted lines), with distribution center d1 at a mean \log_2 ratio of -1.47 and d2 at 0.31. For each protein it was tested, to which distribution it belongs using an equivalence test (p value < 0.01). All proteins with inconclusive classification, i.e. proteins that belonged to neither of the distributions according to the equivalence test, were further analyzed employing a two-sample two-sided t test (p value < 0.01). Statistical analysis resulted in the definition of four distinct classes as shown in Fig. 2D. For more details about the statistical analysis, refer to Supplemental Experimental Procedures.

(I) Additional data filter criteria on class 1 proteins. To exclude low abundant non-mitochondrial contaminants that co-migrated with mitochondria in sucrose density gradients from our mitochondrial core proteome, we disregarded all proteins with a standard deviation of > 0.75 of \log_2 pure/crude ratios across all replicates and a sequence coverage of $\leq 20\%$. The ratio-intensity plot (top) shows the distribution of class 1 proteins after filtering (see also Fig. 2D). Information about subcellular localizations were derived from GO cellular component annotations (Table S2B). Non-mito, class 1 proteins without previous association with mitochondria or mitochondrial subcompartments.




Figure S3

Figure S3. Assessment of proteome-wide absolute quantification experiments

Related to Figure 3

(A) Multiscatter plot showing the reproducibility of protein abundance ratios determined in three biological replicates of proteome-wide absolute quantification experiments. Abundance ratios were determined based on triple SILAC experiments (with label-switch) for proteins extracted from cells grown on galactose versus glucose (gal/glc, top row), glycerol versus glucose (gly/glc, middle row), and glycerol versus galactose (gly/gal, bottom row) (see Table S2E). Log₂-transformed ratios of individual replicates were plotted against each other. Values in the upper left corner of each plot indicate the Pearson correlation coefficient between replicates. MS intensities for light, medium-heavy, and heavy labeled proteins of this experiment were used to calculate absolute protein copy numbers for all three carbon sources according to the proteomic ruler strategy (Wiśniewski et al., 2014). Rep, replicate.

(B) Comparison between protein copy numbers determined in this study and copy numbers reported in previous global proteome studies of *S. cerevisiae*. Copy numbers determined in our study for proteins from cells grown on glucose were compared with data previously published by (Ghaemmaghami et al., 2003), (Chong et al., 2015), and (Kulak et al., 2014) (see Table S2D). The calculation of protein copy numbers in these studies was based on Western blot analyses, single cell imaging, and the 'Total Protein Approach' (Wiśniewski et al., 2012), respectively. Values in the upper left corner of the scatter plots indicate the Pearson correlation coefficient between the data of the studies.

(C) Distribution of estimated copy numbers per cell determined for mitochondrial and non-mitochondrial (Non-mito) proteins extracted from yeast grown on glucose (Glu), galactose (Gal) and glycerol (Gly). Classification of proteins as non-mitochondrial, exclusively mitochondrial (Mito only) or mitochondrial and other subcellular localizations (multiply loc.) was based on GO cellular component annotations (see Table S1).

(D) K-means clustering of proteins with altered expression in yeast grown on galactose or glycerol in relation to glucose. Bar chart indicates the number of proteins per cluster. *, clusters in which mainly mitochondria-related terms are enriched as determined in (E).

(E) GO term enrichment analysis. For clusters C07-C14, no GO term was significantly enriched. Cyto., cytochrome; SDH, succinate dehydrogenase; TM, transmembrane.

(F) Biochemical confirmation of carbon source-dependent effects on mitochondrial protein expression levels revealed by MS-based copy number estimation. Mitochondria were isolated from YPH499 $\Delta arg4$ cells that were grown in the presence of the different carbon sources glycerol, glucose or galactose in SILAC medium. 10, 20, and 40 μ g of mitochondria (protein amount) were lysed, subjected to SDS-PAGE and analyzed by immunoblotting. For direct comparison, immunoblot results are shown together with the corresponding copy numbers. Mito, mitochondria.

(G) Percentage of total protein copy numbers per cell determined for all mitochondrial proteins and mitochondrial proteins of distinct submitochondrial categories quantified in cells grown on different carbon sources as indicated.

Mito, mitochondrial; OM/IM, mitochondrial outer/inner membrane; IMS, intermembrane space; Mito DNA and RNA, associated with mitochondrial DNA and RNA biology; Fe/S biog., iron-sulfur cluster biogenesis.




Figure S4

Figure S4. Fluorescence microscopy and biochemical subcellular localization analysis of mitochondrial proteins

Related to Figure 4

(A-C) Fluorescent images of yeast cells expressing N⁷ GFP-tagged proteins. Scale bar, 5 μ m.

(D) Subcellular fractionation of yeast cells as described in Figure 4E. This supplementary figure shows the fractionations from Figure 4E with the appropriate controls plus further experiments.

(E) Fractionation of two uncharacterized proteins that do not localize to mitochondria. Cmi7 has a negative mean \log_2 ratio (pure/crude) and Cmi8 was not quantified.

PNS, post-nuclear supernatant; Mito, mitochondrial fraction; P100, microsomal fraction; S100, cytosolic fraction; MloX, Mitochondrially localized protein; CmiX, Cytosolic mini protein of ~X kDa.




Figure S5. MS-based subcellular localization analysis of mitochondrial proteins

Related to Figure 4

(A, B) Ratio-intensity plots as shown in Figures 4A and 4B highlighting individual proteins from Tables S4A and S4B. Bar charts represent from left to right the mean of normalized MS intensities of the indicated proteins for total cell lysate, the post mitochondrial supernatant, the mitochondrial fraction and gradient-purified mitochondria of wild-type yeast analyzed by LC-MS as shown in Figure 4E. Error bars indicate SEM for $n \geq 3$ and the range for $n = 2$ (see Table S2A).





A**D****B****C**

Figure S6

Figure S6. Submitochondrial profiling experiments

Related to Figure 5

(A) Multiscatter plots showing the reproducibility of protein abundance ratios across three biological replicates of MS-based global submitochondrial profiling experiments. Gradient-purified mitochondria (M) isolated from light or heavy SILAC-labeled yeast were mixed in equal ratio with mitochondria (S1), mitoplasts (S2) or lysed mitochondria (S3) that had been treated with proteases and were obtained from differentially SILAC-labeled yeast. Samples of three independent replicates were analyzed by SDS-PAGE followed by quantitative MS analysis (see Fig. S1C). For each of 24 gel slices per replicate, protein abundance ratios (S/M) were determined (Table S2H). For each protein quantified, the ratio of the slice with the highest MS intensity for M is plotted (Table S2G). Values in the upper left corner indicate the Pearson correlation coefficient between replicates. Rep, replicate.

(B) Western blots from Figures 5E and 5F together with the respective controls. Tom70, outer membrane protein; Tim10, intermembrane space protein; Tim23, intermembrane space-exposed inner membrane protein; Tim44, matrix protein peripherally attached to the inner membrane; Isd11, soluble matrix protein.

(C) Triton X-100 controls for indicated proteins. WT (for analysis of Tcd2, Pth2, Tim44) or Protein A-tagged strains were treated with 1% Triton X-100 in SEM buffer and incubated for 7 minutes on ice. Where indicated, samples were subjected to proteinase K (Prot. K) treatment at a final concentration of 7 $\mu\text{g}/\text{mL}$ for 15 minutes. Samples were subjected to SDS-PAGE and analyzed by immunoblotting. Tim44, matrix protein.

(D) Overlap of proteins identified in this study to be associated with the mitochondrial outer membrane (OM; left) or exposed to the intermembrane space (IMS; right) with previous proteomics studies targeting these submitochondrial proteomes. Numbers in parentheses indicate the number of proteins assigned to OM or IMS in our study or the work published by Zahedi et al. (2006) and Vögtle et al. (2012), respectively. Our study targets OM proteins exposed to the cytosol; Zahedi et al. (2006) detected OM proteins and a number of (precursor) proteins that are destined for internal mitochondrial compartments. Our study of IMS/IM includes soluble IMS proteins and membrane proteins (mostly of the inner membrane, IM) exposed to the IMS, whereas Vögtle et al. (2012) mainly analyzed soluble IMS proteins (and loosely membrane-attached proteins).




Figure S7

Figure S7. Mitochondrial protein interaction networks

Related to Figure 7

(A) Assembly of [³⁵S]-labeled mitochondrial proteins into high molecular weight complexes. The radiolabeled precursors were mixed with isolated mitochondria and incubated for 45 min at 30°C in the presence (+Δψ) or absence (-Δψ) of the membrane potential. Where indicated, mitochondria were subjected to proteinase K (Prot. K) treatment after import.

(B) SILAC mitochondria isolated from wild-type (WT) and Yor020w-a_{ProtA} yeast strains were solubilized with 1% digitonin and subjected to IgG affinity chromatography, followed by elution of proteins specifically bound to Yor020w-a_{ProtA} using TEV protease. Top, Ratio-versus-ratio plot visualizing the Mco10 (Yor020w-a) interaction network. Data were obtained in q-AP-MS experiments using Protein A-tagged Mco10 (Yor020w-a) as bait (Table S2J). Bottom, Samples were analyzed by SDS-PAGE and immunoblotting using the indicated antisera. Load, 0.5%; Elution, 100%.

(C) Reciprocal interaction networks of Iai11 (left) and its interaction partner Aim11 (right) obtained in q-AP-MS experiments of Protein A-tagged baits (Table S2J).

(D) Left (lanes 1-4), radiolabeled Tmh11 precursor was imported into isolated wild-type (WT) mitochondria. [³⁵S]Tmh11 was incubated with the reaction buffers for 10 min at 25°C followed by centrifugation for 10 min at 14,000 rpm. Import was started by the addition of 60 μg of mitochondria to pre-incubated samples and import reaction was allowed to proceed for the indicated time points at 25°C. Where indicated (-Δψ), membrane potential was dissipated. Non-imported precursor was digested by the addition of proteinase K. Samples were analyzed by Blue-Native electrophoresis followed by autoradiography. Right (lanes 5-12), mitochondria from wild-type (WT) and _{HA}Tmh11 were solubilized with 1% digitonin and incubated with anti-HA affinity matrix. After washing, a denaturing elution was performed and samples were subjected to SDS-PAGE and immunoblotting using antisera directed against the indicated proteins Load, 2%; Elution, 100%. Bottom, Sequence alignment of Tmh11 and the transmembrane protein 14A family members of higher eukaryotes. The conserved GxxxG transmembrane domain interaction motives are highlighted.

Table S4. Mitochondrial proteins identified and validated in this study. Related to Figures 1 and 4. (A) List of identified mitochondrial proteins that were not assigned to mitochondria previously. (B) List of mitochondrial proteins with annotation inferred from high-throughput studies without further verification. Proteins in bold were subjected to additional experimental validation of their mitochondrial localization (outlined in Figures 4, 5, 6, 7, S4, S5, S6, S7 plus Table S7 and summarized in Table S6). Coq21, COQ interacting protein; Dpa10, Delta-Psi dependent mitochondrial assembly, protein of 10 kDa; DpcX, Delta-psi ($\Delta\psi$)-dependent import and cleavage, protein of ~X kDa; DpiX, Delta-psi ($\Delta\psi$)-dependent import, protein of ~X kDa; Iai11, Interactor of Aim11; Mgp12, Mitochondrial glutaredoxin-like protein of 12 kDa; McoX, Mitochondrial class one protein of X kDa; MinX, Mini mitochondrial protein of ~X kDa; MloX, Mitochondrially localized protein, Pdl32, Protein of dual localization, protein of 32 kDa; RciX, Respiratory chain interacting protein of ~X kDa; Rso55, Mitochondrial protein related to spastic paraplegia with optic atrophy and neuropathy SPG55; Tmh11, TMEM14 homolog of 11 kDa; Tmh18, Mitochondrial TMEM205 homolog of 18 kDa; Tml25, Acyl-protein thioesterase with multiple localizations, protein of 25 kDa.

Systematic name	Gene name	kDa	Systematic name	Gene name	kDa	Systematic name	Gene name	kDa	Systematic name	Gene name	kDa
YML050W	AIM32	36.0	YER145C	FTR1	45.7	YJR074W	MOG1	24.3	YDR201W	SPC19	18.9
YNL094W	APP1	66.1	YNL133C	FYV6	20.0	YJL205C	NCE101	6.3	YKR031C	SPO14	195.2
YGR230W	BNS1	15.9	YCL026C-B	HBN1	21.0	YJL126W	NIT2	34.7	YER046W	SPO73	16.6
YJL158C	CIS3	23.2	YKL101W	HSL1	169.6	YOR056C	NOB1	51.7	YGL169W	SUA5	46.5
YOR093C	CMR2	186.9	YPL015C	HST2	40.0	YNL129W	NRK1	27.7	YOR081C	TGL5	84.7
YBR230W-A	COQ21	7.6	YER092W	IES5	14.3	YGL111W	NSA1	51.9	YLR118C	TML25	24.7
YFL001W	DEG1	50.9	YNL106C	INP52	133.3	YBR060C	ORC2	71.3	YDR449C	UTP6	52.4
YLR307C-A	DPA10	9.6	YLL033W	IRC19	27.4	YHR063C	PAN5	42.8	YEL040W	UTR2	49.9
YKL065W-A	DPC7	8.5	YIL156W-B	MCO8	8.2	YLR151C	PCD1	39.8	YGR281W	YOR1	166.7
YGL041W-A	DPC13	18.1	YKL018C-A	MCO12	11.8	YGR087C	PDC6	61.6	YDR349C	YPS7	64.5
YOR114W	DPI34	35.0	YGR053C	MCO32	32.2	YMR087W	PDL32	32.1	YBL055C		47.4
YMR130W	DPI35	35.3	YLR017W	MEU1	37.9	YDR406W	PDR15	172.3	YDL177C		19.1
YDL219W	DTD1	16.7	YMR210W	MGL2	51.4	YBR022W	POA1	19.9	YHL012W		56.0
YML080W	DUS1	48.1	YDR286C	MGP12	13.4	YCL047C	POF1	9.7	YJR149W		45.2
YKL204W	EAP1	69.8	YMR182W-A	MIN3	3.1	YBR087W	RFC5	39.9	YKL071W		28.0
YKR076W	ECM4	43.3	YBL039W-B	MIN6	6.9	YER047C	SAP1	100.3	YMR187C		50.3
YGR200C	ELP2	89.4	YBR201C-A	MIN7	7.7	YLR022C	SDO1	28.3	YNL247W		87.5
YDR512C	EM11	21.1	YPR010C-A	MIN8	7.9	YMR059W	SEN15	14.9	YOR131C		24.8
YDR261C	EXG2	63.5	YKL023C-A	MIN9	8.5	YJL145W	SFH5	34.4	YPL034W		18.9
YMR113W	FOL3	47.8	YFR032C-B	MIN10	10.0	YKL051W	SFK1	40.5			
YHR049W	FSH1	27.3	YLR049C	MLO50	49.5	YNR015W	SMM1	42.8			

Systematic name	Gene name	kDa	Systematic name	Gene name	kDa	Systematic name	Gene name	kDa	Systematic name	Gene name	kDa
YJL200C	ACO2	86.6	YFR044C	DUG1	52.9	YOR020W-A	MCO10	9.6	YDL104C	QRI7	45.5
YKL192C	ACP1	13.9	YDR125C	ECM18	53.2	YDR381C-A	MCO13	12.7	YLR084C	RAX2	133.9
YMR064W	AEP1	59.8	YLR390W	ECM19	12.5	YHL018W	MCO14	14.0	YIL077C	RCI37	37.0
YER080W	AIM9	72.4	YBR163W	EXO5	67.6	YPL109C	MCO76	76.2	YKL133C	RCI50	54.5
YER087W	AIM10	65.9	YFR019W	FAB1	257.4	YGR012W	MCY1	42.8	YOR286W	RDL2	16.7
YER093C-A	AIM11	15.8	YER183C	FAU1	24.1	YJL102W	MEF2	91.3	YDR065W	RRG1	42.9
YHL021C	AIM17	53.1	YDR070C	FMP16	10.9	YML007C-A	MIN4	4.4	YOR305W	RRG7	28.0
YHR198C	AIM18	36.5	YBR047W	FMP23	20.5	YMR252C	MLO1	15.6	YPR116W	RRG8	31.1
YJL131C	AIM23	41.5	YJL161W	FMP33	20.2	YJR039W	MLO127	127.4	YAR008W	SEN34	31.3
YJR100C	AIM25	37.5	YPL222W	FMP40	78.3	YIR021W	MRS1	41.3	YMR066W	SOV1	104.8
YMR003W	AIM34	22.8	YNL168C	FMP41	28.8	YER077C	MRX1	79.6	YGR236C	SPG1	10.5
YMR157C	AIM36	29.1	YKR049C	FMP46	15.7	YPL041C	MRX11	24.2	YLR389C	STE23	117.6
YOL053W	AIM39	45.9	YER004W	FMP52	25.1	YJR003C	MRX12	59.8	YLR305C	STT4	214.6
YOR215C	AIM41	21.2	YOR271C	FSF1	35.4	YPL168W	MRX4	48.9	YHR003C	TCD1	48.9
YHR199C	AIM46	34.1	YDR019C	GCV1	44.5	YJL147C	MRX5	44.9	YKL027W	TCD2	50.3
YER073W	ALD5	56.7	YMR189W	GCV2	114.4	YNL211C	MRX7	10.7	YJR019C	TES1	40.3
YKL157W	APE2	107.8	YLR091W	GEP5	33.9	YDL027C	MRX9	48.3	YJR085C	TMH11	11.3
YGR286C	BIO2	41.9	YGL057C	GEP7	33.0	YGL226W	MTC3	14.5	YPR098C	TMH18	17.7
YBL098W	BNA4	52.4	YDR305C	HNT2	24.8	YNL063W	MTQ1	35.9	YOR251C	TUM1	34.2
YKL208W	CBT1	31.2	YBL059W	IAI11	22.3	YAL029C	MYO4	169.3	YLL040C	VPS13	357.8
YER061C	CEM1	47.6	YER086W	ILV1	63.8	YPR155C	NCA2	70.9	YHL014C	YLF2	45.7
YGR207C	CIR1	28.8	YMR108W	ILV2	74.9	YOL042W	NGL1	42.4	YBR054W	YRO2	38.7
YCR005C	CIT2	51.4	YJR016C	ILV3	62.9	YLR351C	NIT3	32.5	YHR017W	YSC83	44.2
YLR087C	CSF1	338.2	YJL082W	IML2	82.5	YJR062C	NTA1	51.9	YDL157C		13.6
YOR022C	DDL1	81.8	YGL085W	LCL3	32.1	YGR178C	PBP1	78.8	YDR061W		61.2
YOR236W	DFR1	24.3	YLR239C	LIP2	37.2	YPR002W	PDH1	57.7	YGR015C		37.9
YPL107W	DPC25	28.6	YOR196C	LIP5	46.3	YJL023C	PET130	39.8	YKL162C		46.5
YGR021W	DPC29	31.7	YIL094C	LYS12	40.1	YHR189W	PTH1	21.0	YKR070W		39.4
YJL133C-A	DPI8	7.7	YDR234W	LYS4	75.2	YBL057C	PTH2	22.4	YLR283W		36.6
YNR040W	DPI29	28.7	YJL127C-B	MCO6	6.0	YJR111C	PXP2	32.2			

Table S6: Mitochondrial proteins with localization validated in this study. Related to Figures 4-7 and S4-S7.

Systematic name	Gene name	Dual loc.	kDa	Subcell. fractionation	Δy dependent import into mitochondria	Mitochondrial GFP-signal	Submito. fractionation MS (exp. val. ✓)	Additional experiments performed in this study	Copy# Gly/Glc	Pred. no. TM helices (TMHMM)	Previous studies	Name description
									-6 log ₂ ratio 6			
YER093C-A	AIM11		15.8	✓	import	N-term	IMS/IM	Co-IP (Iai11, Mtc3, Gep7)		2	§	Altered inheritance rate of mitochondria
YBR230W-A	COQ21		7.6	✓	import		matrix/IM ✓	BN, Co-IP (Coqx, Cat5)				COQ interacting protein
YFL001W	DEG1	✓	50.9	✓	import				242/363			Depressed growth rate
YLR307C-A	DPA10		9.6	✓	assembly			BN				Delta-Psi dependent mitochondrial assembly, protein of 10 kDa
YKL065W-A	DPC7		8.5	✓	import & cleavage	C-term	matrix/IM	BN	7824/1704	1		Delta-Psi dependent mitochondrial import and cleavage, protein of 7 kDa
YGL041W-A	DPC13		18.1	✓	import & cleavage	N-term	matrix/IM ✓		689/525			Delta-Psi dependent mitochondrial import and cleavage, protein of ~13 kDa
YPL107W	DPC25		28.6	✓	import & cleavage		matrix/IM	BN	367/168		*	Delta-Psi dependent mitochondrial import and cleavage, protein of ~25 kDa
YGR021W	DPC29		31.7	✓	import & cleavage				595/231		‡††	Delta-Psi dependent mitochondrial import and cleavage, protein of ~29 kDa
YJL133C-A	DPI8		7.7	✓	import	N-term	matrix/IM ✓	BN	1139/554		‡††	Delta-Psi dependent mitochondrial Import, protein of 8 kDa
YNR040W	DPI29		28.7	✓	import & assembly	N-term	matrix/IM		779/594		*‡††	Delta-Psi dependent mitochondrial Import, protein of 29 kDa
YOR114W	DPI34		35.0	✓	import & assembly							Delta-Psi dependent mitochondrial Import, protein of ~34 kDa
YMR130W	DPI35		35.2	✓	import		matrix/IM					Delta-Psi dependent mitochondrial Import, protein of 35 kDa
YKR076W	ECM4	✓	43.3	✓					363/467			Extracellular mutant
YLR390W	ECM19		12.5	✓			IMS/IM		7027/1368	1	‡	Extracellular mutant
YDR070C	FMP16		10.9	✓	import & cleavage	N-term	matrix/IM ✓	BN	844/598		‡††	Found in mitochondrial proteome
YJL161W	FMP33		20.2	✓		N-term	IMS/IM		66971/3414		‡††	Found in mitochondrial proteome
YOR271C	FSF1		35.4	✓		N-term			3729/1341		‡†	Found in mitochondrial proteome
YHR059W	FYV4		15.3	✓		N-term	matrix/IM		8923/11250	4	*‡††§	Fungal sideroflexin 1
YCL026C-B	HBN1	✓	21.0	✓	import				391/201		*‡	Function required for yeast viability
YBL059W	IAI11		22.3	✓		N-term	IMS/IM	Co-IP (Aim11, Mtc3, Gep7)	4571/1469	2	*‡§	Homologous to bacterial nitroreductases
YGL085W	LCL3		32.0	✓		N-term	matrix/IM		795/296	1	*	Interactor of Aim11
YJL127C-B	MCO6		6.0	✓		N-term	OM				§	Long chronological lifespan 3
YIL156W-B	MCO8		8.2	✓			IMS/IM ✓		4789/2340			Mitochondrial class one protein of 6 kDa
YOR020W-A	MCO10		9.6	✓			IMS/IM	Co-IP (CV)	2788/853	1	‡††	Mitochondrial class one protein of 8 kDa
YKL018C-A	MCO12		11.8	✓		N-term	OM		522/165		‡††	Mitochondrial class one protein of 10 kDa
YDR381C-A	MCO13		12.7	✓			IMS/IM	BN	2555/1411	1	‡††	Mitochondrial class one protein of 12 kDa
YHL018W	MCO14		14.0	✓	import & assembly		matrix/IM ✓	BN	165/117		*	Mitochondrial class one protein of 13 kDa
YGR053C	MCO32		32.2	✓		N-term	matrix/IM		10/3			Mitochondrial class one protein of 14 kDa
YPL109C	MCO76		76.1	✓		N-term	IMS/IM		100/35		*‡†	Mitochondrial class one protein of 32 kDa
YLR017W	MEU1	✓	37.9	✓	import		matrix/IM		1852/3152		*‡†	Mitochondrial class one protein of 76 kDa
YDR286C	MGP12		13.4	✓	import & cleavage				989/828			Multicopy enhancer of UAS2
YMR182W-A	MIN3		3.1	✓		C-term				1		Mitochondrial glutaredoxin-like protein of 12 kDa
YML007C-A	MIN4		4.4	✓							*	Mini mitochondrial protein of 3 kDa
YBL039W-B	MIN6		6.9	✓		C-term	OM	BN		1		Mini mitochondrial protein of 4 kDa
YBR201C-A	MIN7		7.7	✓			IMS/IM ✓			1		Mini mitochondrial protein of ~6 kDa
YPR010C-A	MIN8		7.9	✓			IMS/IM ✓	BN		1		Mini mitochondrial protein of ~7 kDa
YKL023C-A	MIN9		8.5	✓		C-term	matrix/IM		13017/2464	1		Mini mitochondrial protein of 8 kDa
YFR032C-B	MIN10		10.0	✓		C-term				1		Mini mitochondrial protein of 9 kDa
YMR252C	MLO1		15.6	✓		N-term					*	Mini mitochondrial protein of 10 kDa
YLR049C	MLO50		49.5	✓								Mitochondrially localized protein
YJR039W	MLO127		127.4	✓							‡†	Mitochondrially localized protein of 50 kDa
YPL041C	MRX11		24.2	✓		N-term	matrix/IM			2	§	Mitochondrially localized protein of 127 kDa
YGL226W	MTC3		14.5	✓			IMS/IM		1268/452	1	††	Mitochondrial organization of gene expression (MIOREX)
YJL205C	NCE101		6.3	✓					931/389	1		Maintenance of telomere capping
YMR087W	PDL32	✓	32.1	✓			matrix/IM		221/139			Nonclassical export
YBL057C	PTH2		22.4	✓			OM ✓		4545/6624	1	‡†††	Protein of dual localization, protein of 32 kDa
YJR111C	PXP2	✓	32.2	✓					1786/1763		*	Peptidyl-trna hydrolase
YIL077C	RCI37		37.0	✓	import & assembly	N-term	IMS/IM ✓	BN, Co-IP (CIII, CIV, m-AAA)	540/258	2	*‡	Peroxisomal protein
YKL133C	RCI50		54.5	✓	import & cleavage	N-term	IMS/IM	Co-IP (CIII, CIV, i-AAA)		1	§	Respiratory chain interacting protein of 37 kDa
YOR286W	RDL2		16.7	✓	import & cleavage				6638/4974		‡†††	Respiratory chain interacting protein of ~50 kDa
YNL213C	RRG9		25.3	✓	import & cleavage		matrix/IM ✓	BN			‡††	Rhodanese-like protein
YLR281C	RSO55		18.2	✓	import & cleavage	N-term		BN			‡†	Required for respiratory growth
YJL145W	SFH5		34.4	✓		N-term	OM				*‡	Mitochondrial protein related to spastic paraplegia with optic atrophy and neuropathy SPG55
YNR015W	SMM1	✓	42.8	✓	import		matrix/IM		2441/2547			Sec fourteen homolog
YGL169W	SUA5	✓	46.5	✓			IMS/IM		788/1081			Suppressor of mitochondrial mutation
YKL027W	TCD2		50.3	✓			OM ✓		784/699			Suppressor of upstream AUG
YJR019C	TES1	✓	40.3	✓	import		matrix/IM		4295/3584	1	*‡†††	tRNA Threonylcarbamoyladenosine dehydratase
YJR085C	TMH11		11.3	✓	import	N-term	OM	BN, Co-IP (GxxxG-cont. prot.)	10404/1398		‡††	Thioesterase
YPR098C	TMH18		17.7	✓		N-term			1752/937	3	‡†	TMEM14 homolog of 11 kDa
YLR118C	TML25	✓	24.7	✓	import				5832/2902	3	*‡	Mitochondrial TMEM205 homolog of 18 kDa
YOR251C	TUM1	✓	34.2	✓	import		matrix/IM		2635/2937			Acyl-protein thioesterase with multiple localizations, protein of 25 kDa
									7954/6424			Thiouridine modification

* Huh WK, et al. (2003) Global analysis of protein localization in budding yeast. Nature 425(6959):686-91

† Perocchi F, et al. (2006) Assessing Systems Properties of Yeast Mitochondria through an Interaction Map of the Organelle. PLoS Genet 2(10):e170

‡ Sickmann A, et al. (2003) The proteome of Saccharomyces cerevisiae mitochondria. Proc Natl Acad Sci U S A 100(23):13207-12

†† Reinders J, et al. (2006) Toward the complete yeast mitochondrial proteome: multidimensional separation techniques for mitochondrial proteomics. J Proteome Res 5(7):1543-54

††† Renvoise M, et al. (2014) Quantitative variations of the mitochondrial proteome and phosphoproteome during fermentative and respiratory growth in Saccharomyces cerevisiae. J Proteomics 106:140-50

†††† Wysocki R, et al. (1999) Disruption and basic phenotypic analysis of 18 novel genes from the yeast Saccharomyces cerevisiae. Yeast 15(2):165-71

‡‡ Yofe I, et al. (2016) One library to make them all: streamlining the creation of yeast libraries via a SWAp-Tag strategy. Nat Methods 13(4):371-8

* Zahedi RP, et al. (2006) Proteomic analysis of the yeast mitochondrial outer membrane reveals accumulation of a subclass of preproteins. Mol Biol Cell 17(3):1436-50

Table S7. Dual-localized mitochondrial proteins. Related to Figure 6. To identify soluble mitochondrial proteins with dual/multiple cellular localization, filtered high-confidence mitochondrial class 1 proteins were selected from Table S1 when they displayed a significant amount in the post mitochondrial supernatant (PNS/total > 0.5) and in both mitochondrial fractions (pure mito/crude mito > 0). α -helical membrane proteins (transmembrane helix TMHMM prediction) and β -barrel membrane proteins were removed. The list of 57 mitochondrial proteins includes proteins with known (grey) dual/multiple cellular localizations, proteins validated in this study (green) and high-confidence candidates. Upon completion of this study we noticed that Tum1 was previously imported as y-rho into isolated mitochondria (Dubaquié et al., 1998) in full agreement with our study.

Systematic name	PMS/ Total	Gene name	Mitochondrial proteome class 1	Mitochondrial localization (other evidence)	Extra-mito. annotation	Extra-mitochondrial localization	Gene description
YKL157W	1.42	APE2	✓	HT	Manual	Plasmamembrane	Aminopeptidase yscll
YMR098C	0.87	ATP25	✓	Manual	Manual		Mito. protein required for Oli1p ring formation and stability of Oli1p (Atp9p) mRNA
YJL060W	0.67	BNA3	✓	Manual	Manual	Cytosol	Kynurenine aminotransferase
YMR038C	1.60	CCS1	✓	Manual	Manual	Cytosol, Nucleus	Copper chaperone for superoxide dismutase Sod1p
YDL164C	0.99	CDC9	✓	Manual	Manual	Nucleus	DNA ligase I found in nucleus and mitochondria
YGR255C	5.90	COQ6	✓	Manual	Manual		Flavin-dependent monooxygenase involved in ubiquinone biosynthesis
YFL001W	1.13	DEG1	✓	This study	Manual	Cytosol (this study), Nucleus	tRNA:pseudouridine synthase
YHR011W	6.21	DIA4	✓	Manual	Manual	Cytosol	Probable mitochondrial seryl-tRNA synthetase
YFR044C	1.26	DUG1	✓	HT	Manual		Cys-Gly metallo-di-peptidase
YKR076W	1.16	ECM4	✓	This study	Manual	Cytosol & Microsome (this study)	S-glutathionyl-(chloro)hydroquinone reductase (GS-HQR)
YIL098C	2.45	FMC1	✓	Manual	Manual		Mito. matrix protein that is required for assembly or stability at high temperature of the F1 sector of mitochondrial F1F0 ATP synthase
YNL256W	0.67	FOL1	✓	Manual	Manual		Multifunctional enzyme of the folic acid biosynthesis pathway
YMR113W	0.53	FOL3	✓	Manual	Manual		Dihydrofolate synthetase that is involved in folic acid biosynthesis
YHR049W	2.82	FSH1	✓	Manual	HT	Cytoplasm, Nucleus	Putative serine hydrolase
YHR100C	0.91	GEP4	✓	Manual	Manual		Mitochondrial phosphatidylglycerophosphatase (PGP phosphatase)
YPL091W	1.16	GLR1	✓	Manual	Manual	Cytosol, Nucleus	Cytosolic and mitochondrial glutathione oxidoreductase
YOL059W	1.37	GPD2	✓	Manual	Manual	Cytosol	NAD-dependent glycerol 3-phosphate dehydrogenase
YCL026C-B	1.10	HBN1	✓	This study	HT	Cytosol, Nucleus	Protein of unknown function that is similar to bacterial nitroreductases
YMR207C	1.22	HFA1	✓	Manual	Manual		Mitochondrial acetyl-coenzyme A carboxylase that catalyzes production of malonyl-CoA in mitochondrial fatty acid biosynthesis
YDR305C	1.20	HNT2	✓	HT	HT	Cytoplasm, Nucleus	Dinucleoside triphosphate hydrolase
YPR033C	1.15	HTS1	✓	Manual	Manual	Cytosol	Cytoplasmic and mitochondrial histidine tRNA synthetase
YPR083W	0.66	MDM36	✓	Manual	Manual		Component of the mitochondria-ER-cortex-ancor (MECA)
YLR017W	3.05	MEU1	✓	This study	Manual	Cytosol (this study)	Methylthioadenosine phosphorylase (MTAP)
YMR002W	0.88	MIX17	✓	Manual	HT	Cytoplasm, Nucleus	Mitochondrial intermembrane space protein that is required for normal oxygen consumption
YOR274W	0.71	MOD5	✓	Manual	Manual	Cytosol, Nucleus	Delta 2-isopentenyl pyrophosphate:tRNA isopentenyl transferase
YNL306W	1.70	MRPS18	✓	Manual	Manual		Mitochondrial ribosomal protein of the small subunit
YBR251W	2.15	MRPS5	✓	Manual	Manual		Mitochondrial ribosomal protein of the small subunit
YPL104W	0.62	MSD1	✓	Manual	Manual		Mitochondrial aspartyl-tRNA synthetase that is required for acylation of aspartyl-tRNA
YCL033C	1.21	MXR2	✓	Manual	Manual		Methionine-R-sulfoxide reductase
YAL029C	0.87	MYO4	✓	HT	Manual		Type V myosin motor involved in actin-based transport of cargos
YGL221C	1.19	NIF3	✓	Manual	Manual	Cytosol	Protein of unknown function
YLR351C	1.36	NIT3	✓	HT	Manual		Nit protein
YGR178C	0.58	PBP1	✓	HT	Manual	Cytoplasm, Nucleus (HT)	Component of glucose deprivation induced stress granules that is involved in P-body-dependent granule assembly
YMR087W	1.02	PDL32	✓	This study	Manual	Cytosol (this study)	Putative ADP-ribose-1''-monophosphatase that converts ADP-ribose-1''-monophosphate to ADP-ribose
YCL057W	1.25	PRD1	✓	Manual	Manual	Cytoplasm, Golgi, Vacuole	Zinc metalloendopeptidase
YNL292W	1.25	PUS4	✓	Manual	Manual	Nucleus	Pseudouridine synthase
YJR111C	0.66	PXP2	✓	This study	Manual	Cytosol (this study), Peroxisome	Peroxisomal matrix protein with naturally active promoter
YKL113C	1.41	RAD27	✓	Manual	Manual	Cytosol, Nucleus	5' to 3' exonuclease / 5' flap endonuclease
YLR059C	1.37	REX2	✓	Manual	Manual		3'-5' RNA exonuclease
YBR087W	1.85	RFC5	✓	Manual	Manual	Nucleus	Subunit of heteropentameric Replication factor C (RF-C)
YLR139C	2.31	SLS1	✓	Manual	Manual		Mitochondrial membrane protein that coordinates this studyression of mitochondrially-encoded genes
YNR015W	0.97	SMM1	✓	This study	HT	Cytosol (this study), Nucleus	Dihydrouridine synthase
YMR066W	8.73	SOV1	✓	HT	Manual		Mitochondrial protein of unknown function
YLR389C	1.32	STE23	✓	HT	Manual		Metalloprotease that is involved in N-terminal processing of pro-a-factor to mature form
YLR305C	2.13	STT4	✓	HT	Manual	Plasmamembrane	Phosphatidylinositol-4-kinase that functions in the Pkc1p protein kinase pathway
YGL169W	2.41	SUA5	✓	This study	HT	Cytosol (this study)	Protein involved in threonylcarbamoyl adenosine biosynthesis
YGR046W	2.90	TAM41	✓	Manual	Manual		Mitochondrial phosphatidate cytidyltransferase (CDP-DAG synthase)
YJR019C	0.58	TES1	✓	This study	Manual	Peroxisome	Peroxisomal acyl-CoA thioesterase
YLR118C	0.97	TML25	✓	This study	Manual	Cytosol (this study)	Acyl-protein thioesterase responsible for depalmitoylation of Gpa1p
YHR070W	4.83	TRM5	✓	Manual	Manual	Cytoplasm	tRNA(m(1)G37)methyltransferase
YOR251C	0.87	TUM1	✓	This study	Manual	Cytosol (this study)	Rhodanese domain sulfur transferase
YGR094W	1.12	VAS1	✓	Manual	Manual	Cytoplasm	Mitochondrial and cytoplasmic valyl-tRNA synthetase
YLL040C	1.02	VPS13	✓	HT	Manual	Golgi, Vacuole	Protein involved in prospore membrane morphogenesis
YBL055C	1.33	YBL055C	✓	Manual	Manual		3'-->5' exonuclease and endonuclease
YGR234W	0.81	YHB1	✓	Manual	Manual	Cytosol	Nitric oxide oxidoreductase
YJR149W	2.43	YJR149W	✓	Manual	Manual		Putative protein of unknown function
YNL247W	1.26	YNL247W	✓	Manual	Manual		Cysteinyl-tRNA synthetase

SUPPLEMENTAL EXPERIMENTAL PROCEDURES

Media, growth conditions and metabolic labeling

Yeast cells were grown at 30°C and 160 rpm in in YPG (1% [w/v] yeast extract, 2% [w/v] peptone, 3% [v/v] glycerol), YPD (1% [w/v] yeast extract, 2% [w/v] peptone, 2% [w/v] glucose), or SC medium (0.17% [w/v] yeast nitrogen base [YNB] without amino acids, 0.5% [w/v] ammonium sulfate, 3% [v/v] glycerol, 20 mg/L of L-histidine, L-tryptophan, L-methionine, adenine and uracil, 23 mg/L of L-arginine and L-lysine, 30 mg/L of L-isoleucine and L-tyrosine, 50 mg/L of L-phenylalanine, 100 mg/L of L-leucine, 150 mg/L of L-valine, and 200 mg/L of L-threonine and proline). To assess differences in protein expression levels in YPH499 Δ *arg4* after growth on different carbon sources, cells were grown in SC medium containing 3% (w/v) glycerol, 2% (w/v) galactose or 2% (w/v) glucose. For metabolic labeling of yeast, media were supplemented with stable isotope-coded amino acids (Euriso-Top GmbH), i.e. 'heavy' arginine ($^{13}\text{C}_6/^{15}\text{N}_4$) and lysine ($^{13}\text{C}_6/^{15}\text{N}_2$) or 'medium-heavy' arginine ($^{13}\text{C}_6/^{14}\text{N}_2$) and lysine ($^4\text{H}_2$) instead of the respective 'light' amino acids. To ensure complete incorporation of isotopically labeled amino acids, cells were grown for at least five cell doublings in precultures and four doublings in main cultures. Cells were generally harvested during exponential growth phase.

For selection of yeast strains on solid media, YPD plates (1% [w/v] yeast extract, 2% [w/v] peptone, 2% [w/v] glucose, 2.5% [w/v] Bacto™ Agar) containing antibiotics or minimal medium plates (0.67% [w/v] YNB without amino acids [Becton, Dickinson and Company Inc.], appropriate amino acid drop-out mix [MP Biomedicals LLC], 2% [w/v] glucose, 2.5% Bacto™ Agar [Becton, Dickinson and Company Inc.]) lacking the amino acid used as selection marker were used.

Generation of yeast strains

All *Saccharomyces cerevisiae* strains used in this study have been derived from the wild type strains BY4741 (*MATa*, *his3 Δ 1*, *leu2 Δ 0*, *met15 Δ 0*, *ura3 Δ 0*), BY4742 (*Mata*, *his3 Δ 1*, *leu2 Δ 0*, *met15 Δ 0*, *ura3 Δ 0*, *lys2 Δ 0*) or YPH499 (*MATa*, *ura3-52*, *lys2-801*, *ade2-101*, *trp1- Δ 63*, *his3- Δ 200*, *leu2- Δ 1*).

C-terminal Protein A-, HA- or EGFP-tagging of chromosomal genes was performed by introducing the genetic information for the respective tag, followed by a selection marker, in front of the stop codon of the respective gene. In this study, the *kanMX4*, *kanMX6*, *HIS3MX6* (Knop et al., 1999), and *hphNT1* cassettes (Janke et al., 2004) served as selection markers. Chromosomal deletions were obtained by substitution of the respective gene by a *kanMX6* cassette (Longtine et al., 1998). For generation of YPH499 Δ *arg4* strain, the *kanMX4* selection marker was

introduced (von der Malsburg et al., 2011). Yeast cells were transformed as previously described (Gietz and Woods, 2002) and transformants were selected on minimal medium -HIS plates. Selections of strains containing an antibiotics resistance cassette were performed on YPD plates supplemented with 200 μ M KP_i and 200 mg/L G418 (Enzo Biochem Inc.; ALX-380-013-G005) for selection of strains containing a *kanMX6* cassette or 300 mg/L hygromycin B (Carl Roth GmbH + Co. KG; CP13.3) for selection of strains containing a *hphNT1* cassette. Chromosomal insertion of *hphNT1* and *kanMX6* cassettes at the correct position was confirmed by colony PCR and/or immunoblotting of whole cell yeast extracts using antibodies raised against the Protein A, HA or GFP epitope tag. For colony PCR, a tiny number of cells was transferred with a toothpick to 10 μ L of a 1 mg/mL Zymolyase solution (Zymolyase[®]-20T, Nacalai Tesque Inc.) and incubated for 10 min at 25°C. Afterwards, forward and reverse primers that bind to the 5'-UTR and 3'-UTR of the respective open reading frame were added. For some of the Protein A-tagged strains a reverse primer was employed that binds to the linker region of the Protein A tag. RedMastermix (2X) (Genaxxon Bioscience GmbH; M3029.0500) was used for amplification of DNA fragments. For the generation of N-terminal GFP-tagged proteins, strain BY4741 was genomically transformed to tag proteins at their start codon with GFP under the control of the constitutive NOP1 gene promoter. Using the SWAT approach, excision of the selection cassette and NOP1 promoter resulted in restoration of the gene's endogenous promoter (Yofe et al., 2016). Tagging was verified using a genomic PCR check.

The pRS425-_{HA}YJR085C plasmid was generated as follows: the YJR085C open reading frame including 537 nucleotides upstream (containing the endogenous promoter) and 369 nucleotides downstream (comprising the endogenous terminator) was amplified from yeast genomic DNA. At their 5'-ends, the forward (YJR085C_pRS425_FW) and reverse (YJR085C_pRS425_REV) primers were fused to a HindIII or BamHI cleavage site, respectively. The resulting HindIII-P_{YJR085C}-YJR085C (*S. cerevisiae*)-T_{YJR085C}-BamHI PCR fragment was cloned into pRS425 plasmid using HindIII and BamHI restriction enzymes. Site-directed mutagenesis was employed to introduce the N-terminal HA-tag in front of YJR085C. To this end the forward and reverse primers NHA_YJR085C_FW and NHA_YJR085C_Rev as well as the QuikChange[™] Site-Directed Mutagenesis Kit (Agilent Technologies Inc.) were used.

_{HA}YJR085C and its corresponding wild-type strain (YPH499 + *pRS425*) were generated by transforming YPH499 cells with the pRS425 plasmid encoding for the _{HA}YJR085C protein (under control of its endogenous promoter and terminator) or the empty plasmid followed by several rounds of selection on minimal medium plates depleted from leucine.

To generate the pFA6a-TEV-ProtA-7His-hphNT1 plasmid, the TEV-ProtA-7His module was amplified from pYM10 plasmid (Knop et al., 1999) using TEV-ProtA_fwd and TEV-ProtA_rev primers and cloned into pFA6a-hphNT1 plasmid (Janke et al., 2004) using HindIII and XmaI restriction sites.

To generate the BY4741 $\Delta arg4$ strain, BY4741 and BY4742 $\Delta arg4$ (Euroscarf) were crossed and several tetrads were dissected. Correct genotype (*MATa*; *ura3 Δ 0*, *leu2 Δ 0*, *his3 Δ 1*, *lys2 Δ 0*, *met15 Δ 0*, *arg4::kanMX4*) was selected by replica plating. To this end, minimal media devoid of either arginine, methionine or lysine were used. Mating type was determined by colony PCR using the primers MATlocus_fw, Mat(a)_rv and Mat(alpha)_rv (Huxley et al., 1990).

Preparation of cell lysates, subcellular fractionation, and isolation of mitochondria

Subcellular fractionation and isolation of mitochondria for quantitative MS experiments were performed as described before (Meisinger et al., 2000) with slight modifications. Differentially SILAC-labeled cells were harvested by centrifugation for 5 min at 3,000 x g and RT, washed with deionized water, and incubated for 20 min at 160 rpm and 30°C in 2 mL of DTT buffer (100 mM Tris-H₂SO₄ [pH 9.4], 10 mM DTT) per g wet weight (ww). Following centrifugation (5 min, 3,000 x g, RT), cells were washed with 7 mL/g ww zymolyase buffer (1.2 M sorbitol, 20 mM K₃PO₄ [pH 7.4]) and incubated for 45 min at 160 rpm and 30°C in 7 mL/g ww zymolyase buffer containing 3 mg/g ww Zymolyase 20-T (MP Biomedicals Life Sciences) to digest cell walls. Spheroplasts were harvested by centrifugation (5 min, 3,000 x g, 4°C) and homogenized in 7 mL/g ww ice-cold homogenization buffer (0.6 M sorbitol, 10 mM Tris-HCl [pH 7.4], 1 mM PMSF [dissolved in isopropanol]) containing 1 mM EDTA using a glass-Teflon potter (15 strokes). Cell debris and nuclei were removed from the homogenate (referred to as 'cell lysate' in this work) by centrifugation for 5 min at 1,500 x g and 4°C followed by centrifugation of the supernatants for 5 min at 4,000 x g and 4°C. Crude mitochondrial fractions were pelleted from the resulting postnuclear supernatants (PNS) by centrifugation for 15 min at 12,000 x g and 4°C. The post-mitochondrial supernatants (PMS) were taken off and the pellets were resuspended in SEM buffer (250 mM sucrose, 1 mM EDTA, 10 mM MOPS-KOH [pH 7.2]). To obtain mitochondrial fractions of higher purity, crude mitochondria were loaded onto sucrose density gradients consisting of 2 mL of 60%, 6 mL of 32%, 2 mL of 23%, and 2 mL of 15% (w/v each) sucrose in EM buffer (10 mM MOPS-KOH [pH 7.2], 1 mM EDTA). Following centrifugation for 1 h at 134,000 x g and 2°C, pure mitochondrial fractions were collected from the interface between 60% and 32% sucrose, diluted with the two-fold volume of SEM buffer, pelleted for 10 min at 12,000 x g and 2°C, resuspended in 100 μ L/g ww of SEM buffer, shock-frozen in liquid nitrogen, and stored at -80°C until further use.

To reveal differences in protein abundance between crude mitochondrial fractions and gradient-purified mitochondria and, thus, identify putative mitochondrial proteins, crude and pure mitochondria from differentially SILAC-labeled cells were mixed in equal ratios based on protein concentration and analyzed by LC-MS following different protein and peptide fractionation methods. This experiment (referred to as 'pure/crude' experiment in this work) was performed in four biological replicates including label-switch (see Figure S1A). For subcellular profiling experiments, performed to verify the mitochondrial localization of candidate proteins, aliquots of subcellular fractions (i.e., cell lysates, the post-mitochondrial supernatant PMS, crude and gradient-purified mitochondria; n = 4) generated during pure/crude experiments were analyzed separately by LC-MS following tryptic in-solution digestion. To absolutely quantify the yeast proteome and determine protein copy numbers, cell lysates were prepared from light, medium-heavy, and heavy SILAC-labeled, mixed in 1:1:1 ratios, and analyzed by LC-MS in three biological replicates including a label switch (see Figure S1B).

For medium-scale preparations of mitochondria from up to 20 yeast strains in parallel, performed for validation experiments, the protocol was as follows: *S. cerevisiae* strains were grown to mid-logarithmic phase in 360 mL of YPG medium in a 1-L flask. 400 OD₆₀₀ of cells were pelleted by centrifugation (3,000 x g, 5 min, 25°C, SLA-300 rotor [Sorvall™, Thermo Fisher Scientific Inc.]). Yeast pellets were washed with dH₂O in 50 mL conical tubes, suspended in 3.5 mL of DTT buffer and transferred to 5-mL Eppendorf Tubes® (Eppendorf AG). Samples were incubated at 30°C and 900 rpm for 20 min (ThermoMixer® C, Eppendorf AG) and collected by centrifugation at 3,000 x g for 5 min at 25°C (Rotor FA-45-20-17, Centrifuge 5804 R [Eppendorf AG]). Pellets were dissolved in 4.5 mL of zymolyase buffer, 30 mg Zymolyase®-20T (Nacalai Tesque Inc.) were added to each sample and samples were incubated for 45 min at 30°C. Spheroplasts were pelleted by centrifugation (1,500 x g, 5 min, 25°C), washed with 3.5 mL of 1.2 M sorbitol and dissolved in 4.5 mL homogenization buffer containing 0.2% (w/v) bovine serum albumin. Cells were opened on ice using an Omnifix® 10 mL LL syringe (B. Braun Melsungen AG, 4617100V) and a Sterican® 0.90 x 0.40 mm cannula (B. Braun Melsungen AG, 4657519) by 15-20 repetitions of drawing up the dissolved yeast cells through the cannula into the syringe and pushing the cells back to the 5 mL Eppendorf tube. Cell debris was removed by centrifugation (1,500 x g, 5 min, 4°C) and supernatant was transferred to a fresh 5 mL Eppendorf tube and subjected to two consecutive centrifugation steps at 3,000 x g for 5 min at 4°C. Afterwards mitochondria were pelleted by centrifugation at 20,913 x g for 5 min at 4°C. Mitochondrial pellet was suspended in 3.5 mL SEM buffer and subjected to a low-speed centrifugation (3,000 x g for 5 min at 4°C). Supernatant was transferred to a fresh 5 mL Eppendorf cup and mitochondria were collected by centrifugation at 20,913 x g for 5 min

at 4°C. Mitochondria were dissolved in 200 µL of SEM buffer and protein concentration was determined using the Bradford assay. Afterwards, mitochondria were aliquoted, snap frozen in liquid nitrogen and stored at -80°C.

Subcellular fractionation for biochemical validation of subcellular protein localization

70 OD₆₀₀ of cells (corresponds to 70 mL of cultured cells grown to an OD₆₀₀ of 1.0) were harvested by centrifugation at 3,000 x g for 5 min at 25°C. Cells were suspended in 2 mL DTT buffer and incubated for 20 min at 30°C.

Afterwards, yeast cells were collected by centrifugation (3,000 x g, 5 min, 25°C) and suspended in 1 mL of zymolyase buffer containing 8 mg/mL Zymolyase[®]-20T (Nacalai Tesque Inc.). Spheroplasts were pelleted by centrifugation (1,500 x g, 5 min, 25°C) and suspended in 2 mL of homogenization buffer. Cells were opened by 20 strokes with a PTFE pestle (Sartorius, BBI-8542708) in a 5 mL Homogenizer Vessel (Sartorius, BBI-8542309) on ice. Cell debris, nuclei and intact cells were pelleted (1,500 x g, 5 min, 4°C) and supernatant (containing crude mitochondria) was subjected to another 10 strokes. Half of the post-nuclear supernatant (PNS) was precipitated with trichloroacetic acid (TCA) and dissolved in 2x SDS sample (120 mM Tris-HCl [pH 6.8], 4% [w/v] SDS, 20% [v/v] glycerol, 0.02% [w/v] bromphenolblue) supplemented with 10 mM DTT and 2 mM of PMSF.

Mitochondria were isolated from the other half of the PNS by centrifugation at 13,000 x g for 10 min at 4°C. To remove loosely attached proteins of other cellular compartments from the surface of mitochondria, mitochondria were suspended in 200 µL of SEM buffer, loaded onto 500 µL of S₅₀₀EM (500 mM sucrose, 1 mM EDTA, 10 mM MOPS-KOH [pH 7.2]) and re-isolated by centrifugation (13,000 x g for 10 min at 4°C). The mitochondrial pellet (Mito) was dissolved in 200 µL of 2x SDS sample buffer supplemented with 10 mM DTT and 2 mM of PMSF. For isolation of microsomal fraction, the supernatant was subjected to ultracentrifugation at 100,000 x g for 1 h at 4°C. The microsomal pellet (P100) was suspended in 200 µL of 2x SDS sample buffer supplemented with 10 mM of DTT and 2 mM of PMSF whereas the supernatant (S100) was treated with trichloroacetic acid before addition of SDS sample buffer. 20 µL of PNS, 20 µL of Mito, 23 µL of P100 and 23 µL of S100 fractions were subjected to SDS-PAGE.

Protease accessibility assay

For the global analysis of the suborganellar localization of mitochondrial proteins by quantitative MS, gradient-purified mitochondria (100 µg per replicate, 5 mg/mL) obtained from heavy SILAC-labeled (replicates 1 and 3) or unlabeled cells (replicate 2) were treated with (i) protease only (referred to as S1), (ii) protease following rupture of

the outer membrane (OM) using digitonin (S2), and (iii) protease following lysis of mitochondria using Triton X-100 (S3). Same amounts of mitochondria from differentially SILAC-labeled cells (replicates 1 and 3, unlabeled; replicate 2, heavy labeled) remained untreated and served as control (M). For the perforation of OM, mitochondria were incubated with digitonin (0.1% [w/v] final concentration) for 3 min at 4°C. Membrane perforation was stopped by adding the 14-fold volume of SEM buffer. To lyse mitochondria, Triton X-100 was added to a final concentration of 1% (w/v) and samples were incubated for 10 min at 4°C. Volumes of all samples were adjusted with SEM buffer to the volume of S2. For the digestion of accessible proteins in S1, S2, and S3, trypsin and proteinase K were added (final concentration of 5 µg/mL each) and samples were incubated for 15 min at 4°C. Proteases were inactivated by adding PMSF (1 mM final concentration) and incubation for 15 min at 4°C. Mitochondria (M, S1) and mitoplasts (S2) were collected by centrifugation (10 min, 12,000 x g, 4°C), layered with SEM buffer containing 1 mM PMSF to inactivate residual protease activity, and recollected by centrifugation (10 min, 12,000 x g, 4°C). Pellets were resuspended in 100 µL SDS sample buffer containing 50 mM DTT and 1 mM PMSF and boiled for 20 min at 94°C. Proteins in S3 were precipitated by adding TCA (15% [w/v], final concentration) followed by incubation for 30 min at 4°C and centrifugation (15 min, 14,000 x g, 4°C). The resulting pellets were washed with 80% ice-cold acetone (v/v) and centrifuged again (15 min, 14,000 x g, 4°C). Supernatants were carefully removed and the pellets were resuspended in 100 µL SDS sample buffer containing 50 mM DTT and 1 mM PMSF and boiled for 20 min at 94°C. S1, S2, and S3 were each mixed with equal amounts of M for subsequent quantitative MS analysis (see Figure S1C). This experiment is referred to as 'submitochondrial profiling experiment' in this work.

Swelling assay for determination of submitochondrial protein localization

Crude mitochondria were suspended in SEM buffer and split into samples of equal volume (each sample containing 50-70 µg of mitochondria). Half of the samples were subjected to hypoosmotic swelling by addition of 8 volumes of EM buffer. The other half of the samples were diluted with the same volume of SEM buffer (mitochondria were left intact). Where indicated, samples were treated with proteinase K (final concentration: 7 µg/mL) for 15 min on ice. PMSF was added to all samples to a final concentration of 2.5 mM. Mitochondria were washed with SEM buffer and dissolved in 2x SDS sample buffer supplemented with 2% (v/v) of β-mercaptoethanol and 2 mM of PMSF. Samples were analyzed by SDS-PAGE and immunoblotting. To control for protein aggregates, 70 µg of mitochondria were suspended in 50 µL of 1% Triton X-100 (diluted in SEM buffer) and incubated on ice for 7 minutes. Where indicated, samples were subjected to proteinase K treatment for 15 min at a final concentration of 7 µg/mL. Protease

was inhibited by the addition of PMSF to a final concentration of 2 mM. After addition of SDS sample buffer, samples were separated by SDS-PAGE and analyzed by immunoblotting.

Purification of mitochondrial protein complexes using IgG and HA affinity chromatography

Two milligram of wild-type and Protein A-tagged mitochondria, respectively, were suspended in 2 mL of solubilization buffer (1% [w/v] digitonin, 20 mM Tris-HCl [pH 7.4], 50 mM NaCl, 10% glycerol, 0.1 mM EDTA) supplemented with 1.5 mM PMSF and 1x cOmplete™, EDTA-free Protease Inhibitor Cocktail (F. Hoffmann-La Roche AG). Mitochondria were incubated end-over-end for 30 min at 4°C. Non-solubilized material was removed by centrifugation. Afterwards, 50 µL of solubilized mitochondria were mixed with 12.5 µL of 4x SDS sample buffer containing 4% (v/v) β-mercaptoethanol and 4 mM PMSF (Load fraction).

For co-immunoprecipitation of Protein A-tagged mitochondrial proteins, 150 µL of 50% slurry human IgG-coupled Sepharose beads were equilibrated with solubilization buffer, mixed with solubilized mitochondria and incubated end-over-end for 2 h at 4°C. After binding of Protein A-tagged proteins to IgG, beads were re-collected by centrifugation (100 x g, 1 min, 4°C) and transferred to Mobicol Mini-Columns. To remove unspecifically-bound proteins beads were washed 12 times with 500 µL of wash buffer (0.1-0.3% [w/v] digitonin, 20 mM Tris-HCl [pH 7.4], 60 mM NaCl, 10% glycerol, 0.5 mM EDTA) supplemented with 1.5 mM PMSF and 1x cOmplete™, EDTA-free Protease Inhibitor Cocktail (F. Hoffmann-La Roche AG). Specifically bound proteins were eluted by the addition of 10 µL of AcTEV Protease (Thermo Fisher Scientific Inc.) in 150 µL of wash buffer, followed by vigorous shaking at 4°C for 16 h. For removal of His-tagged AcTEV protease from elution mixture, 10 µL of Ni-NTA agarose (Qiagen N.V.) were equilibrated with wash buffer and added to eluate, followed by vigorous shaking for 30 min at 4°C. Eluted mitochondrial proteins were collected by centrifugation (200 x g, 1 min, 4°C). To remove residual proteins, another 50 µL of wash buffer was added to the IgG-coupled Sepharose beads and collected by centrifugation yielding a total of 215 µL of eluate. 115 µL of eluate were mixed with 30 µL of 4x SDS sample buffer. For the analysis of protein complexes by SILAC-based quantitative MS, 100 µL of eluates obtained from differentially labeled wild-type cells and cells expressing the Protein A-tagged bait were mixed and analyzed by LC-MS.

For co-immunoprecipitation from _{HA}Tmh11 and corresponding wild-type mitochondria one milligram of mitochondria were used per strain. They were suspended in 1 mL of solubilization buffer (supplemented with 1 mM PMSF) and incubated end-over-end for 30 min at 4°C. After removal of non-solubilized material by centrifugation,

20 μL of supernatant were mixed with 20 μL of 2x SDS sample buffer containing 2% (v/v) β -mercaptoethanol and 2 mM PMSF (Load fraction). HA-tagged proteins were bound to 100 μL of pre-equilibrated anti-HA Affinity Matrix slurry (F. Hoffmann-La Roche AG, 11815016001) by incubation end-over-end for 1 h at 4°C. Samples were washed 10 times with 600 μL of HA wash buffer (0.1% [w/v] digitonin, 20 mM Tris-HCl [pH 7.4], 50 mM NaCl, 10% glycerol, 0.1 mM EDTA, 1 mM PMSF) and proteins were eluted in 200 μL of 1x SDS sample buffer and afterwards supplemented with 1% (v/v) of β -mercaptoethanol.

SDS-PAGE and tryptic in-gel digestion

Mixtures of crude mitochondrial fractions and gradient-purified mitochondria obtained in pure/crude experiments ($n = 4$) were boiled in SDS sample buffer for 5 min at 94°C. Proteins (25 μg each per gel lane) were separated on 4 - 12% NuPAGE™ Bis-Tris and Novex™ 16% Tricine gels (Thermo Fisher Scientific) according to the manufacturer's protocol. Following visualization of proteins using colloidal Coomassie Brilliant Blue, gel lanes were cut into 18 slices of equal size. Slices were washed and destained by alternatingly incubating them with 10 mM NH_4HCO_3 and 50% (v/v) acetonitrile (ACN)/10 mM NH_4HCO_3 (10 min at RT each). Cysteine residues were reduced (10 mM DTT/10 mM NH_4HCO_3 , 30 min at 56°C) and alkylated (50 mM iodoacetamide/10 mM NH_4HCO_3 ; 30 min at RT in the dark) followed by proteolytic digestion of proteins using trypsin (60 ng per slice; overnight at 37°C). Peptides were eluted with 0.05% (v/v) trifluoroacetic acid (TFA)/50% (v/v) ACN, dried *in vacuo* and resuspended in 15 μL 0.1% TFA prior to LC-MS analysis. Samples obtained in submitochondrial profiling experiments ($n = 3$; 25 μg of mixed treated and untreated mitochondria per gel lane) were separated using 4 - 12% NuPAGE Bis-Tris gradient gels and processed as described above except that lanes were cut into 24 slices each.

Proteolytic in-solution digestion

Proteins of pure/crude, subcellular profiling, and q-AP-MS experiments were acetone-precipitated and resuspended in 8 M urea/50 mM NH_4HCO_3 . Samples of the absolute quantification experiment were adjusted to 8 M urea and 50 mM NH_4HCO_3 by directly adding the required amounts of the chemicals to the samples. Cysteine residues were reduced with 5 mM Tris(2-carboxyethyl)phosphine (30 min, 37°C) and free thiol groups were subsequently alkylated with 50 mM iodoacetamide/50 mM NH_4HCO_3 (30 min at RT in the dark). The alkylation reaction was quenched by adding DTT to a final concentration of 25 mM. For proteolytic digestion, urea concentration was adjusted to 4 M (LysC), 1.6 M (trypsin), or 1 M (AspN, chymotrypsin, GluC) using 50 mM NH_4HCO_3 . Proteases were added at a

protease-to-protein ratio of 1:50 for trypsin, 1:100 for chymotrypsin, LysC, and GluC, or 1:150 for AspN. Proteins were digested overnight at 37°C. In case of LysC/trypsin double-digestion, incubation with LysC was performed for 4 h (37°C) followed by incubation with trypsin overnight (37°C). Proteolysis was stopped by acidifying the samples with 100% TFA. For the direct analysis (i.e. without further sample fractionation, referred to as '1-shot' analysis) of peptides from pure/crude and subcellular profiling experiments, 10 µg of protein were digested. Acidified peptides were cleared by centrifugation (5 min, 12,000 x g, RT) and one-fifth of each digest was analyzed by LC-MS. For the analysis of peptide samples that were further fractionated by strong cation exchange chromatography (SCX) or high pH reversed-phase (RP) chromatography, 300 µg of protein were digested and peptides were desalted using C18 cartridges (3M Empore, St. Paul, USA) according to the manufacturer's protocol and dried *in vacuo*.

Strong cation exchange chromatography

Dried peptides of tryptic, chymotryptic, AspN, LysC, and GluC digests of differentially SILAC-labeled, mixed crude and gradient-purified mitochondria from pure/crude experiments (n = 4) were resuspended in 500 µL SCX buffer (10 mM KH₂PO₄ [pH 3.0], 25% [v/v] ACN) and loaded onto cation exchange mini-columns (POROS™ 50 HS strong cation exchange resin; 4 x 15 mm; particle size, 50 µm; AB Sciex) equilibrated with SCX buffer. Peptides were eluted step-wise with 35, 55, 75, 100, 125, 150, 200, 250 and 350 mM KCl in SCX buffer (500 µL each). The resulting nine fractions were lyophilized. Peptides were resuspended in 500 µL 0.5% (v/v) acetic acid, desalted using StageTips, dried *in vacuo*, and resuspended in 60 µL of 0.1% TFA, of which 15 µL were analyzed by LC-MS.

High pH reversed-phase chromatography

High pH RP chromatography (Delmotte et al., 2007; Lasaosa et al., 2009) was used for the fractionation of peptides derived from pure/crude samples and from samples of the absolute quantification experiment. Tryptic peptides of mitochondrial fractions were resuspended in 99% solvent A (72 mM triethylamine, 52 mM acetic acid, pH 10) and 1% solvent B (72 mM triethylamine and 52 mM acetic acid in ACN) and loaded onto a Gemini-NX column (150 mm x 2 mm inner diameter, particle size 3 µm, pore size 110 Å; Phenomenex, Aschaffenburg, Germany) using an Ultimate 3000 HPLC system (Thermo Fisher Scientific, Dreieich, Germany) at a flow rate of 200 µL/min and a column temperature of 40°C. For peptide separation, a gradient of 1 - 55% solvent B (starting after 5 min) in 55 min followed by 55 - 70% B in 2 min and 2 min at 70% B was used. Fractions were collected in 30 s intervals starting at minute 8.5 and ending at minute 56.5 and pooled into 30 non-contiguously concatenated fractions by combining

every 30st fraction. Peptides of mixed cell lysates, generated by a LysC/trypsin double digestion, were resuspended in 99% solvent A' (10 mM NH₄OH, pH 10) and 1% solvent B' (10 mM NH₄OH in 90% [v/v] ACN) and loaded onto a Gemini-NX column as described above. Peptides were eluted by applying a gradient of 1 - 54% solvent B' (starting after 5 min) in 53 min followed by 54 - 78% B' in 5 min, 5 min at 78% B', and 78 - 1% B' in 7 min. Fractions were collected in 40 s intervals between minute 1 and minute 73 in a non-contiguous, concatenated way resulting in a total of 12 fractions. Peptides of all high pH RP chromatography experiments were lyophilized, washed with 70% (v/v) ACN, dried again *in vacuo*, and resuspended in 60 µL (pure/crude experiments) or 150 µL (absolute quantification experiments) of 0.1% TFA, of which 15 µL were analyzed by LC-MS.

LC-MS analysis

Peptide mixtures were analyzed by nano-HPLC-ESI-MS/MS using an LTQ Orbitrap XL, an Orbitrap Elite, or a Q Exactive instrument (Thermo Fisher Scientific, Bremen, Germany) each directly coupled to an UltiMate 3000 RSLCnano HPLC system (Thermo Fisher Scientific, Dreieich, Germany). Peptides were washed and pre-concentrated on PepMap™ C18 precolumns (5 mm x 300 µm inner diameter; Thermo Scientific) and separated using AcclaimPepMap™ RSLC columns (50 cm x 75 µm inner diameter; pore size, 100 Å; particle size, 2 µm) at a flow rate of 250 nl/min and 40 - 43°C. For peptide elution, binary solvent systems were used consisting of (i) 0.1% (v/v) formic acid (FA)/4% (v/v) DMSO (solvent A) and 0.1% (v/v) FA/4% (v/v) DMSO/48% (v/v) methanol/30% (v/v) ACN (solvent B) for LC-MS analyses using the LTQ Orbitrap XL and Orbitrap Elite or (ii) 0.1% (v/v) FA (solvent A) and 0.1% (v/v) FA/86% (v/v) ACN (solvent B) for measurements using the Q Exactive. Length and slope of the gradients were adjusted according to the complexity of individual samples. Peptide mixtures of pure/crude experiments, obtained following various protein and peptide fractionation methods, were analyzed at the Orbitrap Elite using the LC gradients detailed in the following. Tryptic peptides, analyzed without fractionation, were eluted with a gradient of 5 - 25% solvent B in 115 min followed by 25 - 45% in 110 min, 45 - 60% in 50 min, 60 - 80% in 20 min, 80 - 99% in 10 min, and 5 min at 99%. Peptides derived from tryptic in-gel digests were separated with 1 - 65% solvent B in 50 min, 65 - 95% in 5 min and 5 min at 95%. For the separation of peptides collected by high pH RP chromatography, a gradient ranging from 1 - 67.5% solvent B in 50 min, 67.5 - 95% in 5 min, and 5 min at 95% was applied, and peptides of SCX samples were eluted with 1 - 40% solvent B in 55 min followed by 40 - 70% in 30 min, 70 - 99% in 10 min, and 5 min at 99%. Separation of tryptic peptides of absolute quantification experiments, analyzed at the Q Exactive, was performed using a gradient of 4 - 39% solvent B' in 195 min, 39 - 54% in 15 min, 54

- 95% in 5 min, and 5 min at 95%'. Tryptic peptides of subcellular profiling experiments, analyzed without fractionation at the Orbitrap Elite, were separated by applying a gradient ranging from 1 - 25% solvent B in 115 min, 25 - 45% in 110 min, 45 - 60% in 50 min, 60 - 80% in 20 min, 80 - 99% in 10 min, and 5 min at 99%. Tryptic peptides derived from submitochondrial profiling experiments were eluted with a gradient ranging from 1 - 65% solvent B in 70 min, 65 - 95% in 5 min, and 5 min 95% followed by MS analysis at the Orbitrap Elite (replicates 1 and 2) or with 4 - 40% solvent B' in 50 min, 40 - 95% in 5 min and 5 min at 95% when analyzed at the Q Exactive (replicate 3). For the separation of peptides from q-AP-MS experiments, a gradient of 1 - 30% solvent B in 65 min, 30 - 45% in 30 min, 45 - 70% in 25 min, 70 - 99% in 15 min, and 5 min at 99% was applied followed by MS analysis at the Orbitrap Elite (Aim11, Coq21, Iai11, Rci37 and Rcf3 complexes) or the LTQ Orbitrap XL (Mco10 and Rci50 complexes). Replicate 1 of Coq21, Rci37, and Rcf3 complexes were analyzed in two technical replicates.

Mass spectrometers were equipped with a Nanospray Flex ion source with DirectJunction (Thermo Scientific; Q Exactive and Orbitrap Elite) or a Finnigan Nanospray ion source with dynamic NSI probe (Thermo Scientific; LTQ-Orbitrap XL) and stainless steel (Thermo Scientific) or fused silica emitters (New Objective, Woburn, USA). MS instruments were externally calibrated using standard compounds. The Orbitrap Elite was operated with the following mass spectrometric parameters: MS survey scans ranging from m/z 370 - 1,700 were acquired in the orbitrap at a resolution (R) of 120,000 (at m/z 400). Automatic gain control (AGC) was set to 1×10^6 ions and the maximum injection time (IT) to 200 ms. A TOP15 (samples of pure/crude experiments fractionated by SDS-PAGE), TOP20 (pure/crude samples fractionated by high pH RP and samples of submitochondrial profiling experiments), or TOP25 method (all other samples) was used for low energy collision-induced dissociation (CID) of multiply charged precursor peptides in the linear ion trap applying a normalized collision energy (NCE) of 35%, an activation q of 0.25, an activation time of 10 ms, an AGC of 5×10^3 , and a max. IT of 150 ms. Singly charged precursor peptides were generally rejected from fragmentation, except for the analysis of samples derived from SCX experiments, which were additionally analyzed allowing fragmentation of singly charged precursor peptides to increase the likelihood to identify small mitochondrial proteins. The dynamic exclusion time (DE) for previously fragmented precursors was set to 45 s. For analyses at the Q Exactive, the following parameters were applied: MS scans, m/z 375 - 1,700; R, 70,000 (at m/z 200); AGC, 3×10^6 ions; max. IT, 60 ms; TOP12 (samples of submitochondrial profiling experiments) or TOP15 method (samples of absolute quantification experiments) for higher energy collisional dissociation (HCD) of precursor peptides ($z \geq 2$) in the orbitrap; NCE, 28%; AGC, 1×10^5 ions, max. IT, 120 ms; DE, 45 s. LTQ Orbitrap XL parameters were as follows: MS scans, m/z 370 - 1,700; R, 60,000 (at m/z 400); AGC, 5

$\times 10^5$ ions; max. IT, 500 ms; TOP5 CID method for fragmentation of precursor peptides ($z \geq 2$); NCE, 35%; activation q, 0.25; activation time, 30 ms; AGC, 1×10^4 ions, max. IT, 100 ms; DE, 45 s.

MS data analysis

For peptide and protein identification and quantification, mass spectrometric raw data were processed using the MaxQuant software package (version 1.5.3.12) with its integrated search engine Andromeda (Cox and Mann, 2008; Cox et al., 2011). MS/MS data of different experiments (i.e., pure/crude, absolute quantification, subcellular and submitochondrial profiling, and q-AP-MS experiments) were searched separately against a custom-made protein database containing all entries of the *Saccharomyces* Genome Database (SGD; <http://www.yeastgenome.org/>, downloaded 02/04/2016; 6,726 entries including protein sequences of verified, uncharacterized, and dubious open reading frames [ORFs]) and 47 entries for putative small ORFs reported by Smith et al. (2014). Data were also correlated with a list of common contaminants provided by MaxQuant. Database searches were performed with mass tolerances of 4.5 ppm for precursor and 0.5 Da (CID data) or 20 ppm (HCD data) for fragment ions and the appropriate enzymatic specificity allowing two (AspN), three (trypsin/P, LysC), or four missed cleavages (chymotrypsin, GluC). The cleavage sites for chymotrypsin were defined as C-terminal to tyrosine, phenylalanine, tryptophan, leucine, and methionine. Acetylation of protein N-termini and oxidation of methionine were considered as variable and carbamidomethylation of cysteine residues as fixed modification. Arg10 and Lys8 were set as heavy labels and Arg6/Lys4 as medium-heavy labels. The options 'requantify', 'match between runs' (match from and to), and 'iBAQ' were generally enabled except for the analysis of data from subcellular profiling experiments ('requantify' disabled) and submitochondrial profiling experiments ('match between runs' and 'iBAQ' disabled). For the analysis of data from submitochondrial profiling experiments, which were analyzed following a gel-based approach, each gel slice ($n = 24$ per replicate; 3 replicates) was defined as individual 'experiment' in the experimental design template. Proteins were identified based on at least one unique peptide with a length of six amino acids and a maximum mass of 4,600 Da or 6,000 Da (for data from pure/crude experiments only). Lists of peptides and proteins identified by MaxQuant were filtered applying a false discovery rate of 0.01. Proteins identified by the same set of peptides were combined to a single protein group by MaxQuant. For SILAC-based protein quantification, at least one unique peptide and a minimum ratio count of one were required. Lists of all proteins identified and quantified in individual experiments are provided in Table S2A (subcellular profiling), S2B and S2C (pure/crude data), S2D and S2E

(absolute proteome quantification under fermentable and non-fermentable growth conditions), S2F - S2H (submitochondrial profiling), and S2I (q-AP-MS data).

Mean protein abundance ratios for pure/crude experiments ($n = 4$) were calculated based on normalized peptide SILAC ratios determined by MaxQuant. For each replicate, the logarithmic pure/crude ratio (normalized) for a protein was computed as the median of \log_2 peptide ratios across values obtained for all different fractionation techniques. Ratios for two of the four replicates were inverted to account for the label switch. MaxQuant additionally reports non-normalized peptide ratios. In cases when the major population of proteins remains unchanged in abundance, the normalization may compensate for slight differences in the mixing of differentially SILAC-labeled samples by shifting the values to a median logarithmic ratio of zero per experiment. In the pure/crude experiment, however, the distribution of logarithmic ratios is not unimodal and includes a major population reflecting proteins of higher abundance in crude mitochondrial fractions compared to gradient-purified mitochondria with \log pure/crude ratios < 0 (see '*Classification of proteins of the pure/crude dataset*' in 'Bioinformatics and statistics'). To take into account the bimodal distribution of the values, \log pure/crude protein ratios based on normalized peptide ratios were subsequently shifted back by calculating the mean difference between normalized and non-normalized protein ratios for each replicate and subtracting the mean of that differences determined across all replicates.

For visualization of pure/crude data, mean \log_2 ratios (pure/crude) were plotted against mean \log_2 iBAQ values. Only proteins that were quantified in all four biological replicates with at least two fractionation techniques were considered for subsequent statistical and bioinformatics data analyses.

Protein copy numbers for cells grown on galactose, glycerin, or glucose were determined based on the MS intensities for light, medium-heavy, and heavy SILAC-labeled proteins according to the 'Proteomic Ruler' strategy (Wiśniewski et al., 2014) using Perseus (Tyanova et al., 2016) with the 'proteomic ruler' plugin.

The analysis of data obtained in subcellular profiling experiments was based on MS intensities determined for proteins identified in cell lysates, the non-mitochondrial fraction PMS, crude and gradient-purified mitochondria. MS intensities of proteins in distinct subcellular fractions were normalized to the total intensity per fraction. For each protein and fraction, the mean of the normalized intensities across all replicates ($n = 2 - 4$) was determined. For visualization, the value of the subcellular fraction with the highest mean normalized intensity was set to one (see Figures 1B, 4E, and S5).

The setup of the MaxQuant analysis of data derived from submitochondrial profiling experiments enabled the quantification of each protein in each of the 24 gel slices per replicate. For each protein, the slice with the highest intensity for M (i.e., untreated gradient-purified mitochondria, serving as control) was determined and the SILAC

ratio for this slice was used for further data processing. In case a protein was missing from treated mitochondria in S1, S2, and S3 due to digestion by the proteases added to the sample and, thus, an S/M protein ratio could not be calculated for a slice, the ratio for this protein was set to 0.01. Mean SILAC ratios across all replicates ($n = 2 - 3$) were determined. For this, only ratios from adjacent gel slices were taken into account. For further bioinformatics analyses, proteins were required to exhibit ratios for all three experimental conditions in at least two replicates. Data used to analyze the topology of membrane proteins were derived from the MaxQuant peptides.txt results file. Putative membrane proteins were determined based on TMHMM predictions (information was retrieved from the SGD). For each peptide of these proteins, the intensities for each S1, S2, S3, and M across all gel slices were summed up and mean S/M intensity ratios per replicate were determined. In case a peptide intensity was given for M but missing for S, the S/M ratio was set to 0.01. For peptides quantified in at least two replicates, mean S/M intensity ratios across replicates were calculated. In peptide plots, the highest value was set to 1.3 (see Figures 5E and 5F).

Bioinformatics and statistics

Gene Ontology annotations. GO annotations for the domains 'biological process' (BP), 'cellular component' (CC) and 'molecular function' (MF) including child terms were downloaded from the SGD (date of download: 12/08/2016). To compile a list of proteins annotated as mitochondrial in the SGD, all entries of the database were searched for the GOCC term 'mitochondrion' (GO:0005739) including its child terms. The resulting list contained 1,178 proteins. To obtain information about mitochondrial subproteomes, we retrieved proteins annotated as OM (GO:0005741; 107 proteins), intermembrane space (IMS; GO:0005758; 61), inner membrane (IM; GO:0005743; 253), and matrix proteins (GO:0005759; 226). To compare the protein composition of crude mitochondrial fractions (Mito) and gradient-purified mitochondria (pMito), proteins identified in pure/crude experiments were assigned to a distinct subcellular localization based on GOCC terms (see Fig. 2B, Table S2B). Intensity-based absolute quantification (iBAQ) values, calculated by MaxQuant, were used as a measure for the abundance of proteins in crude and pure mitochondria. For each protein and replicate ($n = 4$), iBAQ values determined for different fractionation strategies were averaged and the mean across the replicates was calculated for pure and crude mitochondria. Mean iBAQ values of all proteins assigned to a distinct subcellular localization were summed up and the percentage of the overall iBAQ value in crude or pure mitochondria was determined.

Classification of proteins of the pure/crude dataset. Proteins quantified with at least two different fractionation techniques in all four replicates of pure/crude experiments (i.e., 3,365 proteins) were classified based on their

pure/crude SILAC ratios (\log_2 values). To test for multimodal distribution of the pure/crude data, the 'Hartigan's dip test' was performed (using the 'bimodalitytest' package for R) and revealed two distributions with centers at mean \log_2 (pure/crude) of -1.47 for distribution 1 (d1) and 0.31 for d2 (see Fig. S2H). The standard deviation was 0.83 for d1 and 0.08 for d2. For each distribution, 1,000,000 theoretical values were generated taking into account the respective standard deviations. We next determined for each protein present in the dataset to which of the two distributions it belongs using a two-sample two-sided equivalence test (R package 'equivalence') with a significance threshold (p value) of < 0.01 . As a result, 544 proteins were assigned to d1 and 812 proteins to d2 (see Table S2B). The remaining proteins were classified based on a two-sample two-sided t test (p value < 0.01 ; R package 'stats'). 34 of these proteins were determined to be outliers of d1 only and, thus, assigned to d2. *Vice versa*, 1,515 proteins were outliers of d2 only and assigned to d1. 460 proteins were outliers of both d1 and d2. These proteins were distributed at ratios $< d1$ (referred to as OL1), between d1 and d2 (OL2), and at ratios $> d2$ (OL3). Based on this classification, four distinct classes were defined for the pure/crude dataset: proteins of d2 and OL3 were combined to class 1 and proteins of OL2, d1, and OL1 are defined as class 2, class 3, and class 4, respectively. Proteins of class 1 were further filtered based on sequence coverage and standard deviation calculated for mean \log_2 (pure/crude) abundance ratios across all replicates, i.e. proteins with a sequence coverage of $< 20\%$ and a standard deviation of > 0.75 were removed (i.e., 113 proteins).

Clustering analysis. Data obtained in absolute quantification and submitochondrial profiling experiments (see Figs. 3 and 5) were subjected to *k*-means clustering to group proteins showing similar characteristics and low variance in the respective experiment. For absolute quantification experiments, clustering was based on absolute protein copy numbers determined for yeast grown on galactose (Gal), glycerin (Gly), or glucose (Glc). Only proteins for which copy numbers were determined for all carbon sources in all replicates ($n = 3$) were considered. For each protein, ratios of mean \log_2 copy numbers Gal/Glc and Gly/Glc were calculated. To reveal proteins with significant carbon source-dependent changes in copy numbers, an analysis of variance (ANOVA) test was performed between the Gal/Glc and Gly/Glc data (R package 'stats'). 1,576 proteins showed significant differences (p value ≥ 0.05) in protein copy numbers and were used for the *k*-means clustering analysis ('stats' package). The Davies-Bouldin index was used to determine the optimal number of clusters for this dataset (i.e., $k = 14$). Information about the proteins present in the clusters are provided in Table S2D. The clustering analysis of data from submitochondrial profiling experiments was based on SILAC ratios determined in protease accessibility assays for proteins of gradient-purified mitochondria treated with proteases (S1), digitonin and proteases (S2), or Triton X-100 and proteases (S3) versus

untreated mitochondria (M). For clustering, only proteins of class 1 as determined in pure/crude experiments that had S/M ratios for all three experimental conditions (i.e., S1 - S3) in at least two replicates (n = 3) were considered.

Furthermore, proteins were required to (i) exhibit mean S/M ratios ≤ 1.1 (since due to the experimental design S/M ratios should not exceed a value of 1) and (ii) be decreased in abundance with increasing protease-accessibility of proteins ($S3 > S2 > S1$) with a tolerance of ± 0.2 . A total of 624 proteins met all criteria and were subjected to *k*-means clustering into 5 clusters, which was determined to be the optimal cluster number for this dataset. Information about the proteins present in the clusters are provided in Table S2F. Clusters defined for both absolute quantification and submitochondrial profiling experiments were further clustered by hierarchical clustering using the mean ratio of each cluster to reveal (dis)similarities between individual clusters ('stats' package).

Gene Ontology overrepresentation analysis. GO term overrepresentation analyses were based on GO terms retrieved from the SGD and were performed with an in-house developed script using all proteins quantified in the respective dataset as background. The two-sided Fisher's exact test was used to calculate mid p values. Raw mid p values were corrected for multiple testing by Benjamini-Hochberg adjustment and GO terms with a corrected mid p value of < 0.05 were considered enriched.

Principal component analysis. Proteins present in the OM, IMS/IM, and matrix/IMS clusters of the submitochondrial profiling experiment (see Fig. 5C) and further proteins meeting the criteria for signature plots (i.e. $S1/M < 0.25$ for OM proteins; $S1/M > 0.25$ and $S2/M < 0.25$ IMS/IM proteins; $S1/M$ and $S2/M > 0.25$, $S3/M < 0.25$ for matrix/IM proteins) were subjected to principal component analysis. To this end, $S1/M$, $S2/M$, and $S3/M$ protein ratios were \log_2 -transformed, analyzed using the R package 'stats' with the parameters 'center' and 'scale.' set to 'true', and visualized in two-dimensional scores plots depicting the two principal components providing the best visual separation of protein clusters on the x- and y-axes.

Further bioinformatics tools. An in-house developed software based on R was used to process MaxQuant result files, visualize data, and subsequently analyze the data using statistical and bioinformatics means as described. The following R packages were used: base, bimodalitytest, Biobase, clusterSim, clustpro, convert, curl, data.table, datasets, Deducer, devtools, diptest, dplyr, equivalence, flux, gage, GGally, ggbiplot, ggplot2, gplots, graphics, grDevices, grid, gtools, Hmisc, lattice, methods, outliers, pastecs, plotrix, reshape, reshape2, ROCR, scales, silvermantest, stats, stringr, utils, VennDiagram, Vennerable and xtable. Prediction of N-terminal mitochondrial targeting sequences was performed using MitoFates (Fukasawa et al., 2015). For sequence homology searches the HMMER web server was employed (Fin et al., 2015).

Subcellular *in vivo* localization of uncharacterized GFP-tagged proteins

C-terminally GFP-tagged yeast strains were grown over night in 50 mL YPD medium at 30°C to an OD₆₀₀ of 1.0. Subsequently, 2 OD₆₀₀ of yeast cells were harvested, washed with dH₂O and suspended in 500 µL dH₂O. 3 µL of yeast cell solution was dispersed on a microscope slide (Diagonal), covered with a microscope cover glass (Diagonal) and samples were analyzed immediately. Fluorescence microscopy was performed using the Olympus BX61 microscope (Olympus K.K.) equipped with the immersion oil objective UPLFLN 100x/1.3 (Olympus K.K.). Visualization of GFP was carried out using the 470/40 nm bandpass excitation filter, a 495 nm dichromatic mirror and a 525/50 nm bandpass emission filter. All digital recordings of cells were taken with the F-view CCD camera (Olympus Soft-Imaging Solutions GmbH) controlled by the Cell-P software (Olympus K.K.).

Strains expressing N-terminally GFP-tagged mitochondrial proteins under the control of their native or the *spNOP1* promoter were grown in synthetic defined (SD) complete or SD -URA medium. Imaging of the strains was performed using a wide-field epi-fluorescent Olympus microscope with a 60X air objective (excitation, 490/20 nm; emission, 535/50 nm).

***In vitro* import of radiolabeled precursor proteins into mitochondria**

[³⁵S]-labeled precursor proteins were synthesized using the TnT[®] Quick Coupled Transcription/Translation System (Promega GmbH) or the Flexi[®] Rabbit Reticulocyte Lysate System (Promega GmbH). To this end, a PCR template encoding the respective ORF under the SP6 promoter was generated from yeast chromosomal DNA using forward primers containing the SP6 promoter followed by a sequence that corresponds to the 5'-UTR of the respective ORF and reverse primers that bind to the 3'-end of the ORF. The *TMH11* ORF was amplified from yeast chromosomal DNA and cloned into pGEM-4Z plasmid (Promega) using EcoRI and HindIII restriction enzymes. Transcription of respective ORFs from PCR templates was performed using the mMESAGE mMACHINE[®] SP6 Transcription Kit (Thermo Fisher Scientific Inc.) and RNA was purified using the MEGAclean[™] Transcription Clean-Up Kit (Thermo Fisher Scientific Inc.). The RNA served as template for *in vitro* translation reactions. For generation of [³⁵S]Ybl039w-b, [³⁵S]Ykl023c-a, [³⁵S]Ykl065w-a, [³⁵S]Yor114w and [³⁵S]Yjr085c the PCR templates or the pGEM-4Z-Yjr085c plasmid were directly added to the TnT[®] Quick Coupled Transcription/Translation System. For import of radiolabeled precursor proteins, mitochondria were suspended in import buffer (3% [w/v] bovine serum albumin, 250 mM sucrose, 80 mM KCl, 5 mM MgCl₂, 5 mM methionine, 2 mM KP_i, 10 mM MOPS-KOH

[pH 7.2]) supplemented with 4 mM ATP, 4 mM NADH and an ATP regenerating system (200 µg/mL creatine kinase, 10 mM creatine phosphate). Where indicated, membrane potential across the inner mitochondrial membrane was dissipated prior to import ($-\Delta\psi$). To this end, import buffer was instead supplemented with 4 mM ATP, 1 µM valinomycin, 8 µM antimycin A, and 20 µM oligomycin (final concentrations). Import reaction was started by the addition of 5-10% [v/v] of *in vitro* translation/lysate system. If not stated otherwise, import in the presence of membrane potential ($+\Delta\psi$) was stopped after 30 min by the addition of 1 µM valinomycin, 8 µM antimycin A, and 20 µM oligomycin (final concentrations). Mitochondria were pelleted by centrifugation, washed with SEM buffer and suspended in 48 µL of SEM buffer. Where indicated, mitochondria were subjected to hypoosmotic swelling and/or protease treatment (7 µg/mL in SEM or EM buffer, see above). After inhibition of proteinase K activity by 2 mM PMSF, mitochondria were re-isolated and analyzed by SDS-PAGE or blue native gel electrophoresis and digital autoradiography.

Analysis of protein complexes by Blue Native gel electrophoresis

Mitochondria were solubilized in 1% digitonin buffer (20 mM Tris-HCl [pH 7.4], 0.1 mM EDTA, 50 mM NaCl, 10% [v/v] glycerol, 1% [w/v] digitonin, 1 mM PMSF) and incubated for 15 min on ice. Non-solubilized material was removed by centrifugation, blue native loading dye was added (final concentration: 100 mM Bis-Tris-HCl [pH 7.0], 50 mM ϵ -amino n-caproic acid, 0.5% [w/v] Coomassie G-250) and samples were subjected to 6–16.5 or 3–13% discontinuous polyacrylamide gels.

Generation of whole yeast cell extracts of *S. cerevisiae*

S. cerevisiae cells were grown overnight in 5-7 mL of YPG or YPD medium in 13 mL cell culture tubes (Sarstedt AG & Co. KG, 62.515.006). 2.5 OD₆₀₀ of cells were pelleted (4,000 x g for 2 min at 25°C) and suspended in 200 µL of 100 mM NaOH, followed by 5 min incubation at 25°C (Kushnirov, 2000). Cells were collected by centrifugation (4,000 x g for 2 min at 25°C), suspended in 75-150 µL of 2x SDS sample buffer supplemented with 2% (v/v) β -mercaptoethanol and incubated at 95°C for 3 min. Samples were centrifuged at 13,000 x g for 5 min at 25°C and a fraction of the supernatant was subjected to SDS-PAGE.

Miscellaneous

Radiolabeled proteins that had been separated by blue native gel electrophoresis or SDS-PAGE were visualized by autoradiography using the FLA-9000 (Fujifilm Holdings K.K.) and Typhoon FLA 7000 (GE Healthcare AG) image scanners as well as the Multi Gauge (Fujifilm) or ImageJ 1.49v software.

PCR primers used in this study

Name	Sequence (5' → 3')	Description
S3-Aim11_fw	TAAGCAATTGCAAGACCTCCTGT CGAGCGAAAACAACAAGCGTAC GCTGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>AIM11</i> ORF and C-terminal overhang for <i>AIM11</i> terminator
S2-Aim11_rv	CATTATTTACAGTTTAAAGAGAT TAAGCCAATGCGTAGTGATCGAT GAATTCGAGCTCG	
S3-DEG1_fw	ATGGAACCTGTGCAAGTTGTTAA TGCTAAATACTCCAAGAAAAAGA ACAACAAAATAAGCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>DEG1</i> ORF and C-terminal overhang for <i>DEG1</i> terminator
S2-DEG1_rv	AGCACGTGATGAAAAGAAATATA GTCTTCAAGGTTATATTACAG GTTTATATATTATTATCGATGAAT TCGAGCTCG	
S3-ECM4_fw	ACAAGGATCAACCCCTTGGAAT TACGCCCTGGGACCCAAGCCAG ATATTCGTCCTTACGTACGCTGC AGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>ECM4</i> ORF and C-terminal overhang for <i>ECM4</i> terminator
S2-ECM4_rv	GTATAGAAAAAGAGGCAACTCA GGGAGATTATAGACATTTGATTA TTTAATTACAGCTTGATCGATGA ATTCGAGCTCG	
S3-ECM19_fwd	TTATCATCCACACCAGCTGCACC ACCTACACCACCTACACCTCCTA CTCCACCACAACAGCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>ECM19</i> ORF and C-terminal overhang for <i>ECM19</i> terminator
S2-ECM19_rev	TTTACTATTGATGCTATATACAGG AAAAGAAAGTATAGAGGTATTTT CTAGTACGCTTCCATCGATGAAT TCGAGCTCG	
S3-FMP16_fw	TTGAAAAAAAAGGGAGATGACG CTAGAATCGAACAAAACAGGCCA GATGACGGTGTTTATCGTACGCT GCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>FMP16</i> ORF and C-terminal overhang for <i>FMP16</i> terminator
S2-FMP16_rv	GAATATAGATATGTGTATGACAG AAACTACATGCATATACGAGTTT GTACCAAGTGCTTCATCGATGAA TTCGAGCTCG	
S3-FYV4_fwd	TTTGGTGGTGAGAGGAAGAGAAA GGCATTACTGCTAAATGGAAAG CTGAAAACAAGCAACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>FYV4</i> ORF and C-terminal overhang for <i>FYV4</i> terminator
S2-FYV4_rev	TATGGAGTATAATGTGAACATAT ACGTATACATGTATATATTGAGA TCTTGCGAAAGCGTATCGATGAA TTCGAGCTCG	
S3-MEU1_fw	CCAGAGGCTATGTCCAAGGAAAC CTTAGAAAGACTAAGATACTTAT TTCCAAACTATTGGCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>MEU1</i> ORF and C-terminal overhang for <i>MEU1</i> terminator
S2-MEU1_rv	TTCCATAGCTATATATGTTTTCTC TCCTATTTATATTTTACATATGAT	

	TAGCGGCAACCAATCGATGAATT CGAGCTCG	
S3-MLO1_fwd	AATATGAAATCGGGTAGTAGTT CAGTCACCCCAGCTTTAAACAAT TGTTAATACAGAAGCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>MLO1</i> ORF and C-terminal overhang for <i>MLO1</i> terminator
S2-MLO1_rev	ATGATCGGAAGAAACGGTGTGCG CGGCGGAGGGAAGCGAAAAATT GGGAAACGGAAGCAAATCGATG AATTCGAGCTCG	
S3-MTC3_fwd	ACTGCATTTTATAATTGGAAACA AGATAAAAAGCTAGAGGAACAA TTAAGGGATCTTGTACGTACGCT GCAGGTGCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>MTC3</i> ORF and C-terminal overhang for <i>MTC3</i> terminator
S2-MTC3_rev	AAATGAATAGAACGTACTCTGCG CAAGCGCATATATATATACTA TTGTTGCTTCCAATATCGATGAAT TCGAGCTCG	
S3-NCE101_fw	TTTACACGAGCTCATAAAGAAGC GATGGGACGATCGCAAACGTACG CTGCAGGTGCGAC	Amplification of <i>3HA-kanMX6</i> fragment from pYM1 plasmid with N-terminal overhang for 3'-end of <i>FMP16</i> ORF and C-terminal overhang for <i>FMP16</i> terminator
S2-NCE101_rv	ATCGCCAGAACTTATATATACT CCCCTCACGCCGATTAATCGAT GAATTCGAGCTCG	
S3-PXP2_fw	TTGGCTATACTAATATCTCAGTG GTGTGGTGTAGTTGGAAATCTG GTGTTGTAATAATTGCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>PXP2</i> ORF and C-terminal overhang for <i>PXP2</i> terminator
S2-PXP2_rv	TTTTAATGAATAAGGGTACATAT ATTATGAGCATAACGAGTGGCCG ATCGGCAAAGGGGCATCGATGAA TTCGAGCTCG	
S3-RDL2_fwd	TATCCTGGTTCTATTACTGAGTGG TTAGCTAAAGGTGGTGCTGACGT TAAGCCCAAAAACGTACGCTGC AGGTGCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>RDL2</i> ORF and C-terminal overhang for <i>RDL2</i> terminator
S2-RDL2_rev	GAAATACACAAAAGGTTGTCTAT ATACAGGATATATCGATTATACT TGTTTCTTTTTGGCATCGATGAAT TCGAGCTCG	
S3-Rrg9_fwd	CAACAACTATACATTTTGAAGC ATTTGGGCTCGAAACAACGTACG CTGCAGGTGCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>RRG9</i> ORF and C-terminal overhang for <i>RRG9</i> terminator
S2-Rrg9_rev	ACAAGAAGTTTCGTAATAATATA TAAATCCCTCAAAGTGATCGAT GAATTCGAGCTCG	
S3-SMM1_fw	ACCGATCACATAGGCAGTGACAC TAAAAAGCAAAAGGTTGTACCCC TTCCACAGATATACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>SMM1</i> ORF and C-terminal overhang for <i>SMM1</i> terminator
S2-SMM1_rv	CATACATATATATTATGCGCG TTTTCTTTCATAATCCGTTCTTTTT ACTTAGAATATAATCGATGAATT CGAGCTCG	
S3-SUA5_fw	GAAGGATTAGCTGTTATGAACAG ATTGCGAAAGGCGGCTGCAAATA	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-

	ATTGTATACAGTTTCGTACGCTGC AGGTCGAC	ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>SUA5</i> ORF and C-terminal overhang for <i>SUA5</i> terminator
S2-SUA5_rv	CTAGATATAAAATCTCCGTAATC AGTGGTTATGATTTTCAAAGTT AATCACAGTTTTATATCGATGAA TTCGAGCTCG	
S3-TUM1_fw	GGATCCTGGACCGAGTGGGTCTT GAAATCCGGGCCCGAGTGGATTG CTGAAAACAGAGATCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>TUM1</i> ORF and C-terminal overhang for <i>TUM1</i> terminator
S2-TUM1_rv	AGATAGATAATTAATATATGTAG CTAAATAAATCGACTTGTCAAGA ATATATTTCTCTTAATCGATGAAT TCGAGCTCG	
S3-YBL039W-B_f	AATCGACTTCAACTCGAAGAGTA AGAAAAAAATGATAAACGTAC GCTGCAGGTCGAC	Amplification of <i>3HA-kanMX6</i> fragment from pYM1 plasmid with N-terminal overhang for 3'-end of <i>YBL039W-B</i> ORF and C-terminal overhang for <i>YBL039W-B</i> terminator
S2-YBL039W-B_r	TTCCATGGCGTGCTTTTACCAA GTACTGAACAGGGAGAATCGATG AATTCGAGCTCG	
S3-YBL059W_f	AAAAGAAGCCAGCAAGTAGTGG ACAGCTTAGTTAAGACACACAAT TCATCTCTTTGTAACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YBL059W</i> ORF and C-terminal overhang for <i>YBL059W</i> terminator
S2-YBL059W_r	ATGCGGGTAACACATATAAGTAG TGGAATATATTATTGAACTACT ACTATGGTATAACAATCGATGAA TTCGAGCTCG	
S3-YBR201C-A_f	TACTACAAGGACGAATTTTGTTT TCAACGATCATAACCAGGTTTC GTACGCTGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YBR201C-A</i> ORF and C-terminal overhang for <i>YBR201C-A</i> terminator
S2-YBR201C-A_r	CTCGGTAATTTTCTCCCTAATGAT AACCCATATTTTCGAAAGACTAT CGATGAATTCGAGCTCG	
S3-YBR230W-A_fw	GGCCCTATCACAACGGGTCAAGA AAGAGTATGCCGCAATCGTACG CTGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YBR230W-A</i> ORF and C-terminal overhang for <i>YBR230W-A</i> terminator
S2-YBR230W-A_rv	CCAAATATGTAAGTATATAAAAT ATATGTATGTGTGCAATCGAT GAATTCGAGCTCG	
S3-YDR286C_f	GAAGAAGACGATATCAGTGATAA AATAAGGAGAATGCAATCTAGAC GTACGCTGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YDR286C</i> ORF and C-terminal overhang for <i>YDR286C</i> terminator
S2-YDR286C_r	TTATAAATATTTACAGTGTTGTAA CTCTAGTAAAAACAAAAGAGGAT CGATGAATTCGAGCTCG	
S3-YDR381C-A_f	AGTGCTGTAACAAGAAAAAGAG GTGACAAATTAGTTTTTTAGAT AGGAGGAGAAACGAGCGTACGC TGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YDR381C-A</i> ORF and C-terminal overhang for <i>YDR381C-A</i> terminator
S2-YDR381C-A_r	GATGAATATATATACATAATA CTGCATTGAAAAAATGGATAGGT TGATTAACGTGTGATCGATGAA TTCGAGCTCG	
S3-YDR461C-A_fw	CCGCATCCAAGATCCCACAAGGG ATATCCGGGTGCGAAGACCTTA	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-

	CATATACTTCACGGCGTACGCTG CAGGTCGAC	ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YDR461C-A</i> ORF and C-terminal overhang for <i>YDR461C-A</i> terminator
S2-YDR461C-A_rv	ACACGCTCATATATATATATATA TATATATATATGTATATGTACATA TACCGCTTACCACATCGATGAAT TCGAGCTCG	
S3-YGL041W-A_f	CAGTTAGGACCTGAACAACTGGC CCCGCTAATGACCGTTTTAGGCC TTGAGAAGAAAAACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YGL041W-A</i> ORF and C-terminal overhang for <i>YGL041W-A</i> terminator
S2-YGL041W-A_r	ATATTTATACACGGAAGGCGCTA GTAAGACAGATAGGTAATGTTTT TTACTTTACCTCATCGATGAAT TCGAGCTCG	
S3-YGR021W_f	ACCACAGCGCTCGAGGACATCGA CGAAGTGACGTCGTTGTACTACTA ACGCTAGCAACGCTCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a- <i>TEV-ProtA-7His-hphNT1</i> plasmid with N-terminal overhang for 3'-end of <i>YGR021W</i> ORF and C-terminal overhang for <i>YGR021W</i> terminator
S2-YGR021W_r	TGTTTGTTTAGTATATACAGGTAT TTGTGGTATTGATTATTTATTTAC GGTTTGGTTGGAATCGATGAATT CGAGCTCG	
S3-YGR053C_f	GAAGATGTTGAACTGATCCATTA CGAGAAGAAAATTGCCACTCGCG GTGCATTTGCATGTCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a- <i>TEV-ProtA-7His-hphNT1</i> plasmid with N-terminal overhang for 3'-end of <i>YGR053C</i> ORF and C-terminal overhang for <i>YGR053C</i> terminator
S2-YGR053C_r	TTTATGATGGTGTGTTTATTATTT ACAAATAAAGATGCTATCTAATT CTTGTATATCGTAATCGATGAATT CGAGCTCG	
S3-YHL018W_f	AGCGATATAGACGTCCGGATGGC CAAGAGAATAGATTCCTACATCG ATGAGATGACAACCTCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YHL018W</i> ORF and C-terminal overhang for <i>YHL018W</i> terminator
S2-YHL018W_r	AAAATCGATAGGAAAAAAAAAAAA TGGAAGAGCAATAGCCTTTTATG ACCTGCCCTGCCTTAATCGATGA ATTCGAGCTCG	
S3-YIL002W-A_fw	GTTCAGTTAGAAGATCTACACAG GGACAACAATGATTTGGCAAAAA GTTCCAGCCAAAAACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a- <i>TEV-ProtA-7His-hphNT1</i> plasmid with N-terminal overhang for 3'-end of <i>YIL002W-A</i> ORF and C-terminal overhang for <i>YIL002W-A</i> terminator
S2-YIL002W-A_rv	AGATTATATATAAATATATTTATT TAGCGTTCCTTCCCTATGCGGCTG TGAGCGTACTCTATCGATGAATT CGAGCTCG	
S3-YIL077C_fw	TCTTCTGATGACAAATATCAGCG TTTACTGCAGAGCGGGAGATATG GTGGGAACCGCTCCCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YIL077C</i> ORF and C-terminal overhang for <i>YIL077C</i> terminator
S2-YIL077C_rv	TTGGCCCAAACATATACCTATAT ACAAATTAGTGACCATACGCTAT TATTACTCCGTCGTATCGATGAAT TCGAGCTCG	
S3-YIL156W-B_fw	TGTTGCCACTTGTGGCTCCTCGAC GTATTTCCGCTAGGAAACGTACGC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid

	TGCAGGTCGAC	with N-terminal overhang for 3'-end of <i>YIL156W-B</i> ORF and C-terminal overhang for <i>YIL156W-B</i> terminator
S2-YIL156W-B_rv	TATACAAAATATTATCAACATAT ATAGAATATAAACGTACATCGAT GAATTCGAGCTCG	
S3-YJL133C_f	TATGGTCCCTTGAGTGCCTCACTA GCTACCAGAAGACACTTGGCTCA CGCGCCAAAGTTGCGTACGCTGC AGGTTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YJL133C</i> ORF and C-terminal overhang for <i>YJL133C</i> terminator
S2-YJL133C_r	ATAGGTAAAGTCATCATCATT ATAAACCCAGGAAAGAAAAGAGC ATAATTAAGATACGCATCGATGA ATTCGAGCTCG	
S3-YJR085C_f	ACAGCTTTGGGTGGGCTCGGCAG TACTACTATTATAACAAATACA AGGAATTTTACCCTCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YJR085C</i> ORF and C-terminal overhang for <i>YJR085C</i> terminator
S2-YJR085C_r	AGAATATCATATTATATATATAT ATAGGGTCGTATATATATCGTGC GTCTTCTTCTTCTATCGATGAAT TCGAGCTCG	
S31-YKL065W-A_f	TTTATAAGAACGATAGCAAACAT AGTGAAATTAAGATATACCA AAATGAGAAAAAATTTCGTACGC TGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YKL065W-A</i> ORF and C-terminal overhang for <i>YKL065W-A</i> terminator
S21-YKL065W-A_r	GAGCTTCGTGACTCGGTTTACCA TTCTGTGTTATATACGAAAACCCT TATATAACAACCTTATCGATGAA TTCGAGCTCG	
S3-YKL133C_f	TTTAATGAACTCATTGAGGAAGC TCAACGTGAACTTAAAAAGGTTG ATGGTACGCCTATACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YKL133C</i> ORF and C-terminal overhang for <i>YKL133C</i> terminator
S2-YKL133C_r	TCTTTCGGCATTTCACATATTGA TGAAAATTTAGAAGAATATATAT ACAAATGTAAGATATCGATGAAT TCGAGCTCG	
S3-YLR049C_f	AAAATCGATTGTGACTTAGTCAT TCTGCTAGAAGATTTAAGGTCAC GGATTGATTAGATCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YLR049C</i> ORF and C-terminal overhang for <i>YLR049C</i> terminator
S2-YLR049C_r	ACATATATATATATATATTCTT CCAGGAGAAATACCTGCCTTCTC TCTTACCCCTTGCATCGATGAATT CGAGCTCG	
S3-YLR118C_fw	TCTACAGTTCAGATGAATTAGA AGACTTGGCTTCATTTATCAAGA AGAGCTTATCATCACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YLR118C</i> ORF and C-terminal overhang for <i>YLR118C</i> terminator
S2-YLR118C_rv	ATAAATTATTTAACCAAGTATAA TAGCGGTTAAATTGAACTCGCAA TATTAGGAAAAGAGATCGATGAA TTCGAGCTCG	
S3-YLR281C_f	GTCGAAAAAGAGGAACGCGAGG CCCAGACAGAGAAATGGTGC GC GAGTTATTCCGCCGCGTACGCT GCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of

S2-YLR281C_r	AGTTACACTTTTTTTTTTCATTTTTT TTTTTCTTTTCTCCTCCATCTAATT TCACCTGCGGATCGATGAATTCG AGCTCG	<i>YLR281C</i> ORF and C-terminal overhang for <i>YLR281C</i> terminator
S3-YLR307C-A_f	CCAATTGTTATTTGCACTGACAA CGAAGAGGTAGAGACTGTATCGG AGCACGTAAAAGTACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS- hphNT1</i> fragment from pFA6a-TEV- ProtA-7His-hphNT1 plasmid with N- terminal overhang for 3'-end of <i>YLR307C-A</i> ORF and C-terminal overhang for <i>YLR307C-A</i> terminator
S2-YLR307C-A_r	CTTTAGCGTCAAACGTTACAC GTACATTTGAACAGTGTTAAGAG TAGATTAATTCAAAATCGATGAA TTCGAGCTCG	
S3-YML007C-A_f	AGATTGGTGAGAAACCTCCAATA CTTGCTGTTGCCGATAACTTCTTC ATTGCTTTTTATACGTACGCTGCA GGTCGAC	Amplification of <i>TEV-ProtA-7HIS- hphNT1</i> fragment from pFA6a-TEV- ProtA-7His-hphNT1 plasmid with N- terminal overhang for 3'-end of <i>YML007C-A</i> ORF and C-terminal overhang for <i>YML007C-A</i> terminator
S2-YML007C-A_r	AATTACAAAGACGAATGGAAAG AAAAAGAGAGCAGCAGAGGATA GGAGAAAACACCCGAAATCGAT GAATTCGAGCTCG	
S3-YMR087W_fw	AATGTAGAAAAAGATGCAATAG AATTGCTCATTCTAGAAAGGATT TTGACCTTGATTACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS- hphNT1</i> fragment from pFA6a-TEV- ProtA-7His-hphNT1 plasmid with N- terminal overhang for 3'-end of <i>YMR087W</i> ORF and C-terminal overhang for <i>YMR087W</i> terminator
S2-YMR087W_rv	TACTTACAAACGTAGTTCAAGTT CTTGAAAAATTGAAAATAGACTT TATATTATATACCTATCGATGAAT TCGAGCTCG	
S3-YMR130W_f	CAGTTGAGTGAAAGAAAGTACGT TGTTTCGAATCTTGAGGTTTTAGA GGAACCTTTCCCGTACGCTGC AGGTCGAC	Amplification of <i>TEV-ProtA-7HIS- hphNT1</i> fragment from pFA6a-TEV- ProtA-7His-hphNT1 plasmid with N- terminal overhang for 3'-end of <i>YMR130W</i> ORF and C-terminal overhang for <i>YMR130W</i> terminator
S2-YMR130W_r	TATTACTGGCGGATATGAATAAT ATCTATCATACATATTTGTTACTG TAACGTTAGGCGCATCGATGAAT TCGAGCTCG	
S3-YMR182W-A_f	GCATGTAACATAATATTTCTTCCC CTCGTTAAGTGTGCATCAGCAAC CATAATGCTGAATCGTACGCTGC AGGTCGAC	Amplification of <i>γEGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>YMR182W-A</i> ORF and C-terminal overhang for <i>YMR182W-A</i> terminator
S2-YMR182W-A_r	ATGCAGGAGAAAGGGCGAGTTTT GTTTATATGCGATCCTTTATGGTA ACCTTTGCGGTTAATCGATGAAT TCGAGCTCG	
S3-YNR040W_f	AAGACCCTATTTCACTCGGGGAA TTCTAGATCATCCATCAAGAATA TCGTGAAGCCCAAACGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS- hphNT1</i> fragment from pFA6a-TEV- ProtA-7His-hphNT1 plasmid with N- terminal overhang for 3'-end of <i>YNR040W</i> ORF and C-terminal overhang for <i>YNR040W</i> terminator
S2-YNR040W_r	TGTTTAATTGTTCTTTTATTCGAT TATTATTTACATGCATTCTTACGT AGATGTCCGTACATCGATGAATT CGAGCTCG	
S3-YOR020W-A_f	AGTATTGAAGGATTTTTAAATGA TTTAGAGAAAGATACGAGGCAGG ATACGAAAGCCAACCGTACGCTG CAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS- kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YOR020W-A</i> ORF and C-terminal

S2-YOR020W-A_r	CAGTGCATCCCTTCACTGAACGA TGAAGAACACCACCATTTTCAGAA ATTTTTATACATAAATCGATGAA TTCGAGCTCG	overhang for <i>YOR020W-A</i> terminator
S3-YPL107W_f	AAGAAAAGATTACAAAAAATTCG CCGACAGGAAGAAATAAAAAAG AGGACAGCTTTGGTTCGTACGCT GCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YPL107W</i> ORF and C-terminal overhang for <i>YPL107W</i> terminator
S2-YPL107W_r	ACCAAGATCAAAGAAGACCGCA AATATTGTACATAGGCTTTTCAAT AATATATATTTGCTATCGATGAA TTCGAGCTCG	
S3-YPR010C-A_fw	ATTATCTAGTTTAGATGAAGTCCT TGCCAAAGATAAGGATCGTACGC TGCAGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YPR010C-A</i> ORF and C-terminal overhang for <i>YPR010C-A</i> terminator
S2-YPR010C-A_rv	TAAGTGTAGATATTAATATAACA AATCATCAACATGGTTTATCGAT GAATTCGAGCTCG	
S3-YPR098C_f	TGCGGGATGCTTTCGTACGGTGT TTGTTTGTACAGGTGGTTTGTAAAG AAAAATTCCAAAACGTACGCTGC AGGTCGAC	Amplification of <i>TEV-ProtA-7HIS-hphNT1</i> fragment from pFA6a-TEV-ProtA-7His-hphNT1 plasmid with N-terminal overhang for 3'-end of <i>YPR098C</i> ORF and C-terminal overhang for <i>YPR098C</i> terminator
S2-YPR098C_r	ACCTTGGCAAAGGAATGACGAAA AAATGATCTTGCATATATATATTT ACTTGTAATAAATCGATGAAT TCGAGCTCG	
Meo1_S3_fw	GACCCAGAACAAAAAGAGCAA TCAAGCGTCTCCACCAGTTGGAC GGCATTCTCACGCTCGTACGCT GCAGGTCGAC	Amplification of γ <i>EGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>MEO1</i> ORF and C-terminal overhang for <i>MEO1</i> terminator
Meo1_S2_rv	ATCCGTAATTGAAAAAAAAAAAA GAAAAGATCAAGGAACACATC ACCCTGGGCACATCAATCGATGA ATTCGAGCTCG	
YBL039W-B_S3_fw	ATGTTTCATGTCATCACCTACAATC GACTTCAACTCGAAGAGTAAGAA AAAAAATGATAAACGTACGCTGC AGGTCGAC	Amplification of γ <i>EGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>YBL039W-B</i> ORF and C-terminal overhang for <i>YBL039W-B</i> terminator
YBL039W-B_S2_rv	AGGAAGTTCTAAATAATTTTCCA TGCGCTGCTTTACCAAAGTACT GAACAGGGAGATTAATCGATGAA TTCGAGCTCG	
YFR032C-B_S3_fw	ATTTTCCCTTCTCTTTTTTTTCTT TATATCCTCCACTATATGCCACTC AGCGCACCTCAATCGATGAATTC GAGCTCG	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid and γ <i>EGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>YFR032C-B</i> ORF and C-terminal overhang for <i>YFR032C-B</i> terminator
YFR032C-B_S2_rv	AGTACGTTATATTATCATGAAGT CCCCATTTCCGCCATTGGAAATG CAGGTAGCCAAATCCGTACGCTG CAGGTCGAC	
YGL204C_S3_fw	GAAATGTGGAAATATGCCTTTGC TGTTTCGCTACTTCTTAATAGTTT GGCACTATTTTTTCGTACGCTGCA GGTCGAC	Amplification of γ <i>EGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>YGL204C</i> ORF and C-terminal overhang for <i>YGL204C</i> terminator
YGL204C_S2_rv	TATAAGGGAGGAGAAATGATGGT	

	GATATTAATATGCAGAAATATCG ATTCCATTTTTTCAATCGATGAAT TCGAGCTCG	
YKL023C-A_S3_forward	ATTTCCGGTAAACTATCAAAAGAA CGAACCAGTTGAATTTCTTGAAC GTACGCTGCAGGTGCAC	Amplification of <i>3HA-kanMX6</i> fragment from pYM1 plasmid and <i>γEGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>YKL023C-A</i> ORF and C-terminal overhang for <i>YKL023C-A</i> terminator
YKL023C-A_S2_reverse	GCGAGAAAGCTGGCTGTGATGTA GTGGCAGCTGTCATTTGTCTTAAT CGATGAATTCGAGCTCG	
YKL065w-a_fw	TATAAGAACGATAGCAAACATAG TGAAATTA AAAAGATATACCAA ATGAGAAAAAAATTCGTACGCTG CAGGTTCGAC	Amplification of <i>3HA-kanMX6</i> fragment from pYM1 plasmid and <i>γEGFP-kanMX4</i> fragment from pYM12 plasmid with N-terminal overhang for 3'-end of <i>YKL065W-A</i> ORF and C-terminal overhang for <i>YKL065W-A</i> terminator
YKL065w-a_rv	TTCGTGACTCGGTTTACCATTCTG TGTTATATACGAAAACCCTTATA TAACAACTTTTTAATCGATGAATT CGAGCTCG	
YOR114W_S3_fw	CACGATACAAATACACAACCAAA TAATATTCTTCCCATGACGTACCT ACTAAAAAAGAAACGTACGCTGC AGGTGCAC	Amplification of <i>TEV-ProtA-7HIS-kanMX6</i> fragment from pYM9 plasmid with N-terminal overhang for 3'-end of <i>YOR114W</i> ORF and C-terminal overhang for <i>YOR114W</i> terminator
YOR114W_S2_fw	TCTTTATGATAGTGAAGTGCTTTC GTGGACTCTTCTTAAAGCCCTAA AAGTCTATTGTCAATCGATGAAT TCGAGCTCG	
AIM11_seq_fwd	GGATGGGGAACCTTGATTC	Analysis of genomic integration of ProtA tag behind <i>AIM11</i> ORF
AIM11_seq_rev	CGGCTCGTAGTTATACC	
DEG1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCACATGCAATC TTACTGC	Analysis of genomic integration of ProtA tag behind <i>DEG1</i> ORF (together with reverse primer: ProtA_seq_rev)
ECM4_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCCTTACCATCAC TACAGTG	Analysis of genomic integration of ProtA tag behind <i>ECM4</i> ORF (together with reverse primer: ProtA_seq_rev)
ECM19_seq_fwd	CTTCTTCTATCTTTTCCGC	Analysis of genomic integration of ProtA tag behind <i>ECM19</i> ORF (together with reverse primer: ProtA_seq_rev)
FMP16_seq_fwd	CTTGTTCCCTACAATACTCC	Analysis of genomic integration of ProtA tag behind <i>FMP16</i> ORF
FMP16_seq_rev	GACTAAATACGATAGGACC	
FYV4_seq_fwd	CCATCTTCAAACAAGAGC	Analysis of genomic integration of ProtA tag behind <i>FYV4</i> ORF (together with reverse primer: ProtA_seq_rev)
MATlocus_fw	AGTCACATCAAGATCGTTTATGG	Discrimination between mating types MAT α and MAT α Huxley et al., 1990
Mat(a)_rv	ACTCCACTTCAAGTAAGAGTTTG	
Mat(alpha)_rv	GCACGGAATATGGGACTACTTCG	
MEU1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCCTTATTGTCAA TTATGTGAAAAG	Analysis of genomic integration of ProtA tag behind <i>MEU1</i> ORF (together with reverse primer: ProtA_seq_rev)
MLO1_seq_fwd	CCATATAAGCAGCAAAACG	Analysis of genomic integration of ProtA tag behind <i>MLO1</i> ORF (together with reverse primer: ProtA_seq_rev)
MTC3_seq_fwd	GTACTIONCATCCTTCTTCG	Analysis of genomic integration of ProtA tag behind <i>MTC3</i> ORF (together

		with reverse primer: ProtA_seq_rev)
NCE101_seq_fwd	GCATACAAATGTTCACTCC	Analysis of genomic integration of ProtA tag behind <i>NCE101</i> ORF
NCE101_seq_rev	GTATAAAGGTAGATCCTAGG	
ProtA_seq_rev	TGGTGGGAATTCGCGTCTAC	Reverse primer for the analysis of genomic integration of ProtA tag behind <i>DEG1</i> , <i>ECM4</i> , <i>ECM19</i> , <i>FYV4</i> , <i>MEU1</i> , <i>MLO1</i> , <i>MTC3</i> , <i>PXP2</i> , <i>SMM1</i> , <i>SUA5</i> , <i>TUM1</i> , <i>YBL059W</i> , <i>YDR461C-A</i> , <i>YGR021W</i> , <i>YGR053C</i> , <i>YHL018W-A</i> , <i>YIL002W-A</i> , <i>YKL133C</i> , <i>YLR049C</i> , <i>YLR118C</i> , <i>YLR281C</i> , <i>YML007C-A</i> , <i>YMR087W</i> , <i>YMR130W</i> , <i>YNR040W</i> , <i>YPL107W</i> and <i>YPR098C</i> ORFs (together with respective forward primers)
PXP2_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCCATATATACAA GTGCCACAC	Analysis of genomic integration of ProtA tag behind <i>PXP2</i> ORF (together with reverse primer: ProtA_seq_rev)
RDL2_seq_fwd	CTCTCAACAAATGGAAGCG	Analysis of genomic integration of ProtA tag behind <i>RDL2</i> ORF
RDL2_seq_rev	GAATGAATCGGAGAGGTG	
RRG9_seq_fwd	GACTTCCTTCTGAATCATTG	Analysis of genomic integration of ProtA tag behind <i>RRG9</i> ORF
RRG9_seq_rev	GTTCCACTAACGATATTACTG	
SMM1_seq_f	CTATTTTCATCCATCCAAGC	Analysis of genomic integration of ProtA tag behind <i>SMM1</i> ORF (together with reverse primer: ProtA_seq_rev)
SUA5_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGTTAACTTACC ACTAAACCTG	Analysis of genomic integration of ProtA tag behind <i>SUA5</i> ORF (together with reverse primer: ProtA_seq_rev)
TUM1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCCATATAAAGTTGT GAAGAAAATTGC	Analysis of genomic integration of ProtA tag behind <i>TUM1</i> ORF (together with reverse primer: ProtA_seq_rev)
YBL059W_seq_f	GTTCAGCTTCTAACTGTG	Analysis of genomic integration of ProtA tag behind <i>YBL059W</i> ORF (together with reverse primer: ProtA_seq_rev)
YBR201C-A_seq_f	CTGTGACAAGGCGAAAAC	Analysis of genomic integration of ProtA tag behind <i>YBR201C-A</i> ORF
YBR201C-A_seq_r	CCTGTCTCAAATGTATAAATG	
YBR230W-A_seq_f	AGGCAAGAACAGAGGAG	Analysis of genomic integration of ProtA tag behind <i>YBR230W-A</i> ORF
YBR230W-A_seq_r	GGAGAGATTCTATATACCAC	
YDR286C_seq_f	GTAGAGAGTTCGGAGTTG	Analysis of genomic integration of ProtA tag behind <i>YDR286C</i> ORF
YDR286C_seq_r	CCTGCGTAAGAAGTATGC	
YDR381C-A_seq_f	CAATCTTCCTCCTTACAAAC	Analysis of genomic integration of ProtA tag behind <i>YDR381C-A</i> ORF
YDR381C-A_seq_r	GGATAGGTTGATTAACGTG	
YDR461C-A_seq_f	GTGTTGAGTATTCAAAGCAC	Analysis of genomic integration of ProtA tag behind <i>YDR461C-A</i> ORF (together with reverse primer: ProtA_seq_rev)
YGL041W-A_seq_f	GATGTAACAAAACCGACG	Analysis of genomic integration of ProtA tag behind <i>YGL041W-A</i> ORF
YGL041W-A_seq_r	GCTAGTAAGACAGATAGG	
YGR021W_seq_f	GTACGCACGTACGCAAG	Analysis of genomic integration of

		ProtA tag behind <i>YGR021W</i> ORF (together with reverse primer: ProtA_seq_rev)
YGR053C_seq_f	GGATGACAGGTATGAGC	Analysis of genomic integration of ProtA tag behind <i>YGR053C</i> ORF (together with reverse primer: ProtA_seq_rev)
YHL018W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGCACAACAA GATTGTTAGAAT	Analysis of genomic integration of ProtA tag behind <i>YHL018W</i> ORF (together with reverse primer: ProtA_seq_rev)
YIL002W-A_seq_f	CAATTACTCTAGAGAAATCG	Analysis of genomic integration of ProtA tag behind <i>YIL002W-A</i> ORF (together with reverse primer: ProtA_seq_rev)
YIL077C_seq_f	GCAGCCTAATTTCAAGCG	Analysis of genomic integration of ProtA tag behind <i>YIL077C</i> ORF
YIL077C_seq_r	CAAATTAGTGACCATACGC	
YIL156W-B_seq_f	CGGAGAGAAAGATCGAAC	Analysis of genomic integration of ProtA tag behind <i>YIL156W-B</i> ORF
YIL156W-B_seq_r	CTGGGTGCCGTTATACA	
YJL133C-A_seq_f	GTAGGGAGATGTTTAATGTG	Analysis of genomic integration of ProtA tag behind <i>YJL133C-A</i> ORF
YJL133C-A_seq_r	CAGGACCCCAAAGAAG	
YJR085C_seq_f	GGAGACAAGACAGAAACG	Analysis of genomic integration of ProtA tag behind <i>YJR085C</i> ORF
YJR085_seq_r	CGTATATATATCGTGCGTC	
YKL065W-A_seq_f	GGACTTTGGACCTAACTC	Analysis of genomic integration of ProtA tag behind <i>YKL065W-A</i> ORF
YKL065W-A_seq_r	CCACAGAACCGACCATTA	
YKL133C_seq_f	GGCACAAAGTGAGAACG	Analysis of genomic integration of ProtA tag behind <i>YKL133C</i> ORF (together with reverse primer: ProtA_seq_rev)
YLR049C_seq_f	CGTTAGCCAAATTCTTTGG	Analysis of genomic integration of ProtA tag behind <i>YLR049C</i> ORF (together with reverse primer: ProtA_seq_rev)
YLR118C_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGAACACATACA CTATATTTGC	Analysis of genomic integration of ProtA tag behind <i>YLR118C</i> ORF (together with reverse primer: ProtA_seq_rev)
YLR281C_seq_f	CCAACCTACAAGCTATGC	Analysis of genomic integration of ProtA tag behind <i>YLR281C</i> ORF (together with reverse primer: ProtA_seq_rev)
YLR307C-A_seq_f	GACAGCCAAGTATACTTG	Analysis of genomic integration of ProtA tag behind <i>YLR307C-A</i> ORF
YLR307C-A_seq_r	CGTTACACGTACATTTGAAC	
YML007C-A_seq_f	GTAAACTGCTCCACTTCG	Analysis of genomic integration of ProtA tag behind <i>YML007C-A</i> ORF (together with reverse primer: ProtA_seq_rev)
YMR087W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATCTTCGTAA TTTTGAACTG	Analysis of genomic integration of ProtA tag behind <i>YMR087W</i> ORF (together with reverse primer: ProtA_seq_rev)

YMR130W_seq_f	CATCCACAGAAATTGGCTC	Analysis of genomic integration of ProtA tag behind <i>YMR130W</i> ORF (together with reverse primer: ProtA_seq_rev)
YNR040W_seq_f	GTTATCTTATGTGGACTAGG	Analysis of genomic integration of ProtA tag behind <i>YNR040W</i> ORF (together with reverse primer: ProtA_seq_rev)
YOR020W-A_seq_f	GTAAGAAAGGTCGCTACTG	Analysis of genomic integration of ProtA tag behind <i>YOR020W-A</i> ORF
YOR020W-A_seq_r	GAAGAACACCACCATTTC	
YPL107W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCAGTCAGAGGG TGCATG	Analysis of genomic integration of ProtA tag behind <i>YPL107W</i> ORF (together with reverse primer: ProtA_seq_rev)
YPR010C-A_seq_f	GGTAGTTTCCTGAACGAC	Analysis of genomic integration of ProtA tag behind <i>YPR010C-A</i> ORF
YPR010C-A_seq_r	CGATTTGTTCCCGACAATT	
YPR098C_seq_f	GGAGTTGTTTGATGATATAGG	Analysis of genomic integration of ProtA tag behind <i>YPR098C</i> ORF (together with reverse primer: ProtA_seq_rev)
AIM11_SP6_fwd	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGATCGAAGA GAAGAAGG	Synthesis of SP6-AIM11 PCR product as template for subsequent RNA generation
AIM11_SP6_rev	CTACTTGTTGTTTTCGCTC	
DEG1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCACATGCAATC TTACTGTC	Synthesis of SP6-DEG1 PCR product as template for subsequent RNA generation
DEG1_rev	TTACTTATTTTTGTTGTTCTTTTTTC TTG	
FMP16_SP6_fwd	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGTTGAGAAC CACTTTTTTG	Synthesis of SP6-FMP16 PCR product as template for subsequent RNA generation
FMP16_SP6_rev	CCAGATGACGGTGTTTATTAA	
HBN1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGTAGACTGAAG TATCCTATATC	Synthesis of SP6-HBN1 PCR product as template for subsequent RNA generation
HBN1_rev	TTAATTGAAGATTTCAACATCGTT	
MEU1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCCTTTATTGTCAA TTATGTGAAAAG	Synthesis of SP6-MEU1 PCR product as template for subsequent RNA generation
MEU1_rev	TTACCAATAGTTTGAAATAAGT A	
RDL2_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGTTCAAGCA TAGTACAGG	Synthesis of SP6-RDL2 PCR product as template for subsequent RNA generation
RDL2_SP6_r	TTACATTTTTTTGGGCTTAACGTC AG	
RRG9_SP6_fwd	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGAACATTCT GCGAATAGC	Synthesis of SP6-RRG9 PCR product as template for subsequent RNA generation
RRG9_SP6_rev	TTATTGTTTCGAGCCCAAATG	
SMM1_SP6_f	TCGATTTAGGTGACACTATAGAA	Synthesis of SP6-SMM1 PCR product

	TACGCCGCCGCCCTATTTTCATCCA TCCAAGC	as template for subsequent RNA generation
SMM1_rev	TTATATATCTGTGGGAAGGG	
TES1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCCTACAATGAA AAACCACG	Synthesis of SP6-TES1 PCR product as template for subsequent RNA generation
TES1_rev	TCAGAACTTGGCTCGAATG	
TUM1_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATAAAGTTGT GAAGAAAATTGC	Synthesis of SP6-TUM1 PCR product as template for subsequent RNA generation
TUM1_rev	TTAATCTCTGTTTTTCAGCAATC	
YBL039W-B_Sp6_fw	GATCGATTTAGGTGACACTATAG AAGCGGCCACCATGGGCTTTTTT ACAATAATCCGGTAATT	Synthesis of SP6-YBL039W-B PCR product as template for generation of [³⁵ S]Ybl039w-b using the TnT® Quick Coupled Transcription/Translation System
YBL039W-B_Sp6_rv	GATCTTATTTATCATTTTTTTTTCTT ACTCTTCGA	
YBR230W-A_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGAGAAACGA ATTATACCAGT	Synthesis of SP6-YBR230W-A PCR product as template for subsequent RNA generation
YBR230W-A_SP6_r	CTACATATTGGCGGCATACTCTTT	
YDR286C_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGTTGAGAGC GTTCCG	Synthesis of SP6-YDR286C PCR product as template for subsequent RNA generation
YDR286C_SP6_r	TCACATTCTAGATTGCATTCTCCT TATT	
YGL041W-A_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGCTAAGAGT TATATGGAAG	Synthesis of SP6-YGL041W-A PCR product as template for subsequent RNA generation
YGL041W-A_SP6_r	TCATTTTTTCTTCTCAAGGCC	
YGR021W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCAGCAAACCTTG GTATGAAC	Synthesis of SP6-YGR021W PCR product as template for subsequent RNA generation
YGR021W_rev	TTAAGCGTTGCTAGCGTTAG	
YHL018W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGCACAACAA GATTGTTAGAAT	Synthesis of SP6-YHL018W PCR product as template for subsequent RNA generation
YHL018W_SP6_r	TCAAGTTGTCATCTCATCGA	
YIL077C_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGTTGGGAAA AGAAGAAGAA	Synthesis of SP6-YIL077C PCR product as template for subsequent RNA generation (eventually used for generation of radiolabeled Yil077c precursor used for assessment of import and assembly under native conditions; Fig. 6C, lanes 9-18)
YIL077C_SP6_r	CTAGGAGCGGTTCCC	
YIL077C SP6 fw	GATCGATTTAGGTGACACTATAG AAGCGGCCACCATGTTGGGAAAA GAAGAAGAACAGC	Synthesis of SP6-YIL077C PCR product as template for subsequent RNA generation (eventually used for generation of radiolabeled Yil077c precursor used for assessment of import under denaturing conditions; Fig. S5, lanes 1-6)
YIL077C SP6 rv	CTAGGAGCGGTTCCCACC	
YJL133C-A_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATGATCGCCA AAGTACC	Synthesis of SP6-YJL133C-A PCR product as template for subsequent RNA generation

YJL133C-A_SP6_r	CTACATCAACTTTGGCGCGTGA	
YKL023C-A_Sp6_fw	GATCGATTTAGGTGACACTATAG AAGCGGCCACCATGAATCCACGC TACAGGTTTATA	Synthesis of SP6-YKL023C-A PCR product as template for generation of [³⁵ S]Ykl023c-a using the TnT® Quick Coupled Transcription/Translation System
YKL023C-A_Sp6_rv	GATCTTACATCATTTC AAGAAAT TCAACTGGTTC	
YKL065W-A_Sp6_fw	GATCGATTTAGGTGACACTATAG AAGCGGCCACCATGAGATCTAAT ATTTTGAAATTA	Synthesis of SP6-YKL065W-A PCR product as template for generation of [³⁵ S]Ykl065w-a using the TnT® Quick Coupled Transcription/Translation System
YKL065W-A_Sp6_rv	GATCTTAAATTTTTTTTCTCATT GGTA	
YKL133C_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGTAATGAAGAA ATTTATGCTCATTTAG	Synthesis of SP6-YKL133C PCR product as template for subsequent RNA generation
YKL133C_rev	TTATATAGGCGTACCATCAAC	
YLR118C_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGAACACATACA CTATATTTGC	Synthesis of SP6-YLR118C PCR product as template for subsequent RNA generation
YLR118C_rev	CTATGATGATAAGCTCTTCTT	
YLR281C_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGCTATACGTT CTAGAAC	Synthesis of SP6-YLR281C PCR product as template for subsequent RNA generation
YLR281C_rev	TTACCGGCGGAATAACTC	
YMR130W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCGACTTTTGA ACATCCAC	Synthesis of SP6-YMR130W PCR product as template for subsequent RNA generation
YMR130W_rev	TCAGGGAAAGAGTTCCTC	
YNR040W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCATAAGAAGGT GCATTTAGTTATC	Synthesis of SP6-YNR040W PCR product as template for subsequent RNA generation
YNR040W_rev	CTATTTGGGCTTCACGATATTC	
YOR114W_Sp6_fw	GATCGATTTAGGTGACACTATAG AAGCGGCCACCATGAAGGCTACT TFACTGTTGAAGGCCAG	Synthesis of SP6-YOR114W PCR product as template for generation of [³⁵ S]Yor114w using the TnT® Quick Coupled Transcription/Translation System
YOR114W_Sp6_rv	GATCTCATTCTTTTTTAGTAGGT ACGTCATGGG	
YPL107W_SP6_f	TCGATTTAGGTGACACTATAGAA TACGCCGCCGCCAGTCAGAGGG TGCATG	Synthesis of SP6-YPL107W PCR product as template for subsequent RNA generation
YPL107W_rev	TCAAACCAAAGCTGTCCTC	
TEV-ProtA_fwd	CCCAAGCTTGGGCGTACGCTGCA GGTCGAC	Generation of HindIII-TEV-ProtA-7HIS-XmaI PCR product for cloning into pFA6a-hphNT1 plasmid
TEV-ProtA_rev	TCCCCCGGGGGGACTAGGAATT CGCGTCTAC	
YJR085C_pGEM-4Z_FW	GGAATTCAGATCACACTATGGAA CATCCA	Generation of EcoRI-YJR085C (<i>S. cerevisiae</i>)-HindIII PCR product for cloning into pGEM-4Z plasmid
YJR085C_pGEM-4Z_REV	CCCAAGCTTTCTTCTTCTTCTCT CTAAGGGTAA	
YJR085C_pRS425_FW	CCCAAGCTTATCAAACATTTAGA GCCTGAAACAA	Generation of HindIII-P _{YJR085C} -YJR085C (<i>S. cerevisiae</i>)-T _{YJR085C} -BamHI PCR product for cloning into pRS425 plasmid
YJR085C_pRS425_REV	CGCGGATCCAGGTGGTCTTCACT CGGTT	

NHA_YJR085C_FW	CATCAGATCACACTATGTACCCA TACGATGTTCCAGATTACGCTGA ACATCCAGCATATAC	Insertion of a single N-terminal HA-tag behind the start codon of <i>YJR085C</i> in pRS425-YJR085C resulting in pRS425-HA YJR085C plasmid
NHA_YJR085C_Rev	GTATATGCTGGATGTTACGCGTA ATCTGGAACATCGTATGGGTACA TAGTGTGATCTGATG	
AIM11 N' tag CHK R	TACAAACTCTCTTGCCGTTG	Primer for checking N' tagging of the gene. AIM11
AIM11 N' tag pYM F	TTAGTACGACTATCCTACTTCATC AAGAAACGAACTATGcgtacgctgca ggtcgac	Primer for N' tagging of gene using pYM plasmids. AIM11
AIM11 N' tag pYM R	GAAGAACCCTGCGTTTCTTAAGT TCCTTCTTCTCTTCGATcatcgatgaatt ctctgctg	Primer for N' tagging of gene using pYM plasmids. AIM11
FMP16 N' tag CHK R	TTAATAAACACCGTCATCTGG	Primer for checking N' tagging of the gene. FMP16
FMP16 N' tag pYM F	AGTATCACATATATAATACACAG GAATATATTTGATAATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. FMP16
FMP16 N' tag pYM R	GCATCAATTGTCTTGGAGTGCGC AAAAAAGTGGTTCTCAAcatcgatgaa ttctgctg	Primer for N' tagging of gene using pYM plasmids. FMP16
FMP33 N' tag CHK R	CCAAACAATCGAAAATTGAG	Primer for checking N' tagging of the gene. FMP33
FMP33 N' tag pYM F	AAATAAATCAAGTATATCATAGA GTTCTTTCATTCATATGcgtacgctgca ggtcgac	Primer for N' tagging of gene using pYM plasmids. FMP33
FMP33 N' tag pYM R	TGGTGAATTGTGAGTTGTGACGT ACAACCTTGTGTATAGcatcgatgaat tctctgctg	Primer for N' tagging of gene using pYM plasmids. FMP33
FSF1 N' tag CHK R	TTTAGACCTAACGCAACACC	Primer for checking N' tagging of the gene. FSF1
FSF1 N' tag pYM F	CAGCATCGAAAGAAAAAATAA ACAAGGCAGAGAAATATGcgtacgct gcaggtcgac	Primer for N' tagging of gene using pYM plasmids. FSF1
FSF1 N' tag pYM R	GGGATTCGGGCAAAATCGATGGGC CCTGGGACTGATGATGCcatcgatgaa ttctgctg	Primer for N' tagging of gene using pYM plasmids. FSF1
FYV4 N' tag CHK R	AATGCCTTCTCTTCCTCTC	Primer for checking N' tagging of the gene. FYV4
FYV4 N' tag pYM F	AAACAATAACAAACCTTCATTCA ACACACGTTTTACTATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. FYV4
FYV4 N' tag pYM R	TTAAGAATAGAGGAAATTTGTGG CTAATCCGGGAAGGTATcatcgatgaa ttctgctg	Primer for N' tagging of gene using pYM plasmids. FYV4
LCL3 N' tag CHK R	TTCAAGAACTGGCTTTTAGC	Primer for checking N' tagging of the gene. LCL3
LCL3 N' tag pYM F	TTTTTCATAGCTATTAAGGGGGC ATAAATGCACTTATATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. LCL3
LCL3 N' tag pYM R	CAACATCCGCAGATTTTTTTGAAT TAGAATCACCTTCCCTcatcgatgaattc tctgctg	Primer for N' tagging of gene using pYM plasmids. LCL3

MNE1 N' tag CHK R	AGTGCAATTTTGTCAATTTGC	Primer for checking N' tagging of the gene. MNE1
MNE1 N' tag pYM F	CGGCGAGAAAAAATGATAGTAGT GTGCCAAGAAGAATATGcgtacgctg caggtcgac	Primer for N' tagging of gene using pYM plasmids. MNE1
MNE1 N' tag pYM R	TACCAATATGACTAGACGAATAT CTTTTAAAAAGTAACTTcatcgaatgaat tctctgctg	Primer for N' tagging of gene using pYM plasmids. MNE1
MSS116 N' tag CHK R	AACATCATGGTCTTCACTGG	Primer for checking N' tagging of the gene. MSS116
MSS116 N' tag pYM F	GGCAAGAAAATACCAGAGTTGCA CGTTAGGCTGATAAATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. MSS116
MSS116 N' tag pYM R	CAAGAACAGGTGTGCGACCTTTT ATCAATATAGAGGTCAAcacatcgaatgaat tctctgctg	Primer for N' tagging of gene using pYM plasmids. MSS116
SFH5 N' tag CHK R	GTGACAGCCTTTTTATTTGC	Primer for checking N' tagging of the gene. SFH5
SFH5 N' tag pYM F	CCAATCGTAATTAATTCATAAAA TTAACATCCTTAAAATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. SFH5
SFH5 N' tag pYM R	GCTTATCGAAAACCTGCTTCTCA CTGTCATTGTGCAATTTcatcgaatgaat ctctgctg	Primer for N' tagging of gene using pYM plasmids. SFH5
TOM5 N' tag CHK R	ATTTCCATTGCTTTTTTCACC	Primer for checking N' tagging of the gene. TOM5
TOM5 N' tag pYM F	AGGCGTCCATTGGCATCAAATAA CTAGATAGTATAAAAATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. TOM5
TOM5 N' tag pYM R	TTTTCTCCTCTTCGGAGACTTCTT GTTGAGGTAGACCAAACatcgaatgaat ctctgctg	Primer for N' tagging of gene using pYM plasmids. TOM5
YBL059W N' tag CHK R	ACCTGCAAGACTCTTTTGTG	Primer for checking N' tagging of the gene. YBL059W
YBL059W N' tag pYM F	ATCTTGCTTTTGTTCAACTGCACT TGTAATCAGTGAATGcgtacgctgca gggtcgac	Primer for N' tagging of gene using pYM plasmids. YBL059W
YBL059W N' tag pYM R	TAGAGAAAGACCTATTA AAAAAT CGCTGCGGTCCAATAAcacatcgaatgaat tctctgctg	Primer for N' tagging of gene using pYM plasmids. YBL059W
YGL041W-A N' tag CHK R	CGTGTCTCAGATCTCTGTCC	Primer for checking N' tagging of the gene. YGL041W-A
YGL041W-A N' tag pYM F	CACAAGAAGCAGAAAGTAGATA CTACAGCCACACATAATGcgtacgct gcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YGL041W-A
YGL041W-A N' tag pYM R	AACGAGTGA CTCTGGA ACTATGC TTCCATATAACTCTTAGcgtacgaatgaat ctctgctg	Primer for N' tagging of gene using pYM plasmids. YGL041W-A
YIL077C N' tag CHK R	GTA CTGCCTTTGATTTCTCG	Primer for checking N' tagging of the gene. YIL077C
YIL077C N' tag pYM F	TTAAGGAGCAGCCTAATTTCAAG CGTAGAACAAGGTGATGcgtacgctg caggtcgac	Primer for N' tagging of gene using pYM plasmids. YIL077C
YIL077C N' tag pYM R	TACCATTCTGGCCATATTGCTGTT	Primer for N' tagging of gene using

	CTTCTCTTTTTCCCAAcatcgatgaattctctgtcg	pYM plasmids. YJL077C
YJL127C-B N' tag CHK R	TACTGCACGTATTTTCGATTG	Primer for checking N' tagging of the gene. YJL127C-B
YJL127C-B N' tag pYM F	TGCAGCAGTGTACAGTTTACGCGACAATAAAGAAAGCATGcgtacgctgcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YJL127C-B
YJL127C-B N' tag pYM R	AAGCTGTAAATATTGAACGGATCTGGTTGAAAAAGAAAATcatcgatgaattctgtcg	Primer for N' tagging of gene using pYM plasmids. YJL127C-B
YJL133C-A N' tag CHK R	CCATATGTCTTGGCATTTC	Primer for checking N' tagging of the gene. YJL133C-A
YJL133C-A N' tag pYM F	AATATTACATTCATAAGACAGTAATAAACGTATTAATGcgtacgctgcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YJL133C-A
YJL133C-A N' tag pYM R	AAGAAGAGACGGCGGGCTAACTCTGGTACTTTGGGCGATcatcgatgaattctgtcg	Primer for N' tagging of gene using pYM plasmids. YJL133C-A
YJR085C N' tag CHK R	AGGGTAAAATTCCTTGTATTTG	Primer for checking N' tagging of the gene. YJR085C
YJR085C N' tag pYM F	AAACGAAGTTTGAGAGAAAGGAACATCAGATCACACTATGcgtacgctgcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YJR085C
YJR085C N' tag pYM R	CGGCTGTAGTCAAAGACTCAATGTATATGCTGGATGTTcctcgatgaattctgtcg	Primer for N' tagging of gene using pYM plasmids. YJR085C
YKL018C-A N' tag CHK R	AGATTGATCTCTCCCCTTC	Primer for checking N' tagging of the gene. YKL018C-A
YKL018C-A N' tag pYM F	AAAAGAGAGTGAAGATCAGATCGTAACATATTGCAAGATGcgtacgctgcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YKL018C-A
YKL018C-A N' tag pYM R	CTACCAACGTCCCCTCTACTACCCACCTTATCATTCCCAAcatcgatgaattctgtcg	Primer for N' tagging of gene using pYM plasmids. YKL018C-A
YKL044W N' tag CHK R	TATAGGCTGGGAAGTGGAGG	Primer for checking N' tagging of the gene. YKL044W
YKL044W N' tag pYM F	CTGTGGCCCAGGTAGCGTAGGCAAAATCAAGCTCAGAATGcgtacgctgcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YKL044W
YKL044W N' tag pYM R	CTGACATCCTCGCACTCGAAAACGTCATAATTACGTAACCcatcgatgaattctgtcg	Primer for N' tagging of gene using pYM plasmids. YKL044W
YKL133C N' tag CHK R	AAAGTAGATCCACGTTTTTCG	Primer for checking N' tagging of the gene. YKL133C
YKL133C N' tag pYM F	AAAGATATTGTAATGAAGAAATTATGCTCATTTAGTATGcgtacgctgcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YKL133C
YKL133C N' tag pYM R	CGGTCCCTTCATTTTTGACTGATCTATGAAGTATTTCCAcatcgatgaattctgtcg	Primer for N' tagging of gene using pYM plasmids. YKL133C
YLR281C N' tag CHK R	CGTTCCTCTTTTTCGACTC	Primer for checking N' tagging of the gene. YLR281C
YLR281C N' tag pYM F	CTGGAAAAAGCGCAGAAGAATCCGAGACTAGTAAACTATGcgtacgct	Primer for N' tagging of gene using pYM plasmids. YLR281C

	gcaggtcgac	
YLR281C N' tag pYM R	CTGCGGCACTGCTGATTGACCTC TFACTCGCTCCCCTCATcatcgaatgaatt ctctgtcg	Primer for N' tagging of gene using pYM plasmids. YLR281C
YMR252C N' tag CHK R	CTGGGGTGACTGAACCTAC	Primer for checking N' tagging of the gene. YMR252C
YMR252C N' tag pYM F	AAGCAGCAAAAACGAAGAAGATC GAGTGTGAAGTGTAAATGcgtacgct gcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YMR252C
YMR252C N' tag pYM R	CAATTCTTGTCTGATGTAGCTTA CGAAAACCTTACCAAcatcgaatgaatt ctctgtcg	Primer for N' tagging of gene using pYM plasmids. YMR252C
YNR040W N' tag CHK R	CTCCCTCGTGATAGAAACTG	Primer for checking N' tagging of the gene. YNR040W
YNR040W N' tag pYM F	AAGAAGGTGCATTTAGTTATCTT ATGTGGACTAGGGTATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. YNR040W
YNR040W N' tag pYM R	AGCTGCTTCCTTTGAACTGATTCT TGGCAGCCATGTTTGTcatcgaatgaatt tctgtcg	Primer for N' tagging of gene using pYM plasmids. YNR040W
YPL041C N' tag CHK R	CCAACAAATTTCTCCATAGC	Primer for checking N' tagging of the gene. YPL041C
YPL041C N' tag pYM F	GGATAAGTGTTACACCCAAGGCA CCCTCACCAGGAACATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. YPL041C
YPL041C N' tag pYM R	TCTGCAGTTGACAGGGCCTGAAG AAAAGATTCATGACAGTcatcgaatgaatt ttctgtcg	Primer for N' tagging of gene using pYM plasmids. YPL041C
YPL109C N' tag CHK R	TTGGTGTCCGTAATTTTGGAG	Primer for checking N' tagging of the gene. YPL109C
YPL109C N' tag pYM F	CAGAATAGAGATAAAGAACATC AGAACCATCTGGGCAATGcgtacgct gcaggtcgac	Primer for N' tagging of gene using pYM plasmids. YPL109C
YPL109C N' tag pYM R	AGTATCGCCAAGAGTTTCTATAA GCGAACTTTAAAAATGAcacgaatgaatt ttctgtcg	Primer for N' tagging of gene using pYM plasmids. YPL109C
YPR098C N' tag CHK R	CGACTTGCCAAATTCCTTAC	Primer for checking N' tagging of the gene. YPR098C
YPR098C N' tag pYM F	ACTACGCTTTACAGTCCAGTAAA CTTAACAACGAAAAATGcgtacgctgc aggtcgac	Primer for N' tagging of gene using pYM plasmids. YPR098C
YPR098C N' tag pYM R	AGGAATAAAAAAGCAAATGAGC CGTAGTTTTGACTAAACAcacgaatgaatt ttctgtcg	Primer for N' tagging of gene using pYM plasmids. YPR098C
S4 reverse complement	cgacagagaattcatcgaat	Primer for checking N' tagging of the genes.

Yeast strains used in this study

Strain	Description	Genotype	Source or Reference	Number
Wild-type (WT)	BY4741	<i>MATa; ura3Δ0, leu2Δ0, his3Δ1, met15Δ0</i>	Brachmann et al., 1998	1354
Wild-type (WT)	BY4742 Δ <i>arg4</i>	<i>MATa; ura3Δ0, leu2Δ0, his3Δ1, lys2Δ0, arg4::kanMX4</i>	Euroscarf	2708
Wild-type (WT)	BY4741 Δ <i>lys2</i> Δ <i>arg4</i>	<i>MATa; ura3Δ0, leu2Δ0, his3Δ1, lys2Δ0, met15Δ0, arg4::kanMX4</i>	this study	3559
Wild-type (WT)	YPH499	<i>MATa; ura3-52, lys2-801, ade2-101, trp1-Δ63, his3-Δ200, leu2-Δ1</i>	Sikorski and Hieter, 1989	1501
Wild-type (WT)	YPH499 Δ <i>arg4</i>	<i>MATa; ura3-52, lys2-801, ade2-101, trp1-Δ63, his3-Δ200, leu2-Δ1, arg4::kanMX4</i>	von der Malsburg et al., 2011	2799
Wild-type (WT)	YPH499 pRS425 (empty)	YPH499 + pRS425	this study	4312
Aim11 _{ProtA}	Aim11 _{ProtA} (chromosomal)	BY4741 <i>aim11::AIM11-TEV-ProtA-7HIS-kanMX6</i>	this study	4950
Deg1 _{ProtA}	Deg1 _{ProtA} (chromosomal)	BY4741 <i>deg1::DEG1-TEV-ProtA-7HIS-hphNT1</i>	this study	5044
Ecm4 _{ProtA}	Ecm4 _{ProtA} (chromosomal)	BY4741 <i>ecm4::ECM4-TEV-ProtA-7HIS-hphNT1</i>	this study	5045
Ecm19 _{ProtA}	Ecm19 _{ProtA} (chromosomal)	BY4741 <i>ecm19::ECM19-TEV-ProtA-7HIS-hphNT1</i>	this study	4951
Fmp16 _{ProtA}	Fmp16 _{ProtA} (chromosomal)	BY4741 <i>fmp16::FMP16-TEV-ProtA-7HIS-kanMX6</i>	this study	4952
Fyv4 _{ProtA}	Fyv4 _{ProtA} (chromosomal)	BY4741 <i>fyv4::FYV4-TEV-ProtA-7HIS-hphNT1</i>	this study	4953
Meo1 _{GFP}	Meo1 _{GFP} (chromosomal)	BY4741 <i>meo1::MEO1-γEGFP-kanMX4</i>	this study	4113
Meu1 _{ProtA}	Meu1 _{ProtA} (chromosomal)	BY4741 <i>meu1::MEU1-TEV-ProtA-7HIS-hphNT1</i>	this study	5046
Mlo1 _{ProtA}	Mlo1 _{ProtA} (chromosomal)	BY4741 <i>mlo1::MLO1-TEV-ProtA-7HIS-hphNT1</i>	this study	4954
Mtc3 _{ProtA}	Mtc3 _{ProtA} (chromosomal)	BY4741 <i>mtc3::MTC3-TEV-ProtA-7HIS-hphNT1</i>	this study	4955
Nce101 _{HA}	Nce101 _{HA}	BY4741 <i>nce101::NCE101-3HA-</i>	this study	4956

	(chromosomal)	<i>kanMX6</i>		
Pxp2 _{ProtA}	Pxp2 _{ProtA} (chromosomal)	BY4741 <i>pxp2::PXP2-TEV-ProtA-7HIS-hphNT1</i>	this study	5047
Rdl2 _{ProtA}	Rdl2 _{ProtA} (chromosomal)	BY4741 <i>rdl2::RDL2-TEV-ProtA-7HIS-kanMX6</i>	this study	4957
Rrg9 _{ProtA}	Rrg9 _{ProtA} (chromosomal)	BY4741 <i>rrg9::RRG9-TEV-ProtA-7HIS-kanMX6</i>	this study	4958
Smm1 _{ProtA}	Smm1 _{ProtA} (chromosomal)	BY4741 <i>smm1::SMM1-TEV-ProtA-7HIS-hphNT1</i>	this study	5048
Sua5 _{ProtA}	Sua5 _{ProtA} (chromosomal)	BY4741 <i>sua5::SUA5-TEV-ProtA-7HIS-hphNT1</i>	this study	5049
Tum1 _{ProtA}	Tum1 _{ProtA} (chromosomal)	BY4741 <i>tum1::TUM1-TEV-ProtA-7HIS-hphNT1</i>	this study	5050
Ybl039w-b _{GFP}	Ybl039w-b _{GFP} (chromosomal)	BY4741 <i>ybl039w-b::YBL039W-B-γEGFP-kanMX4</i>	this study	4112
Ybl039w-b _{HA}	Ybl039w-b _{HA} (chromosomal)	BY4741 <i>ybl039w-b::YBL039W-B-3HA-kanMX6</i>	this study	4983
Ybl059w _{ProtA}	Ybl059w _{ProtA} (chromosomal)	BY4741 <i>ybl059w::YBL059W-TEV-ProtA-7HIS-hphNT1</i>	this study	4959
Ybr201c-a _{ProtA}	Ybr201c-a _{ProtA} (chromosomal)	BY4741 <i>ybr201c-a::YBR201C-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4960
Ybr230w-a _{ProtA}	Ybr230w-a _{ProtA} (chromosomal)	BY4741 <i>ybr230w-a::YBR230W-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4961
Ydr286c _{ProtA}	Ydr286c _{ProtA} (chromosomal)	BY4741 <i>ydr286c::YDR286C-TEV-ProtA-7HIS-kanMX6</i>	this study	4962
Ydr381c-a _{ProtA}	Ydr381c-a _{ProtA} (chromosomal)	BY4741 <i>ydr381c-a::YDR381C-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4970
Ydr461c-a _{ProtA}	Ydr461c-a _{ProtA} (chromosomal)	BY4741 <i>ydr461c-a::YDR461C-A-TEV-ProtA-7HIS-hphNT1</i>	this study	4971
Yfr032c-b _{GFP}	Yfr032c-b _{GFP} (chromosomal)	BY4741 <i>yfr032c-b::YFR032C-B-γEGFP-kanMX4</i>	this study	4118
Yfr032c-b _{ProtA}	Yfr032c-b _{ProtA} (chromosomal)	BY4741 <i>Yfr032c-b::YFR032C-B-TEV-ProtA-7HIS-kanMX6</i>	this study	5054
Ygl041w-a _{ProtA}	Ygl041w-a _{ProtA} (chromosomal)	BY4741 <i>ygl041w-a::YGL041W-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4972

Ygl204c _{GFP}	Ygl204c _{GFP} (chromosomal)	BY4741 <i>Ygl204c::YGL204C-γEGFP-kanMX4</i>	this study	4117
Ygr021w _{ProtA}	Ygr021w _{ProtA} (chromosomal)	BY4741 <i>ygr021w::YGR021W-TEV-ProtA-7HIS-hphNT1</i>	this study	4973
Ygr053c _{ProtA}	Ygr053c _{ProtA} (chromosomal)	BY4741 <i>ygr053c::YGR053C-TEV-ProtA-7HIS-hphNT1</i>	this study	4974
Yhl018w _{ProtA}	Yhl018w _{ProtA} (chromosomal)	BY4741 <i>yhl018w::YHL018W-TEV-ProtA-7HIS-kanMX6</i>	this study	4975
Yil002w-a _{ProtA}	Yil002w-a _{ProtA} (chromosomal)	BY4741 <i>yil002w-a::YIL002W-A-TEV-ProtA-7HIS-hphNT1</i>	this study	4976
<i>yil077cΔ</i>	Deletion of <i>YIL007C</i> in BY4741 yeast cells	BY4741 <i>yil077c::kanMX4</i>	Euroscarf	3407
Yil077c _{ProtA}	Yil077c _{ProtA} (chromosomal)	BY4741 <i>yil077c::YIL077C-TEV-ProtA-7HIS-kanMX6</i>	this study	4977
Yil156w-b _{ProtA}	Yil156w-b _{ProtA} (chromosomal)	BY4741 <i>yil156w-b::YIL156W-B-TEV-ProtA-7HIS-kanMX6</i>	this study	4978
Yjl133c-a _{ProtA}	Yjl133c-a _{ProtA} (chromosomal)	BY4741 <i>yjl133c-a::YJL133C-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4979
^{HA} YJR085C (YPH499)	YPH499 <i>pRS425-^{HA}YJR085C</i>	YPH499 + <i>pRS425-^{HA}YJR085C</i>	this study	4378
Yjr085c _{ProtA}	Yjr085c _{ProtA} (chromosomal)	BY4741 <i>yjr085c::YJR085C-TEV-ProtA-7HIS-kanMX6</i>	this study	4980
Ykl023c-a _{GFP}	Ykl023c-a _{GFP} (chromosomal)	BY4741 <i>ykl023c-a::YKL023C-A-γEGFP-kanMX4</i>	this study	4114
Ykl023c-a _{HA}	Ykl023c-a _{HA} (chromosomal)	BY4741 <i>ykl023c-a::YKL023C-A-3HA-kanMX6</i>	this study	4108
Ykl065w-a _{GFP}	Ykl065w-a _{GFP} (chromosomal)	BY4741 <i>ykl065w-a::YKL065W-A-γEGFP-kanMX4</i>	this study	4115
Ykl065w-a _{ProtA}	Ykl065w-a _{ProtA} (chromosomal)	BY4741 <i>ykl065w-a::YKL065W-A-TEV-ProtA-7HIS-hphNT1</i>	this study	4981
Ykl133c _{ProtA}	Ykl133c _{ProtA} (chromosomal)	BY4741 <i>ykl133c::YKL133C-TEV-ProtA-7HIS-hphNT1</i>	this study	4982
Ylr049c _{ProtA}	Ylr049c _{ProtA} (chromosomal)	BY4741 <i>ylr049c::YLR049C-TEV-ProtA-7HIS-hphNT1</i>	this study	4984
Ylr118c _{ProtA}	Ylr118c _{ProtA} (chromosomal)	BY4741 <i>ylr118c::YLR118C-TEV-ProtA-7HIS-hphNT1</i>	this study	5051

Ylr281c _{ProtA}	Ylr281c _{ProtA} (chromosomal)	BY4741 <i>ylr281c::YLR281C-TEV-ProtA-7HIS-hphNT1</i>	this study	4984
Ylr307c-a _{ProtA}	Ylr307c-a _{ProtA} (chromosomal)	BY4741 <i>ylr307c-a::YLR307C-A-TEV-ProtA-7HIS-hphNT1</i>	this study	4986
Yml007c-a _{ProtA}	Yml007c-a _{ProtA} (chromosomal)	BY4741 <i>yml007c-a::YML007C-A-TEV-ProtA-7HIS-hphNT1</i>	this study	4987
Ymr087w _{ProtA}	Ymr087w _{ProtA} (chromosomal)	BY4741 <i>ymr087w::YMR087W-TEV-ProtA-7HIS-hphNT1</i>	this study	5052
Ymr130w _{ProtA}	Ymr130w _{ProtA} (chromosomal)	BY4741 <i>ymr130w::YMR130W-TEV-ProtA-7HIS-hphNT1</i>	this study	4988
Ymr182w-a _{GFP}	Ymr182w-a _{GFP} (chromosomal)	BY4741 <i>ymr182w-a::YMR182W-A-γEGFP-kanMX4</i>	this study	5055
Ynr040w _{ProtA}	Ynr040w _{ProtA} (chromosomal)	BY4741 <i>ynr040w::YNR040W-TEV-ProtA-7HIS-hphNT1</i>	this study	4989
Yor020w-a _{ProtA}	Yor020w-a _{ProtA} (chromosomal)	BY4741 <i>yor020w-a::YOR020W-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4990
Yor114w _{ProtA}	Yor114w-a _{ProtA} (chromosomal)	BY4741 <i>yor114w::YOR114W-TEV-ProtA-7HIS-kanMX6</i>	this study	5053
Ypl107w _{ProtA}	Ypl107w _{ProtA} (chromosomal)	BY4741 <i>ypl107w::YPL107W-TEV-ProtA-7HIS-hphNT1</i>	this study	4991
Ypr010c-a _{ProtA}	Ypr010c-a _{ProtA} (chromosomal)	BY4741 <i>ypr010c-a::YPR010C-A-TEV-ProtA-7HIS-kanMX6</i>	this study	4992
Ypr098c _{ProtA}	Ypr098c _{ProtA} (chromosomal)	BY4741 <i>ypr098c::YPR098C-TEV-ProtA-7HIS-hphNT1</i>	this study	4993
NOP1pr-sfGFP-Aim11	NOP1pr-sfGFP-Aim11 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Aim11	this study	
NOP1pr-sfGFP-Fsf1	NOP1pr-sfGFP-Fsf1 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Fsf1	this study	
NOP1pr-sfGFP-Mmo1	NOP1pr-sfGFP-Mmo1 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Mmo1	this study	
NOP1pr-sfGFP-Mrx11	NOP1pr-sfGFP-Mrx11 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Mrx11	this study	
NOP1pr-sfGFP-Tom5	NOP1pr-sfGFP-Tom5 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Tom5	Yofe et al., 2016	
NOP1pr-sfGFP-Ybl059w	NOP1pr-sfGFP-Ybl059w (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1	this study	

		pr-sfGFP-Ybl059w		
NOP1pr-sfGFP-Yil077c	NOP1pr-sfGFP-Yil077c (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Yil077c	this study	
NOP1pr-sfGFP-Yjl127c-b	NOP1pr-sfGFP-Yjl127c-b (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Yjl127c-b	this study	
NOP1pr-sfGFP-Yjr085c	NOP1pr-sfGFP-Yjr085c (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Yjr085c	this study	
NOP1pr-sfGFP-Ykl133c	NOP1pr-sfGFP-Ykl133c (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Ykl133c	this study	
NOP1pr-sfGFP-Ynr040w	NOP1pr-sfGFP-Ynr040w (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-Ynr040w	this study	
NOP1pr-sfGFP-YPR098C	NOP1pr-sfGFP-YPR098C (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 hphΔn::URA3::NOP1 pr-sfGFP-YPR098C	this study	
SFH5pr-sfGFP-Sfh5	SFH5pr-sfGFP-Sfh5 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; SFH5pr-sfGFP-Sfh5	this study	
FMP16pr-sfGFP-Fmp16	FMP16pr-sfGFP-Fmp16 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; FMP16pr-sfGFP-Fmp16	this study	
MSS116pr-sfGFP-Mss116	MSS116pr-sfGFP-Mss116 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; MSS116pr-sfGFP-Mss116	this study	
YGL041W-Apr-sfGFP-Ygl041w-a	YGL041W-Apr-sfGFP-Ygl041w-a (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; YGL041W-Apr-sfGFP-Ygl041w-a	this study	
FYV4pr-sfGFP-Fyv4	FYV4pr-sfGFP-Fyv4 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; FYV4pr-sfGFP-Fyv4	this study	
YKL018C-Apr-sfGFP-Ykl018c-a	YKL018C-Apr-sfGFP-Ykl018c-a (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; YKL018C-Apr-sfGFP-Ykl018c-a	this study	
FMP33pr-sfGFP-Fmp33	FMP33pr-sfGFP-Fmp33 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5	this study	

		lyp1Δ::STE3pr-LEU2 ; FMP33pr-sfGFP-Fmp33		
LCL3pr-sfGFP-Llc3	LCL3pr-sfGFP-Llc3 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; LCL3pr-sfGFP-Llc3	this study	
YLR281Cpr-sfGFP-Ylr281c	YLR281Cpr-sfGFP-Ylr281c (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; YLR281Cpr-sfGFP-Ylr281c	this study	
YMR252Cpr-sfGFP-Ymr252c	YMR252Cpr-sfGFP-Ymr252c (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; YMR252Cpr-sfGFP-Ymr252c	this study	
MNE1pr-sfGFP-Mne1	MNE1pr-sfGFP-Mne1 (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; MNE1pr-sfGFP-Mne1	this study	
YPL109Cpr-sfGFP-Ypl109c	YPL109Cpr-sfGFP-Ypl109c (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; YPL109Cpr-sfGFP-Ypl109c	this study	
YJL133C-Apr-sfGFP-Yjl133c-a	YJL133C-Apr-sfGFP-Yjl133c-a (chromosomal)	his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 lys+ can1Δ::GAL1pr-SceI::STE2pr-SpHIS5 lyp1Δ::STE3pr-LEU2 ; YJL133C-Apr-sfGFP-Yjl133c-a	this study	

Plasmids used in this study

Name	Description	Source or Reference	Number
pYM1	template for amplification of <i>3HA-kanMX6</i> cassette	Knop et al., 1999	1449
pYM9	template for amplification of <i>TEV-ProtA-7HIS-kanMX6</i> cassette	Knop et al., 1999	1457
pYM10	template for amplification of <i>TEV-ProtA-7HIS</i> cassette that was cloned into pFA6a-hphNT1 vector to generate pFA6a-TEV-ProtA-7His-hphNT1 plasmid.	Knop et al., 1999	1458
pYM12	template for amplification of γ <i>EGFP-kanMX4</i> cassette	Knop et al., 1999	1460
pFA6a-hphNT1	Target plasmid for TEV-ProtA-7His module (that has been amplified from pYM10 plasmid) for generation of pFA6a-TEV-ProtA-7His-hphNT1 plasmid.	Janke et al., 2004	2722
pFA6a-TEV-ProtA-7His-hphNT1	template for amplification of <i>TEV-ProtA-7HIS-hphNT1</i> cassette	this study	2723
pGEM-4Z-YJR085C	Backbone: pGEM-4Z Insert: EcoRI-YJR085C (<i>S. cerevisiae</i>)-HindIII template for in vitro synthesis of [³⁵ S]Yjr085c using the TnT® Quick Coupled Transcription/ Translation System	this study	2610
pRS425	Vector used for overexpression of proteins in <i>S. cerevisiae</i> . origin of replication: 2 μ yeast selectable marker: LEU2	Christianson et al., 1992	X30
pRS425-YJR085C	Backbone: pRS425 Insert: HindIII-P _{YJR085C} -YJR085C (<i>S. cerevisiae</i>)-T _{YJR085C} -BamHI	this study	2607
pRS425 _{-HA} -YJR085C	Backbone: pRS425 Insert: HindIII-P _{YJR085C} -1HA-YJR085C (<i>S. cerevisiae</i>)-T _{YJR085C} -BamHI	this study	2608
pST-N2	SWAT-GFP	Yofe et al., 2016	555
pSD-N9	Seamless-GFP	Yofe et al., 2016	561

Antibodies used in this study

Antigen	Dilution	Number	Secondary antibody
Abf2	1:200 TBS-T + 5% milk	GR B2072	anti-rabbit
Aco1	1:1000 TBS-T + 5% milk	GR945-7	anti-rabbit
Afg3	1:500 TBS-T + 5% milk	GR1551-5	anti-rabbit
Atp2	1:50 TBS-T + 5% milk	GR863 affinity purified e4	anti-rabbit
Atp4	1:250 TBS-T + 5% milk	GR1970-6	anti-rabbit
Atp5	1:250 TBS-T + 5% milk	GR1546-4	anti-rabbit
Atp17	1:250 TBS-T + 5% milk	GR1968-3	anti-rabbit
Atp20	1:250 TBS-T + 5% milk	GR1517-5	anti-rabbit
Cdc48	1:250 TBS-T + 5% milk	GR5015-4	anti-rabbit
Cor1	1:300 TBS-T + 5% milk	GR371-5	anti-rabbit
Cox1	1:400 TBS-T + 5% milk	GR1538-4	anti-rabbit
Cox2	1:250 TBS-T + 5% milk	GR1949-2	anti-rabbit
Cox4	1:1000 TBS-T + 5% milk	GR578-5	anti-rabbit
Cox5a	1:400 TBS-T + 5% milk	GR1540-5	anti-rabbit
Cox6	1:250 TBS-T + 5% milk	GR2015-2	anti-rabbit
Cox8	1:300 TBS-T + 5% milk	GR3609-5	anti-rabbit
Cox9	1:250 TBS-T + 5% milk	GR3612-3	anti-rabbit
Cox13	1:250 TBS-T + 5% milk	GR1542-4	anti-rabbit
Cta1	1:20000 TBS-T + 5% milk	12	anti-goat
Cyt1	1:750 TBS-T + 5% milk	GR541-6	anti-rabbit
Dic1	1:200 TBS-T + 5% milk	GR2054-5	anti-rabbit
Erg6	1:500 TBS-T + 5% milk	GR3034-6	anti-rabbit
Fmp10	1:500 TBS-T + 5% milk	GR3338-2	anti-rabbit
GFP	1:1000 TBS-T + 5% milk	GFP Antibody (Novus Biologicals LLC; NB600-308)	anti-rabbit
Goat IgG	1:50000 TBS-T + 5% milk	Peroxidase antibody, rabbit (Sigma-Aldrich Corp.; A8919)	—
HA	1:1000 TBS	Anti-HA-Peroxidase (F. Hoffmann-La Roche AG; 11667475001)	—
Isd11	1:250 TBS-T + 5% milk	336-7	anti-rabbit
Mdj1	1:500 TBS-T + 5% milk	GR1839-6	anti-rabbit
Mdl1	1:200 TBS-T + 5% milk	GR1518-7	anti-rabbit
Mdm38	1:1000 TBS-T + 5% milk	342-6	anti-rabbit
Mge1	1:250 TBS-T + 5% milk	GR1838-6	anti-rabbit
Mgr2	1:250 TBS-T + 5% milk	GR3121-3/4	anti-rabbit
Mia40	1:750 TBS-T + 5% milk	B315	anti-rabbit
Mic10	1:500 TBS-T + 5% milk	GR3343-2	anti-rabbit
Mic19	1:400 TBS-T + 5% milk	GR3358-2	anti-rabbit
Mic26	1:250 TBS-T + 5% milk	GR3335-2	anti-rabbit
Mic27	1:250 TBS-T + 5% milk	GR3357-2	anti-rabbit

Mic60	1:500 TBS-T + 5% milk	GR857-5	anti-rabbit
Ndi1	1:200 TBS-T + 5% milk	GR809-4	anti-rabbit
Ola1	1:1000 TBS-T + 5% milk	7173	anti-rabbit
OM14	1:500 TBS-T + 5% milk	GR3040-1	anti-rabbit
Pam16	1:200 TBS-T + 5% milk	GR750-6	anti-rabbit
Pam18	1:500 TBS-T + 5% milk	GR752-3	anti-rabbit
Pdi1	1:2000 TBS-T + 5% milk	GR1946	anti-rabbit
Pgk1	1:5000 TBS-T + 5% milk	GR753-1	anti-rabbit
Phb2	1:300 TBS-T + 5% milk	B295-10	anti-rabbit
Por1	1:500 TBS-T + 5% milk	GR3622-3	anti-rabbit
ProtA	1:200 TBS + 5% milk	Peroxidase Anti-Peroxidase Soluble Complex antibody produced in rabbit (Sigma-Aldrich Corp.; P1291)	—
Pth2	1:500 TBS-T + 5% milk	GR797-3	anti-rabbit
Qcr6	1:250 TBS-T + 5% milk	GR1054-6	anti-rabbit
Qcr8	1:5000 TBS-T + 5% milk	GR1038-1	anti-rabbit
Rcf2	1:2000 TBS-T + 5% milk	GR3113-1	anti-rabbit
Rip1	1:500 TBS-T + 5% milk	GR543-5	anti-rabbit
Rpl19	1:7000 TBS-T + 5% milk	106	anti-rabbit
Sac1	1:2500 TBS-T + 5% milk	GR1487-6	anti-rabbit
Sam50	1:500 TBS-T + 5% milk	B312-14	anti-rabbit
Scm4	1:500 TBS-T + 5% milk	GR1473-2	anti-rabbit
Sdh1	1:1000 TBS-T + 5% milk	GR1849-3	anti-rabbit
Sdh4	1:2000 TBS-T + 5% milk	GR1855-3	anti-rabbit
Sec61	1:1000 TBS-T + 5% milk	GR759-2	anti-rabbit
Ssa1	1:1000 TBS-T + 5% milk	GR1011-4	anti-rabbit
Sss1	1:100 TBS-T + 5% milk	GR787-7 / GR788-1	anti-rabbit
Tcd2	1:500 TBS-T + 5% milk	GR1396-1	anti-rabbit
Tim10	1:500 TBS-T + 5% milk 1:250 TBS-T + 5% milk	217-8 GR2041-7	anti-rabbit
Tim11	1:400 TBS-T + 5% milk	138-9	anti-rabbit
Tim12	1:250 TBS-T + 5% milk	GR906-7	anti-rabbit
Tim17	1:300 TBS-T + 5% milk	GR1845-3	anti-rabbit
Tim21	1:500 TBS-T + 5% milk	GR3899-4	anti-rabbit
Tim23	1:500 TBS-T + 5% milk	GR3878-4	anti-rabbit
Tim44	1:200 TBS-T + 5% milk	GR1836-4	anti-rabbit
Tim50	1:500 TBS-T + 5% milk	GR3881-1	anti-rabbit
Tim54	1:1000 TBS-T + 5% milk	215-6	anti-rabbit
Tom7	1:250 TBS-T + 5% milk	230-9	anti-rabbit
Tom20	1:5000 TBS-T + 5% milk	GR3225	anti-rabbit
Tom22	1:5000 TBS-T + 5% milk	GR3227	anti-rabbit
Tom40	1:500 TBS-T + 5% milk	168-4	anti-rabbit
Tom70	1:500 TBS-T + 5% milk	GR657-4	anti-rabbit
Yil077c (Rci37)	1:250 TBS-T + 5% milk	GR3731-5	anti-rabbit

Yme1	1:400 TBS-T + 5% milk	GR1435-3	anti-rabbit
------	-----------------------	----------	-------------

SUPPLEMENTAL REFERENCES

- Brachmann, C.B., Davies, A., Cost, G.J., Caputo, E., Li, J., Hieter, P., and Boeke, J.D. (1998). Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* *14*, 115–132.
- Christianson, T.W., Sikorski, R.S., Dante, M., Shero, J.H., and Hieter, P. (1992). Multifunctional yeast high-copy-number shuttle vectors. *Gene* *110*, 119–122.
- Cox, J., and Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* *26*, 1367–1372.
- Cox, J., Neuhauser, N., Michalski, A., Scheltema, R.A., Olsen, J.V., and Mann, M. (2011). Andromeda: a peptide search engine integrated into the MaxQuant environment. *J. Proteome Res.* *10*, 1794–1805.
- Delmotte, N., Lasasoa, M., Tholey, A., Heinzle, E., and Huber, C.G. (2007). Two-dimensional reversed-phase x ion-pair reversed-phase HPLC: an alternative approach to high-resolution peptide separation for shotgun proteome analysis. *J. Proteome Res.* *6*, 4363–4373.
- Dubaquié, Y., Looser, R., Fünfschilling, U., Jenö, P., Rospert, S. (1998). Identification of in vivo substrates of the yeast mitochondrial chaperonins reveals overlapping but non-identical requirement for hsp60 and hsp10. *EMBO J.* *17*, 5868–5876.
- Fin, R.D., Clements, J., Arndt, W., Miller, B.L., Wheeler, T.J., Schreiber, F., Bateman, A., and Eddy, S.R. (2015). HMMER web server: 2015 update. *Nucleic Acids Res.* *43*, W30–W38.
- Fukasawa, Y., Tsuji, J., Fu, S.-C., Tomii, K., Horton, P., and Imai, K. (2015). MitoFates: improved prediction of mitochondrial targeting sequences and their cleavage sites. *Mol. Cell. Proteomics* *14*, 1113–1126.
- Gietz, R.D., and Woods, R.A. (2002). Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Meth. Enzymol.* *350*, 87–96.
- Huxley, C., Green, E.D., and Dunham, I. (1990). Rapid assessment of *S. cerevisiae* mating type by PCR. *Trends Genet.* *6*, 236.
- Janke, C., Magiera, M.M., Rathfelder, N., Taxis, C., Reber, S., Maekawa, H., Moreno-Borchart, A., Doenges, G., Schwob, E., Schiebel, E., et al. (2004). A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* *21*, 947–962.
- Knop, M., Siegers, K., Pereira, G., Zachariae, W., Winsor, B., Nasmyth, K., and Schiebel, E. (1999). Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical routines. *Yeast* *15*, 963–972.
- Kushnirov, V.V. (2000). Rapid and reliable protein extraction from yeast. *Yeast* *16*, 857–860.
- Lasasoa, M., Delmotte, N., Huber, C.G., Melchior, K., Heinzle, E., and Tholey, A. (2009). A 2D reversed-phase x ion-pair reversed-phase HPLC-MALDI TOF/TOF-MS approach for shotgun proteome analysis. *Anal. Bioanal. Chem.* *393*, 1245–1256.
- Longtine, M.S., McKenzie, A., Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., and Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* *14*, 953–961.
- Meisinger, C., Sommer, T., and Pfanner, N. (2000). Purification of *Saccharomyces cerevisiae* mitochondria devoid of microsomal and cytosolic contaminations. *Anal. Biochem.* *287*, 339–342.
- Sikorski, R.S., and Hieter, P. (1989). A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* *122*, 19–27.
- Smith, J.E., Alvarez-Dominguez, J.R., Kline, N., Huynh, N.J., Geisler, S., Hu, W., Collier, J., Baker, K.E. 2014. Translation of Small Open Reading Frames within Unannotated RNA Transcripts in *Saccharomyces cerevisiae*. *Cell Rep.* *7*, 1858–1866.
- Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and Cox, J. (2016). The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods* *13*, 731–740.
- von der Malsburg, K., Müller, J.M., Bohnert, M., Oeljeklaus, S., Kwiatkowska, P., Becker, T., Loniewska-Lwowska, A., Wiese, S., Rao, S., Milenkovic, D., et al. (2011). Dual role of mitofilin in mitochondrial membrane organization and protein biogenesis. *Dev. Cell* *21*, 694–707.
- Wiśniewski, J.R., Ostasiewicz, P., Duś, K., Zielinska, D.F., Gnäd, F., and Mann, M. (2012). Extensive quantitative remodeling of the proteome between normal colon tissue and adenocarcinoma. *Mol. Syst. Biol.* *8*, 611.
- Wysocki, R., Roganti, T., Van Dyck, E., de Kerchove D'Exaerde, A., Foury, F. 1999. Disruption and basic phenotypic analysis of 18 novel genes from the yeast *Saccharomyces cerevisiae*. *Yeast* *15*, 165–171.