

Supplementary Table 1. Adjusted ORs for pancreatic cancer for a unit increment of the WCRF/AICR score, by strata of selected covariates. Italy 1991-2008.

	Cases/controls	Adjusted OR (95% CI) <sup>a</sup>
Sex		
Males	174/348	0.67 (0.52-0.86)
Females	152/304	0.80 (0.61-1.07)
Age (years)		
<60	121/242	0.90 (0.68-1.19) <sup>c</sup>
≥60	205/410	0.62 (0.49-0.79) <sup>c</sup>
Education (years) <sup>b</sup>		
<7	166/350	0.78 (0.61-1.00)
≥7	158/300	0.66 (0.51-0.87)
Body mass index (kg/m <sup>2</sup> )		
<25	139/264	0.73 (0.60-0.88)
≥25	185/385	0.67 (0.55-0.83)
Tobacco smoking <sup>b</sup>		
Never smoker	137/328	0.80 (0.61-1.05)
Ex smoker	86/189	0.56 (0.39-0.80)
Current smoker	102/133	0.79 (0.56-1.12)
Diabetes		
No	279/615	0.76 (0.62-0.92)
Yes	47/37	0.51 (0.29-0.88)

AICR: American Institute of Cancer Research; OR: odds ratio; CI: confidence interval; WCRF: World Cancer Research Fund.

<sup>a</sup> Covariates include: sex, study center, year of interview, age, education, body mass index, tobacco smoking, and history of diabetes.

<sup>b</sup> The sum is not the total number of subjects due to missing values.

<sup>c</sup> p-value for heterogeneity between strata <0.05.