

Additional table 2. Relative mRNA concentrations of NK cell receptors and cytokines in spleen of rats in short-term experiment

	<i>activating NK cell receptors</i>			<i>cytokines</i>
Groups	NCR1/NKp46	NCR3/NKp30	Klrk1/NKG2D	TNFα
One-way ANOVA	<i>relative mRNA concentrations</i>			
control/NaCl	1.0 ± 0.1	1.0 ± 0.1	1.0 ± 0.05	1.0 ± 0.2
DIO/NaCl	0.9 ± 0.1	1.1 ± 0.1	0.8 ± 0.1	0.7 ± 0.1
control/MADB106	1.1 ± 0.1	1.1 ± 0.05	1.1 ± 0.1	1.0 ± 0.1
DIO/MADB106	1.1 ± 0.1	1.4 ± 0.3	1.0 ± 0.05	1.1 ± 0.1
Two-way ANOVA	<i>p</i>			
diet	0.53	0.26	0.21	0.64
MADB106	0.15	0.34	0.09	0.17
diet x MADB106	0.51	0.52	0.33	0.10

Values represent means ± SEM, n=8 rats/group for one-way ANOVA. For two-way ANOVA p-values are shown for the main factors diet, MADB106 and the interaction of both main factors.