

**Additional table 3.** Relative mRNA concentrations of NK cell receptors and cytokines in spleen of rats in long-term experiment

	<i>NK cell receptors</i>		<i>cytokines</i>	
	<i>activating</i>	<i>inhibiting</i>		
<b>Groups</b>	<b>NCR3/NKp30</b>	<b>Klra1/Ly49</b>	<b>TNF<math>\alpha</math></b>	<b>TNFSF10/TRAIL</b>
<b>One-way ANOVA</b>	<i>relative mRNA concentrations</i>			
control/NaCl	1.0 ± 0.1	1.0 ± 0.05	1.0 ± 0.1	1.0 ± 0.1
DIO/NaCl	1.3 ± 0.3	0.9 ± 0.1	1.5 ± 0.3	1.5 ± 0.5
control/MADB106	0.8 ± 0.1	1.0 ± 0.1	1.0 ± 0.2	0.9 ± 0.1
DIO/MADB106	1.1 ± 0.1	0.9 ± 0.1	1.0 ± 0.1	1.0 ± 0.2
<b>Two-way ANOVA</b>	<i>p</i>			
diet	0.11	0.20	0.21	0.22
MADB106	0.21	0.79	0.22	0.26
diet x MADB106	0.74	0.89	0.29	0.38

Values represent means ± SEM, n=8 rats/group for one-way ANOVA. For two-way ANOVA p-values are shown for the main factors diet, MADB106 and the interaction of both main factors.