


Appendix 1 (as supplied by the authors)

Table of contents

Study timeframe definitions – Page 2

Incidence of supracondylar humerus fractures treated in Ontario at SickKids, other specialized paediatric centres, and community hospitals between 2002-2015 adapted from *Khoshbin et al.* – Page 3

Supplementary Figure 1: Study timeframe definitions



Supplementary Table 1: Incidence of supracondylar humerus fractures treated in Ontario between 2002-2015 using population based data adapted from *Khoshbin et al.* Presented by their location of treatment.

Year	Location of treatment, N (%)		
	SickKids	Other specialized paediatric centre	Community
April 2002 – March 2003	43 (11.14%)	138 (35.75%)	205 (53.17%)
April 2003 – March 2004	58 (14.61%)	143 (36.02%)	196 (49.38%)
April 2004 – March 2005	80 (18.48%)	148 (34.18%)	205 (47.3%)
April 2005 – March 2006	84 (19.81%)	148 (34.9%)	192 (45.31%)
April 2006 – March 2007	68 (15.32%)	168 (37.85%)	208 (46.9%)
April 2007 – March 2008	60 (15.23%)	146 (37.05%)	188 (47.69%)
April 2008 – March 2009	67 (17.45%)	144 (37.5%)	173 (45.01%)
April 2009 – March 2010	70 (19.02%)	152 (41.31%)	146 (39.69%)
April 2010 – March 2011	66 (20.12%)	132 (40.24%)	130 (39.55%)
April 2011 – March 2012	137 (35.13%)	115 (29.49%)	138 (35.37%)
April 2012 – March 2013	116 (29.82%)	128 (32.91%)	145 (37.3%)
April 2013 – March 2014	119 (29.97%)	157 (39.55%)	121 (30.46%)
April 2014 – March 2015	186 (37.73%)	148 (30.02%)	159 (32.29%)
April 2015 – March 2016	225 (39.54%)	201 (35.33%)	143 (25.14%)