

Diagnosis of cytomegalovirus pneumonia by quantitative polymerase chain reaction using bronchial washing fluid from patients with hematologic malignancies

Hwa Young Lee^{1,*}, Chin Kook Rhee^{1,*}, Joon Young Choi¹, Hea Yon Lee¹, Jong Wook Lee^{2,3} and Dong Gun Lee^{3,4,5}

¹ Department of Internal medicine, Division of Pulmonary and Critical Care Medicine, The Catholic University of Korea, Seoul, Korea

² Department of Internal Medicine, Division of Hematology, The Catholic University of Korea, Seoul, Korea

³ The Catholic Blood and Marrow Transplantation Center, The Catholic University of Korea, Seoul, Korea

⁴ Department of Internal medicine, Division of infectious diseases, The Catholic University of Korea, Seoul, Korea

⁵ Vaccine Bio Research Institute, The Catholic University of Korea, Seoul, Korea

* These authors have contributed equally to this work

Correspondence to: Dong Gun Lee, **email:** symonlee@catholic.ac.kr

Keywords: cytomegalovirus, pneumonia, real-time polymerase chain reaction, hematologic neoplasms

Received: August 04, 2016

Accepted: December 27, 2016

Published: January 04, 2017




Figure S1: Flow chart of 101 patients who harbored > 380 copies/mL qRT-PCR results in bronchial washing fluid.