

Oxidation of Secondary Methyl Ethers to Ketones

Pieter J. Gilissen, Daniel Blanco-Ania, Floris P. J. T. Rutjes*

Institute for Molecules and Materials, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, the Netherlands


Supporting Information

Substrate synthesis schemes	S3
Copies of ^1H NMR and ^{13}C NMR spectra	S5


Substrate synthesis schemes

Substrate **8** was synthesized from undecan-2-ol (**42**) in a common Williamson ether synthesis (Scheme S1). Substrates **16–20** were synthesized from 5-hydroxypentan-2-one (**10**) using regular synthetic transformations (Scheme S2). Substrate **21** was synthesized in two steps from γ -valerolactone (**45**; Scheme S3). Substrate **22** was synthesized from 1-phenylethanol (**46**; Scheme S4). Substrates **23–25** were synthesized from cyclohexane-1,4-diol (**47**; Scheme S5) and substrates **26–28** were synthesized from benzyl 4-oxopiperidine-1-carboxylate (**40**; Scheme S6).


Scheme S1: Synthesis of substrate **8**.




Scheme S2: Synthesis of substrates **16–20**.




Scheme S3: Synthesis of substrate **21**.




Scheme S4: Synthesis of substrate 22.





Scheme S5: Synthesis of substrates 23–25.







Scheme S6: Synthesis of substrates 26–28.




Copies of ^1H NMR and ^{13}C NMR spectra








S8





— 166.6480

132.8356
130.4341
129.5342
128.3243

— 76.2763


— 65.0738



— 56.0395



— 32.7918

— 24.7975

— 19.0093










S12



X = impurity found in the eluent







18

76.6128
72.0018
71.5076
69.9770
59.0802
55.9753
32.7426
25.5812
19.0393






S14











19











S20



X = heptane

7.3825
7.2463

4.3185
4.3023
4.2861
4.2700

3.2259

1.4459
1.4298



S21

X = heptane


— 143.5087

128.4248
127.4519
126.1874


— 79.6485

— 56.4454



— 23.8870




22



S22






23

170.6564
170.6165



77.2674
75.7872
71.7750
70.6345



55.9221
55.5845


28.3188
28.2738
27.1279
27.1122
21.4103
21.3857




S24












25




S27













26














— 207.6611

— 166.5388



132.9745
130.1827
129.5377
128.3824


— 64.1183

— 39.9746


— 30.0475



— 22.9227






30







S38




31




S39




31




S40








32






S42




S43










36






S46






37













38



— 2.7196














S54


X = methanesulfonyl chloride



S55



44



S56

X = impurity from the eluent

