

Supplemental table 3: Subgroup HADS Individual Item Scores for SST Participants in Whom CNV Developed in the Second Eye															
Before the last SST examination, Comparing Scores from (1) Baseline to prior to CNV diagnosis, and (2) Prior to and following CNV diagnosis															
Subfoveal CNV subgroup (n=42)															
Comparison of mean and median HADS scores															
Baseline to prior to CNV diagnosis								Prior to and following CNV diagnosis							
HADS Item	N	Baseline Mean	Baseline Median	Prior Mean	Prior Median	p-value*	Adjusted p-value**	N	Prior Mean	Prior Median	Following Mean	Following Median	p-value*	Adjusted p-value**	
hads_tense	34	3.26	3.00	3.41	3.50	0.26	1.00	32	3.34	3.00	3.50	4.00	0.23	1.00	
hads_enjoy	34	1.53	1.00	1.62	1.00	0.66	1.00	32	1.59	1.00	1.81	2.00	0.18	1.00	
hads_awful	34	3.59	4.00	3.59	4.00	1.00	1.00	32	3.56	4.00	3.56	4.00	1.00	1.00	
hads_laugh	34	1.32	1.00	1.44	1.00	0.46	1.00	32	1.38	1.00	1.28	1.00	0.56	1.00	
hads_worrying	33	3.55	4.00	3.33	3.00	0.18	1.00	28	3.21	3.00	3.61	4.00	<b>0.02</b>	0.23	
hads_cheerful	34	3.85	4.00	3.68	4.00	0.14	1.00	32	3.84	4.00	3.62	4.00	0.09	1.00	
hads_relaxed	34	1.76	2.00	1.68	2.00	0.45	1.00	32	1.66	2.00	1.69	2.00	0.86	1.00	
hads_slowed_down	34	2.79	3.00	2.85	3.00	0.71	1.00	32	2.84	3.00	2.78	3.00	0.72	1.00	
hads_butterflies	34	1.44	1.00	1.35	1.00	0.48	1.00	32	1.38	1.00	1.25	1.00	0.25	1.00	
hads_appearance	34	3.76	4.00	3.76	4.00	1.00	1.00	32	3.78	4.00	3.91	4.00	0.16	1.00	
hads_restless	34	3.21	3.00	3.35	3.00	0.26	1.00	32	3.41	3.50	3.28	3.00	0.40	1.00	
hads_look_forward	34	1.59	1.00	1.59	1.00	1.00	1.00	32	1.59	1.50	1.62	2.00	0.83	1.00	
hads_panic	34	3.62	4.00	3.65	4.00	0.79	1.00	32	3.59	4.00	3.62	4.00	0.81	1.00	
hads_book	34	1.12	1.00	1.29	1.00	0.18	1.00	32	1.25	1.00	1.59	1.00	0.07	0.98	
Probably subfoveal or juxtafoveal CNV subgroup (n=17)															
Comparison of mean and median HADS scores															
Baseline to prior to CNV diagnosis								Prior to and following CNV diagnosis							
HADS Item	N	Baseline Mean	Baseline Median	Prior Mean	Prior Median	p-value*	Adjusted p-value**	N	Prior Mean	Prior Median	Following Mean	Following Median	p-value*	Adjusted p-value**	
hads_tense	9	3.67	4.00	3.89	4.00	0.62	1.00	15	3.80	4.00	3.73	4.00	1.00	1.00	
hads_enjoy	9	1.11	1.00	1.56	1.00	0.50	1.00	15	1.47	1.00	1.67	1.00	0.63	1.00	
hads_awful	9	3.89	4.00	4.00	4.00	1.00	1.00	15	4.00	4.00	3.80	4.00	0.50	1.00	
hads_laugh	9	1.00	1.00	1.00	1.00	NA	NA	15	1.00	1.00	1.33	1.00	0.25	1.00	
hads_worrying	7	3.71	4.00	3.57	4.00	1.00	1.00	12	3.50	4.00	3.58	4.00	1.00	1.00	
hads_cheerful	9	4.00	4.00	4.00	4.00	NA	NA	15	3.87	4.00	3.80	4.00	1.00	1.00	
hads_relaxed	9	1.56	2.00	1.11	1.00	0.12	1.00	15	1.13	1.00	1.33	1.00	0.53	1.00	
hads_slowed_down	9	3.11	3.00	2.89	3.00	0.62	1.00	15	2.93	3.00	2.80	3.00	0.82	1.00	
hads_butterflies	9	1.22	1.00	1.00	1.00	0.50	1.00	15	1.00	1.00	1.07	1.00	1.00	1.00	
hads_appearance	9	3.67	4.00	4.00	4.00	0.50	1.00	15	4.00	4.00	3.80	4.00	0.25	1.00	
hads_restless	9	3.56	4.00	3.78	4.00	0.50	1.00	15	3.73	4.00	3.27	3.00	0.09	1.00	
hads_look_forward	9	1.00	1.00	1.11	1.00	1.00	1.00	15	1.13	1.00	1.33	1.00	0.38	1.00	
hads_panic	9	3.78	4.00	3.89	4.00	1.00	1.00	15	3.87	4.00	3.93	4.00	1.00	1.00	
hads_book	9	1.22	1.00	1.44	1.00	0.75	1.00	15	1.33	1.00	1.67	1.00	0.46	1.00	

Extrafoveal CNV subgroup (n=30)														
Comparison of mean and median HADS scores														
Baseline to prior to CNV diagnosis														
HADS Item	N	Baseline Mean	Baseline Median	Prior Mean	Prior Median	p-value*	Adjusted p-value**	Prior to and following CNV diagnosis						
								N	Prior Mean	Prior Median	Following Mean	Following Median	p-value*	Adjusted p-value**
<b>hads_tense</b>	20	3.65	4.00	3.55	4.00	0.69	1.00	28	3.36	3.50	3.50	4.00	0.59	1.00
<b>hads_enjoy</b>	20	1.5	1.00	1.65	1.00	0.69	1.00	28	1.64	1.00	1.86	2.00	0.15	1.00
<b>hads_awful</b>	20	3.45	4.00	3.80	4.00	0.09	1.00	28	3.64	4.00	3.75	4.00	0.50	1.00
<b>hads_laugh</b>	20	1.50	1.00	1.30	1.00	0.47	1.00	28	1.57	1.00	1.32	1.00	0.27	1.00
<b>hads_worrying</b>	19	3.37	3.00	3.42	3.00	1.00	1.00	24	3.33	3.00	3.54	4.00	0.46	1.00
<b>hads_cheerful</b>	20	3.75	4.00	3.65	4.00	0.69	1.00	28	3.75	4.00	3.75	4.00	1.00	1.00
<b>hads_relaxed</b>	20	1.60	1.50	1.65	2.00	1.00	1.00	28	1.64	2.00	1.79	2.00	0.53	1.00
<b>hads_slowed_down</b>	20	3.10	3.00	3.00	3.00	0.75	1.00	28	3.00	3.00	2.93	3.00	0.85	1.00
<b>hads_butterflies</b>	20	1.50	1.00	1.40	1.00	0.75	1.00	28	1.46	1.00	1.46	1.00	1.00	1.00
<b>hads_appearance</b>	20	4.00	4.00	3.95	4.00	1.00	1.00	28	3.86	4.00	3.96	4.00	0.50	1.00
<b>hads_restless</b>	20	3.45	3.50	3.25	3.00	0.56	1.00	28	3.21	3.00	3.32	3.00	0.69	1.00
<b>hads_look_forward</b>	20	1.40	1.00	1.55	1.00	0.45	1.00	28	1.61	1.00	1.36	1.00	0.25	1.00
<b>hads_panic</b>	20	3.60	4.00	3.70	4.00	0.75	1.00	28	3.64	4.00	3.50	4.00	0.43	1.00
<b>hads_book</b>	20	1.15	1.00	1.40	1.00	0.31	1.00	28	1.71	1.00	1.71	1.00	1.00	1.00
* Paired t-test (n ≥30) or Wilcoxon signed-rank test (n <30)														
** Bonferroni adjustment for multiple comparisons														
Bold font indicates statistical significance (p value < 0.05)														