

Combining molecular typing and spatial pattern analysis to identify areas of high tuberculosis transmission in a moderate-incidence county in Taiwan

Yih-Yuan Chen^a, Jia-Ru Chang^b, Chih-Da Wu^c, Yen-Po Yeh^f, Shiu-Ju Yang^b, Chih-Hao Hsu^b, Ming-Ching Lin^b, Ching-Fang Tsai^d, Ming-Shian Lin^e, Ih-Jen Su^b and Horng-Yunn Dou^{b#}

^aDepartment of Biochemical Science and Technology, National Chiayi University, Chia-Yi, Taiwan

^bNational Institute of Infectious Diseases and Vaccinology, National Health Research Institutes, Zhunan, Miaoli, Taiwan

^cDepartment of Forestry and Natural Resources, National Chiayi University, Chia-Yi, Taiwan; The Center for Health and the Global Environment, Harvard T. H. Chan School of Public Health, Boston, MA, USA

^dDepartment of Medical Research, Ditmanson Medical Foundation Chia-Yi Christian Hospital, Chia-Yi, Taiwan

^eDepartment of Internal Medicine, Ditmanson Medical Foundation Chia-Yi Christian Hospital, Chia-Yi, Taiwan

^fChang-Hua County Public Health Bureau, Taiwan

[#]Corresponding author

Horng-Yunn Dou, Ph.D.,

E-mail address: hydou@nhri.org.tw

Mailing address: National Institute of Infectious Diseases and Vaccinology, National Health Research Institutes, Taiwan

No.35, Keyan Road, Zhunan Town, Miaoli County 350, Taiwan, ROC.

Tel: 886-37-246166 ext 35529 Fax: 886-37-583009


Key words: *Mycobacterium tuberculosis*, geographical information system (GIS), Beijing lineage

Running title: Spatial analysis of high TB transmission

Figure legends

Supplementary Figure 1. Spatial distribution of TB cases in the study sites. (A) Total TB cases. (B) Random samples of molecularly analyzed TB cases. Esri ArcGIS 10.2 was used to create this figure (<http://www.esri.com/software/arcgis>).

(A)



(B)

