

SUPPLEMENTARY INFORMATION

Inhibition of *Staphylococcus aureus* cell wall biosynthesis by desleucyl-oritavancin: a quantitative peptidoglycan composition analysis by mass spectrometry

James D. Chang,¹ Erin E. Foster,¹ Aanchal N. Thadani,¹ Alejandro J. Ramirez,¹ and Sung Joon Kim^{1,2}

¹Department of Chemistry and Biochemistry, Baylor University, Waco, Texas 76798, United States.

²To whom the correspondence should be addressed: Tel: 1-254-710-2531; e-mail address: Sung_J_Kim@baylor.edu; Fax: 1-254-710-4272.

Running title: LC-MS analysis of oritavancin-treated *S. aureus*

Keywords: *S. aureus*, peptidoglycan, oritavancin, and LC-MS

Muropeptides	Cross-linking efficiency (ρ_{CL}) [†]		P value**
	des-Ori (0 µg/mL) [*]	des-Ori (10 µg/mL) [*]	
All muropeptides	0.636 ± 0.005	0.630 ± 0.002	0.024
With a terminal pentaglycine bridge structure	0.632 ± 0.006	0.6274 ± 0.002	0.092
Without a pentaglycine terminal bridge	0.686 ± 0.003	0.667 ± 0.010	0.004
Tripeptide stem structure	0.649 ± 0.016	0.625 ± 0.023	0.053
Tetrapeptide stem structure	0.593 ± 0.016	0.625 ± 0.002	0.003
Pentapeptide stem structure	0.639 ± 0.003	0.631 ± 0.004	0.008

[†] Cross-linking efficiencies calculated based on equation in Fig. 3b.

^{*} All error bars represent 95% confidence interval (n = 3).

^{**} Unpaired Student's t-test.

Table S1. Cross-linking efficiency of muramidase digested PG from isolated cell walls of *S. aureus* grown in absence or in presence of des-Ori (10 µg/ml) classified based on PG-stem modifications.

Muropeptide Acetylation states	Cross-linking efficiency (ρ_{CL}) [†]		P value ^{**}
	des-Ori (0 µg/mL) [*]	des-Ori (10 µg/mL) [*]	
-1	0.631 ± 0.015	0.607 ± 0.023	0.056
0	0.628 ± 0.008	0.622 ± 0.005	0.117
1 (or higher)	0.642 ± 0.003	0.638 ± 0.001	0.011

[†] Normalized percentile composition based on the sum of integrated XIC of selected ions.

^{*} All error bars represent 95% confidence interval (n = 3).

^{**} Unpaired Student's t-test.

Table S2. Cross-linking efficiency of muramidase digested PG from isolated cell walls of *S. aureus* grown in absence or in presence of des-Ori (10 µg/ml) classified based on acetylation state.

Muropeptides		Composition (%) [†]		P value ^{**}
PG-stem length	Acetylation state	des-Ori (0 µg/mL) [*]	des-Ori (10 µg/mL) [*]	
3	-1	9.764 ± 2.964	15.863 ± 2.964	0.010
	0	19.450 ± 10.808	10.733 ± 0.808	0.063
	1	70.787 ± 10.121	73.405 ± 2.443	0.469
4	-1	2.838 ± 0.356	5.227 ± 3.638	0.106
	0	35.386 ± 6.176	42.133 ± 3.375	0.038
	1	61.776 ± 5.895	52.641 ± 0.295	0.008
5	-1	6.063 ± 1.487	5.674 ± 2.598	0.700
	0	39.892 ± 2.287	40.235 ± 3.910	0.821
	1	54.045 ± 2.869	54.091 ± 1.984	0.968

[†] Normalized percentile composition based on the sum of integrated XIC of selected ions.

^{*} All error bars represent 95% confidence interval (n = 3).

^{**} Unpaired Student's t-test.

Table S3. Normalized PG composition of muramidase digested cell walls isolated from *S. aureus* grown in absence or in presence of des-Ori (10 µg/ml) classified based on PG-stem length and acetylation state.