

Table e-1. Schedule of procedures for diagnosis and research outcomes

		Baseline	6 mo	12 mo	18 mo	24 mo	30 mo	36 mo
Diagnostic Tests	History	X	X		X		X	
	Family History	X						
	Caregiver interview	X	X		X		X	
	Neurological Exam	X	X		X		X	
	EMG	X	X*		X*		X*	
	MiniMental State Exam ¹	X	X		X		X	
	Frontal Assessment Battery ²	X	X		X		X	
	Mattis Dementia Rating Scale -2 ^{3, 4}	X	X		X		X	
	Research Outcome Measures							
Motor	ALSFRS-R⁵	X	X	X	X	X	X	X
	Timed gait, timed reading, finger and foot tapping, 9-hole peg test	X	X		X		X	
	Physiology: TMS, EIM	X	X		X		X	
Cognitive	Letter Fluency Score	X	X	X	X	X	X	X
	DRS, FAB, MMSE, D-KEFS Trails, Sorting, Fluency, 20 Questions ⁶	X	X		X		X	
	Language: ADRC FTLD module ⁷	X	X		X		X	
Behavioral	Frontal Behavioral Inventory⁸	X	X	X	X	X	X	X
Mood, Psuedo-bulbar affect	ALS depression Inventory, ⁹ CNS lability ¹⁰	X	X		X		X	
Quality of Life	SF-8 Health Survey ¹¹	X	X		X		X	
Imaging	MRI							
Biospecimens	CSF, blood	X	X		X		X	
	Skin biopsy	X	X**					

*EMG repeated when El Escorial certainty less than definite

** Skin Biopsy performed only once, usually first, occasionally second visit

References

1. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. J Psychiatr Res 1975;12:189-198.

2. Dubois B, Slachevsky A, Litvan I, Pillon B. The FAB: a Frontal Assessment Battery at bedside. *Neurology* 2000;55:1621-1626.
3. Jurica SJ, Leitten CL, Mattis S. *Dementia Rating Scale: Professional manual*. Odessa: Psychological Assessment Resources, 2001.
4. Rascovsky K, Salmon DP, Hansen LA, Galasko D. Distinct cognitive profiles and rates of decline on the Mattis Dementia Rating Scale in autopsy-confirmed frontotemporal dementia and Alzheimer's disease. *J Int Neuropsychol Soc* 2008;14:373-383.
5. Cedarbaum JM, Stambler N, Malta E, et al. The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments of respiratory function. BDNF ALS Study Group (Phase III). *J Neurol Sci* 1999;169:13-21.
6. Delis DC, Kramer JH, Kaplan E, Holdnack J. Reliability and validity of the Delis-Kaplan Executive Function System: an update. *J Int Neuropsychol Soc* 2004;10:301-303.
7. ADRC FTLD module. Available at: https://www.alz.washington.edu/WEB/forms_ftld.html.
8. Milan G, Lamenza F, Iavarone A, et al. Frontal Behavioural Inventory in the differential diagnosis of dementia. *Acta Neurol Scand* 2008;117:260-265.
9. Hammer EM, Hacker S, Hautzinger M, Meyer TD, Kubler A. Validity of the ALS-Depression-Inventory (ADI-12)--a new screening instrument for depressive disorders in patients with amyotrophic lateral sclerosis. *Journal of affective disorders* 2008;109:213-219.
10. Moore SR, Gresham LS, Bromberg MB, Kasarkis EJ, Smith RA. A self report measure of affective lability. *J Neurol Neurosurg Psychiatry* 1997;63:89-93.
11. SF-8 Health Survey. Available at <https://campaign.optum.com/optum-outcomes/what-we-do/health-surveys.html>