

Supplementary Table 2. Demographic and pathological characteristics according to availability of information on medication use.

Characteristic	Statin status known (n=680) No. (%)	Statin status unknown (n=60) No. (%)	P value
<i>Age category</i>			
<50	47 (6.9)	0 (0)	<0.001
50-<60	71 (10.4)	4 (6.7)	
60-<70	197 (29.0)	8 (13.3)	
70-<80	239 (35.2)	23 (38.3)	
>80	126 (18.5)	25 (41.7)	
<i>Gender</i>			
Male	370 (54.4)	28 (46.7)	0.25
Female	310 (45.6)	32 (53.3)	
<i>Year of diagnosis</i>			
2004	101 (14.9)	20 (33.3)	<0.001
2005	134 (19.7)	21 (35.0)	
2006	131 (19.3)	5 (8.3)	
2007	146 (21.5)	8 (13.3)	
2008	168 (24.7)	6 (10.0)	
<i>Stage</i>			
II	389 (57.2)	37 (61.7)	0.50
III	291 (42.8)	23 (38.3)	
<i>Grade</i>			
Well-moderate	577 (84.9)	54 (90.0)	0.51
Poor	99 (14.6)	6 (10.0)	
Unknown	4 (0.59)	0 (0.0)	
<i>MSI status</i>			
Non MSI-high	437 (64.3)	34 (56.7)	0.50
MSI-high	123 (18.1)	13 (21.7)	
Unknown	120 (17.7)	13 (21.7)	

Abbreviations: MSI – microsatellite instability.