Supplementary information for

A GalNAc/Gal-specific lectin from the sea mussel Crenomytilus grayanus

modulates immune response in macrophages and in mice

Oleg V. Chernikov^{1,#}, Wei-Ting Wong^{2,3,#}, Lan-Hui Li⁴, Irina V. Chikalovets^{1,5},

Valentina I. Molchanova¹, Shih-Hsiung Wu⁶, Jiahn-Haur Liao⁶, Kuo-Feng Hua^{2,7,*}

¹G.B. Elyakov Pacific Institute of Bioorganic Chemistry FEB RAS, Vladivostok,

Russia; ²Department of Biotechnology and Animal Science, National Ilan University,

Ilan, Taiwan; ³Graduate Institute of Life Sciences, National Defense Medical Center,

Taipei, Taiwan; ⁴Department of Laboratory, Kunming Branch, Taipei City Hospital,

Taipei, Taiwan; ⁵School of Natural Sciences, Far Eastern Federal University,

Vladivostok, Russia; ⁶Institute of Biological Chemistry, Academia Sinica, Taipei,

Taiwan; ⁷Department of Pathology, Tri-Service General Hospital, National Defense

Medical Center, Taipei, Taiwan

*Correspondence to:

Dr. Kuo-Feng Hua

Department of Biotechnology and Animal Science,

National Ilan University, Ilan, Taiwan

No. 1, Sec. 1, Shennong Rd., Ilan 260, Taiwan

Tel: +(886)-3-935-7626

Fax: +(886)-3-931-1526

E-mail: kuofenghua@gmail.com

^{*}These authors contributed equally to this work.




Fig. S1. The full-length gels of Figs.1A and 1B.




Fig. S2. The full-length gels of Figs. 3C and 3E.

Fig. S3
A (full gels of Fig. 4A)
B (full gels of Fig. 4B)




Fig. S3. The full-length gels of Figs. 4A and 4B.

Fig. S4 (full gels of Fig. 5B)




Fig. S4. The full-length gels of Fig. 5B.

Fig. S5 (full gels of Fig. 6A)





Fig. S5. The full-length gels of Fig. 6A.

Fig. S6



C (full gels of Fig. 7E)




Fig. S6. The full-length gels of Figs. 7C, 7D and 7E.