

Supplementary information

Toll-like Receptor 4 Deficiency Reduces Oxidative Stress and Macrophage Mediated Inflammation in Hypertensive Kidney

Sathnur Pushpakumar¹, MD, PhD, Lu Ren¹, PhD, Sourav Kundu², PhD, Alejandra Gamon³,
Suresh C. Tyagi¹, PhD, Utpal Sen¹, PhD*

¹Department of Physiology, University of Louisville School of Medicine, Louisville, KY-40202

²Institute of Advanced Study in Science and Technology, Guwahati, Assam – 781035, India

³University of California, San Diego, CA 92093

Running title: TLR4 induced oxidative stress and renal inflammation

Keywords: TLR4, hypertension, oxidative stress, macrophage, bone marrow-derived fibroblast

Word count: 5487


Number of Figures: 12

No. of tables: 1

Correspondence:


Utpal Sen, Ph.D.
500 S Preston St.
HSC-A, Room 802,
University of Louisville,
KY-40202,
Ph: 502-852-2030;
Fax: 502-852-6239;
Email: utpal.sen@louisville.edu

Figure 5C




Supplementary figure S1: Uncropped images for Nox4 and p22-phox.

Figure 8A




Supplementary figure S2: Uncropped images for MCP-1 and MIP-2.

Figure 8A




Supplementary figure S3: Uncropped images for TNF-α and IL-6.

Figure 9A



Supplementary figure S4: Uncropped images for TGF-β and p-Smad2/3.

Figure 9C



Supplementary figure S5: Uncropped images for fibronectin, collagen 1 and collagen IV.