

Supplementary Table S8**Multivariate analysis of clinicopathological characteristics, BCSS endpoint**

Characteristic	Uppsala (n=233)			Stockholm (n=113)		
	HR	95% CI	P value	HR	95% CI	P value
ER*	0.54	0.25 – 1.16	0.11	0.30	0.08 – 1.15	0.08
PR*	1.58	0.73 – 3.42	0.24	2.25	0.88 – 5.77	0.09
Age†	0.92	0.54 – 1.57	0.75	0.71	0.30 – 1.64	0.42
Tumour size‡	2.39	1.31 – 4.37	0.005	1.52	0.64 – 3.61	0.34
Elston grade§	1.41	0.65 – 3.05	0.39	2.77	0.60 – 12.85	0.20
Nodal status¶	2.31	1.39 – 3.84	0.001	1.19	0.52 – 2.75	0.68
Ki67¥	1.71	0.98 – 2.98	0.06	2.5	0.87 – 7.20	0.09

HR= Hazard ratio, CI= Confidence interval

*ER; PR positive vs. negative

†Age <50 vs. ≥50

‡Size <20 vs. ≥20

§Grade I vs. II and III

¥Nodal status, negative vs. positive

¥Ki67 <10 vs. ≥10 Uppsala, <8 vs. ≥8 Stockholm