
Permian
Triassic
Jurassic
Cretaceous
Paleogene
Neogene
164 species dataset this study
309 species dataset this study
Biju and Bossuyt (2003)
San Mauro et al. (2005)
Bossuyt et al. (2006)
van Bocxlaer et al. (2006)
van der Meijden et al. (2007)
Roelants et al. (2007)
Heinicke et al. (2009)
Wiens (2011)
Kurabayashi et al. (2011)
Irisarri et al. (2012)
de S á (2012)
Zhang et al. (2013), Multidivtime
Zhang et al. (2013), BEAST
Fraz o et al. (2015)
Cannatella (2015)
Anura
Neobatrachia
Hyloidea
Microhylidae
Natatanura
300
250
200
150
100
50
0
Age (Mya)

References

Biju SD, Bossuyt F (2003) New frog family from India reveals an ancient biogeographical link with the Seychelles. *Nature*, 425, 711–714.

Bossuyt F, Brown R, Hillis D, Cannatella D, Milinkovitch M (2006) Phylogeny and biogeography of a cosmopolitan frog radiation: Late Cretaceous diversification resulted in continent-scale endemism in the family Ranidae. *Systematic Biology*, 55, 579–594.

Cannatella D (2015) Xenopus in space and time: fossils, node calibrations, tip-dating, and paleobiogeography. *Cytogenetic and Genome Research*, 145, 283–301.

de Sá RO, Streicher JW, Sekonyela R *et al.* (2012) Molecular phylogeny of microhylid frogs (Anura: Microhylidae) with emphasis on relationships among New World genera. *BMC Evolutionary Biology*, 12, 241.

Frazão A, Silva HR da, Russo CA de M (2015) The gondwana breakup and the history of the Atlantic and Indian oceans unveils two new clades for early neobatrachian diversification (A Rzhetsky, Ed,). *PLOS ONE*, 10, e0143926.

Heinicke MP, Duellman WE, Trueb L *et al.* (2009) A new frog family (Anura: Terrarana) from South America and an expanded direct-developing clade revealed by molecular phylogeny. *Zootaxa*, 35, 1–35.

Irisarri I, Mauro DS, Abascal F *et al.* (2012) The origin of modern frogs (Neobatrachia) was accompanied by acceleration in mitochondrial and nuclear substitution rates. *BMC Genomics*, 13, 626.

Kurabayashi A, Matsui M, Belabut DM *et al.* (2011) From Antarctica or Asia? New colonization scenario for Australian-New Guinean narrow mouth toads suggested from the findings on a mysterious genus Gastrophrynoides. *BMC Evolutionary Biology*, 11, 175.

van Bocxlaer I, Roelants K, Biju SD, Nagaraju J, Bossuyt F (2006) Late Cretaceous vicariance in Gondwanan amphibians. *PLOS ONE*, 1, e74.

van der Meijden A, Vences M, Hoegg S *et al.* (2007) Nuclear gene phylogeny of narrow-mouthed toads (Family: Microhylidae) and a discussion of competing hypotheses concerning their biogeographical origins. *Molecular Phylogenetics and Evolution*, 44, 1017–1030.

Roelants K, Gower DJ, Wilkinson M *et al.* (2007) Global patterns of diversification in the history of modern amphibians. *Proceedings of the National Academy of Sciences*, 104, 887–892.

San Mauro D, Vences M, Alcobendas M, Zardoya R, Meyer A (2005) Initial diversification of living amphibians predated the breakup of Pangaea. *The American Naturalist*, 165, 590–599.

Wiens JJ (2011) Re-evolution of lost mandibular teeth in frogs after more than 200 million years, and re-evaluating dollo's law. *Evolution*, 65, 1283–1296.

Zhang P, Liang D, Mao R-L *et al.* (2013) Efficient sequencing of anuran mtdnas and a mitogenomic exploration of the phylogeny and evolution of frogs. *Molecular Biology and Evolution*, 30, 1899–1915.