


Supplementary information for SREP-17-11580A

***In vitro* significance of SOCS-3 and SOCS-4 and potential mechanistic links to wound healing**


Yi Feng^{1#}, Andrew J. Sanders^{1#}, Liam D. Morgan¹, Sioned Owen¹, Fiona Ruge^{1,2}, Keith G. Harding²,

***Wen G. Jiang¹**


¹Cardiff China Medical Research Collaborative and ²Wound Healing Research Unit, Cardiff University School of Medicine, Cardiff University, Cardiff, CF14 4XN, UK, jiangw@cf.ac.uk




Full gel profile of composite used in figure 1a




1 = HaCaT pEF6
 2 = HaCaT SOCS-4 ribozyme 1-2
 3 = HaCaT SOCS-4 ribozyme 1-4
 4 = HaCaT SOCS-4 ribozyme 2-1 (used in study)
 5 = HaCaT SOCS-4 ribozyme 2-2



1 = HECV pEF6
 2 = HECV ribozyme 2-1 (used in study)



1 = HaCaT pEF6
 2 = HaCaT SOCS-3 exp1
 3 = HaCaT SOCS-3 exp2 (used in study)



1 = HECV pEF6
 2 = HECV SOCS-3 exp1
 3 = HECV SOCS-3 exp2 (used in study)

Full length blots used in composites in figure 1d,e,h,i