

Supplementary Table 1. Demographics and baseline characteristics of the warfarin-initiated cohort

Variable	Warfarin-initiated cohort (n=1,814)
Age (years), mean (SD)	71.3 (9.8)
<65	388 (21.4)
65–74	715 (39.4)
≥75	711 (39.2)
Female	821 (45.3)
Height (cm), mean (SD)	162.1 (8.94)
Weight (kg), mean (SD)	62.2 (11.4)
NIHSS score*	
Mean (SD)	7.1 (6.6)
Median (IQR)	5.0 (2.0–11.0)
<8	1,130 (62.7)
8–16	457 (25.4)
>16	215 (11.9)
Creatinine (mg/dL), mean (SD)	1.03 (0.67)
CHADS ₂ score, mean (SD)	3.46 (0.96)
CHA ₂ DS ₂ -VASc score, mean (SD)	4.88 (1.42)
Congestive heart failure	216 (11.9)
Hypertension	1,242 (68.5)
Diabetes mellitus	477 (26.3)
Stroke/TIA	1,814 (100.0)
Vascular disease	324 (17.9)
Total number of INR measurements included in TTR analysis	42,634
INR measurements included in TTR analysis per patient, mean (SD)	23.6 (20.7)
Total duration included in TTR analysis (month)	29,512
Duration included in TTR analysis per patient (month), mean (SD)	16.4 (14.3)
Interval between consecutive INR measurements included in TTR analysis (day), mean (SD)	16.5 (13.2)

Values are n (%) otherwise indicated.

*NIHSS score, not available in 12 patients.

INR: international normalized ratio, IQR: interquartile range, NIHSS: National Institutes of Health Stroke Scale, TTR: time in therapeutic range.