

Supplementary Table 1. Zero-order correlation coefficients between overall (global) pain intensity, pain extent, and pain intensity at each pain site and measures of pain interference and psychological functioning

Pain intensity and extent measures	Entire sample (N = 182)		Only those with pain at a specific site		
	Pain Interference (BPI)	Psychological Functioning (MHI-5)	N	Pain Interference (BPI)	Psychological Functioning (MHI-5)
Global pain intensity	.63**	-.30**			
Pain extent	.39**	-.19*			
Intensity at each pain site					
Head	.19**	-.20**	28	.31	-.19
Neck	.26**	-.18*	93	.47**	-.20
Shoulder	.35**	-.25**	110	.48**	-.36**
Upper back	.24**	-.14	67	.39**	-.34**
Lower back	.39**	-.14	128	.49**	-.16
Arms	.37**	-.22**	69	.47**	-.35**
Elbows	.20**	-.01	29	.39*	-.04
Wrists	.27**	-.14	46	.61**	-.30*
Hands	.40**	-.18*	70	.61**	-.39**
Buttocks	.28**	-.15	34	.62**	-.27
Hips	.42**	-.15*	85	.57**	-.38**
Chest	.13	-.06	19	.12	-.36
Abdomen/Pelvis	.19*	-.09	35	.41*	-.18
Legs	.53**	-.26**	125	.54**	-.28**
Knees	.26**	-.21**	86	.48**	-.34**
Ankles	.36**	-.20**	59	.48**	-.37**
Feet	.45**	-.20**	67	.48**	-.25*
Other	.26**	-.18*	19	.37	-.21

*p < .05, **p < .01