

Supplementary Table1. Characteristics of Study Population between the follow up and lost groups

Characteristics	Baseline population		P value
	Follow up group (n=1499)	lost to follow up group (n=181)	
Demographic			
Age (years)	61.4±11.4	57.6±10.5	<0.01
Male Sex n (%)	629 (42.0)	83(45.9)	0.316
BMI (kg/m ²)	25.6±3.3	25.8±3.4	0.273
Medical history			
Hypertension n (%)	690(46.0)	72(39.8)	0.111
Diabetes mellitus n (%)	292(19.5)	30(16.6)	0.348
Current smoking n (%)	247(16.5)	35(19.3)	0.331
Systolic BP (mm Hg)	132.7±17.0	130.6±16.7	0.136
Diastolic BP (mm Hg)	76.6±10.3	77.8±10.7	0.077
Laboratory values			
FBG (mmol/L)	5.4±1.6	5.3±1.5	0.474
PBG (mmol/L)	8.0±4.2	7.5±3.7	<0.05
Total cholesterol (mmol/L)	5.1±0.9	5.0±0.9	0.321
Triglycerides (mmol/L)	1.8±1.2	1.9±1.1	0.054
LDL cholesterol (mmol/L)	3.0±0.7	2.9±0.6	0.21
HDL cholesterol (mmol/L)	1.4±0.3	1.4±0.3	0.386
Uric acid (μmol/L)	291.4±73.3	276.9±70.8	<0.01
eGFR (mL/min/1.73m ²)	88.9±15.5	92.1±16.0	0.009
NT-proBNP (pg/mL)	41.7(19.8,81.9)	37.4(17.3,73.9)	<0.01
Hs-CRP (mg/L)	2.3 (1.4, 3.5)	2.2 (1.3, 3.4)	0.189
Medication use			
Antihypertensives n (%)	528 (35.2)	55 (30.4)	0.197
Antidiabetic n (%)	189 (12.6)	19 (10.5)	0.415

Lipid lowering	n (%)	262 (17.5)	29(16.0)	0.625
Aspirin	n (%)	360 (24.0)	37 (20.4)	0.285
