

A critical courier role of volatile oils from *Dalbergia odorifera* for cardiac protection *in vivo* by QiShenYiQi

Jiahui Yu^{1, 2#}, Wen Zhang^{1, 3#}, Yiqian Zhang^{3, 4}, Yadong Wang¹, Boli Zhang¹, Guanwei Fan^{1, 5*}, Yan Zhu^{1, 2*}

¹Tianjin State Key Laboratory of Modern Chinese Medicine, Tianjin University of Traditional Chinese Medicine, Tianjin, China

²Research and Development Center of TCM, Tianjin International Joint Academy of Biotechnology & Medicine, Tianjin, China


³State Key Laboratory of Core Technology in Innovative Chinese Medicine, Tianjin Tasly Holding Group Co., Ltd., Tianjin, China

⁴State Key Laboratory of Natural Medicines, China Pharmaceutical University, Nanjing, China

⁵The First Teaching Hospital of Tianjin University of Traditional Chinese Medicine Tianjin, China,

Equal contribution to this work

* Correspondence should be addressed to Yan Zhu, yanzhu.harvard@iCloud.com (YZ) and Guanwei Fan, fgw1005@hotmail.com (GWF)



Supplementary Fig S1. Determination of *in vitro* QSYQ dose by cell viability assay. (A) Toxicity concentration of QSYQ to untreated cells. (B) IC₅₀ curve of QSYQ. (C) QSYQ increased cell viability in concentration-dependent manner. Data are expressed as the mean \pm SD, n=3. ^{##}p<0.01 vs. Control group. ^{**}p<0.01 vs. H₂O₂ group.