

SUPPLEMENTAL MATERIAL

THE ACTIVITIES OF THE C-TERMINAL REGIONS OF THE FORMIN PROTEIN DISHEVELED-ASSOCIATED ACTIVATOR OF MORPHOGENESIS (DAAM) IN ACTIN DYNAMICS

Andrea Teréz Vig¹, István Földi², Szilárd Szikora², Ede Migh², Rita Gombos², Mónika Ágnes Tóth¹, Tamás Huber¹, Réka Pintér¹, Gábor Csaba Talián¹, József Mihály², Beáta Bugyi^{1,3,*}

* to whom the correspondence should be addressed: Beáta Bugyi, University of Pécs, Medical School, Department of Biophysics, Pécs, Szigeti str. 12, H-7624, Hungary, Phone: +36-72-536-265, E-mail: beata.bugyi@aok.pte.hu

Institutions:

¹ University of Pécs, Medical School, Department of Biophysics, Szigeti str. 12, Pécs, H-7624, Hungary

² Biological Research Centre, Hungarian Academy of Sciences, Institute of Genetics, MTA-SZBK NAP B Axon Growth and Regeneration Group, Temesvári krt. 62, Szeged, H-6726, Hungary

³ Szentágothai Research Center, Ifjúság str. 34, Pécs, H-7624, Hungary

Supplemental movie 1. Filopodia dynamics in *Drosophila* primary neurons.

Time lapse sequence of filopodia dynamics of a neuronal growth cone expressing Actin5C::GFP. Scale bar = 5 μ m, elapsed time is given in seconds.