

Longitudinal observations of progressive cardiac dysfunction in a cardiomyopathic animal model by self-gated cine imaging based on 11.7-T magnetic resonance imaging

Shigeyoshi Saito PhD^{1,2*}, Minori Tanoue BSc¹, Kasumi Masuda PhD³, Yuki Mori PhD^{2,4}, Satoshi Nakatani MD, PhD³, Yoshichika Yoshioka PhD^{2,4}, Kenya Murase PhD^{1,2}

¹ Department of Medical Physics and Engineering, Division of Health Sciences, Osaka University Graduate School of Medicine, Suita, Osaka, Japan, 560-0871

² Center for Information and Neural Networks (CiNet), National Institute of Information and Communications Technology, and Osaka University, Suita, Osaka, Japan, 560-0871

³ Department of Functional Diagnostics, Division of Health Sciences, Osaka University Graduate School of Medicine, Suita, Osaka, Japan 560-0871

⁴ Biofunctional Imaging Laboratory, Immunology Frontier Research Center (IFReC), Osaka University, Suita, Osaka, Japan, 560-0871

Supplemental files (6 files)

Supplementary 1

J2N_k_20weeks.avi

Typical movie of a 4-chamber view obtained using magnetic resonance imaging for a
J2N_k hamster

Supplementary 2

J2N_n_20weeks.avi

Typical movie of a 4-chamber view obtained using magnetic resonance imaging for a
J2N_n hamster

Supplementary 3

K_Mpeg.wmv

Typical movie of an SA view obtained using b-mode echocardiography for a J2N_k
hamster

Supplementary 4

N_Mpeg.wmv

Typical movie of an SA view obtained using b-mode echocardiography for a J2N_n
hamster

Supplementary 5

k_SA.avi

Typical movie of an SA view obtained using magnetic resonance imaging for a J2N_k
hamster

Supplementary 6

n_SA.avi

Typical movie of an SA view obtained using magnetic resonance imaging for a J2N_n
hamster