

Sterol targeting drugs reveal life cycle stage-specific differences in trypanosome lipid rafts

Aabha I. Sharma^{1,2}, Cheryl L. Olson^{1,2}, João I. Mamede³, Felipe Gazos-Lopes⁵,
Conrad L. Epting^{1,4}, Igor C. Almeida⁵, David M. Engman^{1,2,6}

Departments of ¹Pathology, ²Microbiology-Immunology, ³Cell and Molecular Biology and ⁴Pediatrics, Northwestern University, Chicago, Illinois, USA. ⁵Department of Biological Sciences, University of Texas at El Paso, El Paso, TX, USA. ⁶Department of Pathology and Laboratory Medicine, Cedars-Sinai Medical Center, Los Angeles, CA, USA. Correspondence should be addressed to I.C.A. or D.M.E. (email: icalmeida@utep.edu or david.engman@csmc.edu).

Supplementary Figure 1. GC-MS fragmentation spectra of sterol species from whole cell and flagellar extracts of procyclic *T. brucei* cells. A-F, Cholesterol, zymosterol, cholesta-5-7-24-trienol, ergosta-5,7,25(27)-trienol, lanosterol, and stigmastanol (internal standard). $[M]^+$, molecular ion; OTMS, O-trimethylsilyl fragment; CH_3 , methyl group.

