Supplementary Table 1 First MRI assessment of tumour burden in Tsc1 +/- mice before treatment

(Mann Whitney test)

A. Lesion number (all cystic, papillary and solid lesions)

	Number of			P (compared	P (compared with	P (compared with
Treatment*	mice	Median	Range	with Vehicle)	Rapamycin)	Rapamycin+Atorvastatin)
Vehicle	6	14	8-24			
Atorvastatin	6	15.5	10-29	0.5628	0.8398	0.9134
Rapamycin	6	18	10-27	0.5130		
Rapamycin+Atorvastatin	6	17	14-22	0.4935	0.8463	

B. Lesion volume (all cystic, papillary and solid lesions)

	Number of	Median		P (compared	P (compared with	P (compared with
Treatment*	mice	(mm^3)	Range (mm ³)	with Vehicle)	Rapamycin)	Rapamycin+Atorvastatin)
Vehicle	6	20.205	10.18-93.95			
Atorvastatin	6	15.435	5.04-38.16	0.5887	0.8182	0.8182
Rapamycin	6	16.52	10.59-47.65	0.6991		
Rapamycin+Atorvastatin	6	21.11	7.61-29.78	0.9372	0.6991	

^{*} The 1st MRI scan was performed before treatment at the age of 12 months.