

Grape-seed Polyphenols Play a Protective Role in Elastase-induced Abdominal Aortic Aneurysm in Mice

Chao Wang^{1,2,3,#}, Yunxia Wang^{1,2,4,#}, Maomao Yu^{1,2,#}, Cong Chen^{1,2,4}, Lu Xu^{1,2},
Yini Cao^{1,2,4}, Rong Qi^{1,2,4,*}

¹ Peking University Institute of Cardiovascular Sciences, Peking University Health Science Center, Peking University, Beijing, 100191, China

² Key Laboratory of Molecular Cardiovascular Sciences, Ministry of Education, Beijing, 100191, China

³ Second School of Clinical Medicine, Peking University, Beijing, 100044, China

⁴ Beijing Key Laboratory of Molecular Pharmaceutics and New Drug Delivery Systems, Beijing 100191, China

These three authors contributed equally to this work.

* Corresponding author: ronaqi@bjmu.edu.cn

Supplemental Data




Fig. S1 Effects of GSP on normal aorta. Effects of GSP on (a) morphology, (b) maximal abdominal aortic diameter, (c) H&E staining for aorta structure, $n=3$. PBS, GSP-1 and GSP-2 were represented the mice groups administrated by 0.01M PBS, 800 mg/kg GSP and 1600 mg/kg GSP, respectively.