

1 **Social support helps protect against perinatal bonding failure and depression among**
2 **mothers: a prospective cohort study – Supplementary Information**

3

4 Masako Ohara¹⁾, Takashi Okada^{1)*}, Branko Aleksic¹⁾, Mako Morikawa¹⁾, Chika Kubota¹⁾,
5 Yukako Nakamura¹⁾, Tomoko Shiino¹⁾, Aya Yamauchi¹⁾, Yota Uno¹⁾, Satomi Murase¹⁾, Setsuko
6 Goto²⁾, Atsuko Kanai³⁾, Tomoko Masuda⁴⁾, Masahiro Nakatochi⁵⁾, Masahiko Ando⁵⁾, and Norio
7 Ozaki¹⁾

8

9 1) Department of Psychiatry, Nagoya University Graduate School of Medicine, Nagoya, Japan

10 2) Sugiyama Jogakuen University, Nagoya, Japan

11 3) Graduate School of Education and Human Development, Nagoya University, Nagoya, Japan

12 4) Graduate School of Law, Nagoya University, Nagoya, Japan

13 5) Center for Advanced Medicine and Clinical Research, Nagoya University Hospital, Nagoya,
14 Japan

15

16 *Corresponding author

17 Takashi Okada, M.D., Ph.D.

18 Department of Psychiatry, Nagoya University Graduate School of Medicine

19 65 Tsurumai-cho, Showa-ku, Nagoya, Aichi 466-8550, Japan

20 E-mail: okada@med.nagoya-u.ac.jp

21 Tel: +81-52-744-2282, Fax: +81-52-744-2293

- 1 **Table of contents**
- 2 Supplementary Table S1. Demographic characteristics of included and all mothers

1 **Supplementary Table S1. Demographic characteristics of included and all mothers**

	Included mothers (n= 477-494)		all mothers (n= 973-1011)		t	p
	Mean (SD)		Mean (SD)			
Age (years)	32.4	(4.5)	32.1	(4.5)	1.13	0.65
Number of children	0.24	(0.53)	0.25	(0.53)	0.42	0.54
Partner's age (years)	34.7	(5.7)	34.5	(5.8)	0.43	0.68

2

3