S2 Table. Change within and between groups of children in sedentary behavior and physical activity over time as a proportion of measurement time (estimates, 95% confidence intervals and p-value).

Including children (n=111) who had acceptable accelerometer use in all three weeks.

| CHILDREN                                 | Unadjusted                |                 | Adjusted*                  |                 |
|--|---------------------------|-----------------|----------------------------|-----------------|
|  | estimate (95% CI)         | <i>p</i> -value | estimate (95% CI)          | <i>p</i> -value |
| Sedentary behavior**                     |                           |                 |                            |                 |
| difference at baseline (ref = control)   | -0.191 (-2.612 to 2.230)  | 0.876           | -0.099 (-2.513 to 2.315)   | 0.935           |
| change in time, control                  | 0.013 (-0.013 to 0.040)   | 0.329           | 0.016 (-0.011 to 0.043)    | 0.251           |
| change in time, intervention             | 0.016 (-0.013 to 0.044)   | 0.283           | 0.015 (-0.013 to 0.043)    | 0.280           |
| intervention effect (ref = control)      | 0.002 (-0.037 to 0.041)   | 0.902           | -0.0005 (-0.040 to 0.039)  | 0.980           |
| Standing still**                         |                           |                 |                            |                 |
| difference at baseline (ref = control)   | 0.108 (-0.723 to 0.940)   | 0.797           | -0.103 (-0.951 to 0.745)   | 0.810           |
| change in time, control                  | -0.001 (-0.011 to 0.010)  | 0.907           | -0.003 (-0.013 to 0.008)   | 0.606           |
| change in time, intervention             | -0.005 (-0.016 to 0.006)  | 0.400           | -0.004 (-0.015 to 0.006)   | 0.424           |
| intervention effect (ref = control)      | -0.004 (-0.019 to 0.011)  | 0.591           | -0.002 (-0.017 to 0.013)   | 0.828           |
| Light physical activity**                |                           |                 |                            |                 |
| difference at baseline (ref = control)   | -0.117 (-1.332 to 1.097)  | 0.848           | -0.050 (-1.277 to 1.177)   | 0.936           |
| change in time, control                  | -0.013 (-0.025 to -0.001) | 0.037           | -0.013 (-0.255 to -0.0002) | 0.046           |
| change in time, intervention             | -0.014 (-0.027 to -0.001) | 0.041           | -0.014 (-0.027 to -0.001)  | 0.038           |
| intervention effect (ref = control)      | -0.001 (-0.019 to 0.017)  | 0.934           | -0.001 (-0.019 to 0.017)   | 0.916           |
| Moderate-to-vigorous physical activity** |                           |                 |                            |                 |
| difference at baseline (ref = control)   | 0.197 (-1.225 to 1.620)   | 0.784           | 0.247 (-1.156 to 1.650)    | 0.728           |
| change in time, control                  | -0.001 (-0.016 to 0.015)  | 0.948           | -0.001 (-0.017 to 0.015)   | 0.887           |
| change in time, intervention             | 0.002 (-0.014 to 0.019)   | 0.784           | 0.002 (-0.014 to 0.019)    | 0.765           |
| intervention effect (ref = control)      | 0.003 (-0.020 to 0.025)   | 0.807           | 0.004 (-0.019 to 0.026)    | 0.754           |

<sup>\*</sup> Adjusted for child's BMI, daycare or preschool, and number of siblings

<sup>\*\*</sup> Proportion of measurement time