

Pleiotropic Effects of Apolipoprotein C3 on HDL functionality and Adipose Tissue

Metabolic Activity

Evangelia Zvintzou¹, Marie Lhomme², Stella Chasapi³, Serafoula Filou¹, Vassilis Theodoropoulos¹, Eva Xapapadaki¹, Anatol Kontush⁴, George Spyroulias³, Constantinos C. Tellis⁵, Alexandros D. Tselepis⁵, Caterina Constantinou¹ and Kyriakos E. Kypreos^{1,*}

¹Pharmacology Department, University of Patras Medical School, Rio Achaias, TK 26500, Greece; ²ICANalytics, ICAN, 83 Bd de l'hopital, 75013 Paris; ³Department of Pharmacy, University of Patras, 26504, Patras, Greece; ⁴INSERM UMR_S 1166 – ICAN, Faculté de Médecine Pitié-Salpêtrière, 91 Bd de l'Hôpital, 75013 Paris, France; ⁵Laboratory of Biochemistry, Department of Chemistry, University of Ioannina, 45110 Ioannina, Greece.

SUPPLEMENTAL TABLE S1

TABLE S1. Primers used in the real -time PCR analysis.

Primer Name	Sequence (5'-3')
mapoa-I_F	GGAGCTGCAAGGGAGACTGT
mapoa-I_R	TGCGCAGAGAGTCTACGTGTGT
mapoe_F	CTGACAGGATGCCTAGCCG
mapoe_R	CGCAGGTAATCCCAGAAGC
mapoa2_F	GCAGACGGACCGGATATGC
mapoa2_R	GCTGCTCGTGTGTCTTCTCA
mapoc1_F	TCCTGTCCTGATTGTGGTCGT
mapoc1_R	CCAAAGTGTTCCCAAACCTCCTT