

Pleiotropic Effects of Apolipoprotein C3 on HDL functionality and Adipose Tissue

Metabolic Activity

Evangelia Zvintzou¹, Marie Lhomme², Stella Chasapi³, Serafoula Filou¹, Vassilis Theodoropoulos¹, Eva Xapapadaki¹, Anatol Kontush⁴, George Spyroulias³, Constantinos C. Tellis⁵, Alexandros D. Tselepis⁵, Caterina Constantinou¹ and Kyriakos E. Kypreos^{1,*}

¹Pharmacology Department, University of Patras Medical School, Rio Achaias, TK 26500, Greece; ²ICANalytics, ICAN, 83 Bd de l'hopital, 75013 Paris; ³Department of Pharmacy, University of Patras, 26504, Patras, Greece; ⁴NSERM UMR_S 1166 – ICAN, Faculté de Médecine Pitié-Salpêtrière, 91 Bd de l'Hôpital, 75013 Paris, France; ⁵Laboratory of Biochemistry, Department of Chemistry, University of Ioannina, 45110 Ioannina, Greece.

SUPPLEMENTAL TABLE S2

TABLE S2. Lipid classes in HDL expressed as mol% of total lipids

	APOC3-HDL_1	APOC3-HDL_2	APOC3-HDL_3	control-HDL_1	control-HDL_2	control-HDL_3
LPC mol%	8,84	8,45	7,42	5,59	5,27	5,72
LPE mol%	0,34	0,45	0,41	0,20	0,16	0,17
PA mol%	0,16	0,17	0,18	0,24	0,23	0,24
PS mol%	0,06	0,06	0,06	0,01	0,01	0,02
PG mol%	0,01	0,01	0,01	0,01	0,01	0,01
PI mol%	2,53	2,90	2,82	4,25	4,21	4,77
PE mol%	1,41	1,30	1,36	0,77	0,74	0,73
PC mol%	80,29	80,98	81,44	83,49	84,23	82,70
SM mol%	6,26	5,60	6,21	5,37	5,07	5,57
Cer mol%	0,10	0,10	0,09	0,07	0,07	0,08

PC, Phosphatidylcholine; LPC, Lysophosphatidylcholine; SM, Sphingomyelin; PI, Phosphatidylinositol; PE, Phosphatidylethanolamine; LPE, Lysophosphatidylethanolamine; PA, Phosphatidic acid; PS, Phosphatidylserine; PG, Phosphatidylglycerol; Cer, Ceramide