

***Pseudomonas aeruginosa* ExIA and *Serratia marcescens* ShIA trigger cadherin cleavage by promoting calcium influx and ADAM10 activation**

Emeline Reboud¹, Stéphanie Bouillot¹, Sabine Patot², Benoît Béganton¹, Ina Attrée¹ and Philippe

Huber^{1*}

Supporting Information

S1 Table: *P. aeruginosa*, *S. marcescens* and *E. coli* strains used in this work

Strains	Description	Abbreviations	Refs
<i>Pseudomonas aeruginosa</i>			
PAO1F (RP1831)	Wild-type laboratory strain T3SS+ (<i>exoS,T,Y</i>)	PAO1F	[1]
CHA	Wild-type laboratory strain T3SS+ (<i>exoS,T,Y</i>)	CHA	[2]
PA14	Wild-type laboratory strain T3SS+ (<i>exoU,T,Y</i>)	PA14	[3]
CLJ1	Clinical strain T3SS-, ExlA+	CLJ1	[4]
IHMA879472	Clinical strain T3SS-, ExlA+	IHMA87	[5]
IHMA87Δ <i>exlA</i>	IHMA87 with <i>exlA</i> inactivated	IHMA87Δ <i>exlA</i>	[6]
IHMA87Δ <i>exlA/exlA</i>	IHMA87Δ <i>exlA</i> with plasmidic expression of <i>exlA</i>	IHMA87Δ <i>exlA/exlA</i>	This work
IHMA87Δ <i>exlA::exlBA</i>	IHMA87Δ <i>exlA</i> with chromosomal insertion of <i>exlBA</i>	IHMA87Δ <i>exlA::exlBA</i>	[6]
IHMA87Δ <i>pilA</i>	IHMA87 with <i>pilA</i> deletion	IHMA87Δ <i>pilA</i>	[6]
<i>Serratia marcescens</i>			
Db11	Clinical strain expressing ShlA		[7]
21C4	<i>shlB</i> mutant of Db11 with no secretion of ShlA		[8]
<i>Escherichia coli</i>			
TOP10 pSW196	Strain containing pSW196 vector only	<i>E. coli</i> -vector	[4]
TOP10 pSW196 <i>exlBA</i>	Strain expressing the <i>exlBA</i> operon	<i>E. coli-exlBA</i>	[4]

References

1. Cisz M, Lee PC, Rietsch A (2008) ExoS controls the cell contact-mediated switch to effector secretion in *Pseudomonas aeruginosa*. *J Bacteriol* 190: 2726-2738.
2. Toussaint B, Delic-Attree I, Vignais PM (1993) *Pseudomonas aeruginosa* contains an IHF-like protein that binds to the *algD* promoter. *Biochem Biophys Res Commun* 196: 416-421.
3. Berthelot P, Attree I, Plesiat P, Chabert J, de Bentzmann S, et al. (2003) Genotypic and phenotypic analysis of type III secretion system in a cohort of *Pseudomonas aeruginosa* bacteremia

isolates: evidence for a possible association between O serotypes and exo genes. *J Infect Dis* 188: 512-518.

4. Elsen S, Huber P, Bouillot S, Coute Y, Fournier P, et al. (2014) A type III secretion negative clinical strain of *Pseudomonas aeruginosa* employs a two-partner secreted exolysin to induce hemorrhagic pneumonia. *Cell Host Microbe* 15: 164-176.
5. Kos VN, Deraspe M, McLaughlin RE, Whiteaker JD, Roy PH, et al. (2015) The resistome of *Pseudomonas aeruginosa* in relationship to phenotypic susceptibility. *Antimicrob Agents Chemother* 59: 427-436.
6. Basso P, Ragno M, Elsen S, Reboud E, Golovkine G, et al. (2017) *Pseudomonas aeruginosa* Pore-Forming Exolysin and Type IV Pili Cooperate To Induce Host Cell Lysis. *MBio* 8.
7. Flyg C, Kenne K, Boman HG (1980) Insect pathogenic properties of *Serratia marcescens*: phage-resistant mutants with a decreased resistance to *Cecropia* immunity and a decreased virulence to *Drosophila*. *J Gen Microbiol* 120: 173-181.
8. Kurz CL, Chauvet S, Andres E, Aurouze M, Vallet I, et al. (2003) Virulence factors of the human opportunistic pathogen *Serratia marcescens* identified by in vivo screening. *EMBO J* 22: 1451-1460.

Supplemental Method

Bacterial transmigration across Transwell devices

Three days before infection, A549 cells (6×10^4) were seeded in DMEM medium on Transwell filters (Greiner ThinCert 24 well, 3.0 μm). At day 0, the medium was replaced and bacteria (MOI 10) were added in the upper compartment. At 4 h.p.i., 100 μL of medium in the lower compartment were withdrawn, serially diluted and spotted onto LB-agar plates to determine the number of transmigrated bacteria.