

Supplement table:

Table 11: Characteristics of patients not included in study (n=73) or CT scans (n=12) were not analyzable.

Categories		N (%) No scan/no data	N (%) Patient's volume prevents evaluation of scan
Age			
	Mean (SD)	64.4 (13.1)	59.5 (10.0)
	< 61 years	29 (25.0)	7 (58.3)
	>61 years	87 (75.0)	5 (41.7)
Sex			
	Male	37 (50.7)	6 (50.0)
	Female	36 (49.3)	6 (50.0)
BMI (kg/m²)			
	Mean (SD)	25.1 (3.9)	32.4 (4.13)
	Underweight (<18.5)	2 (2.7)	0 (0.0)

	Normal weight (18.5 – 24.99)	27 (37.0)	1 (8.3)
	Overweight (25 – 29.99)	28 (38.4)	3 (25.0)
	Obese Class I (30 – 34.99)	9 (12.3)	5 (41.7)
	Obese Class II & III (> 34.99)	1 (1.4)	3 (25.0)
	Unknown	6 (8.2)	0 (0.0)
Site of CRC			
	Colon	42 (57.5)	7 (58.3)
	Recto sigmoid	5 (6.9)	1 (8.3)
	Rectum	20 (27.4)	4 (33.3)
	Unknown	6 (8.2)	0 (0.0)
Stage			
	0	0 (0.0)	0 (0.0)
	I	21 (28.8)	3 (37.3)
	II	19 (26.0)	4 (36.4)
	III	13 (17.8)	2 (18.2)
	IV	3 (4.1)	0 (0.0)
	Unknown	17 (23.3)	2 (18.2)
Time of CT Scan			
	Pre-op	5 (6.9)	11 (91.7)
	Post-op	5 (6.9)	1 (8.3)
	No-scan	63 (86.3)	0 (0.0)
Metabolic Equivalents			
	Unknown	73	12