

Supplemental Table S3. Coagulation and anticoagulation factor levels according to WHO bleeding grade in patients with severe thrombocytopenia (platelet count $\leq 10 \times 10^9/L$)

Factors	Total patients with platelet $\leq 10 \times 10^9/L$ (n = 13)	WHO bleeding grade ≥ 2 (n = 7)	WHO bleeding grade 0 or 1 (n = 6)	P value*
Fibrinogen (mg/dL)	311.2 (185.2-390.7)	281.9 (185.2-345.9)	323.6 (265.8-390.7)	0.153
Factor II (%)	86.2 (74.3-116.0)	86.2 (74.3-113.2)	94.7 (75.7-116.0)	0.668
Factor V (%)	108.2 (75.8-138.3)	109.9 (86.3-138.3)	91.9 (75.8-118.2)	0.074
Factor VII (%)	113.7 (57.9-153.1)	113.7 (57.9-124.5)	112.2 (93.2-153.1)	0.253
Factor VIII (%)	135.8 (91.6-230.5)	114.0 (91.6-154.9)	149.4 (94.8-230.5)	0.199
Factor IX (%)	116.1 (51.1-150.2)	115.0 (51.1-150.2)	119.5 (100.7-146.9)	0.391
Factor X (%)	75.0 (57.3-113.7)	74.8 (57.3-113.7)	82.2 (67.8-112.4)	0.568
Factor XI (%)	83.0 (57.0-127.4)	83.0 (63.3-127.4)	73.8 (57.0-124.1)	0.391
Factor XII (%)	60.0 (34.9-129.8)	63.9 (40.8-129.8)	55.1 (34.9-74.6)	0.568
AT (%)	81.4 (67.5-124.6)	90.8 (67.5-106.0)	79.1 (67.7-124.6)	0.775
PC (%)	117.9 (57.9-148.2)	92.6 (57.9-148.2)	118.9 (90.6-130.2)	0.568
PS (%)	86.7 (52.3-179.8)	86.7 (71.9-102.7)	86.6 (52.3-179.8)	0.999

*P value: Man-Whitney *U* test for results from patients with WHO bleeding grading ≥ 2 or those with WHO bleeding grade 0 or 1.

Values are expressed as median (range).

Abbreviations: AT, antithrombin; PC, protein C; PS, protein S.