

BMJ Open

Daycare attendance and respiratory tract infections: a prospective birth cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-014635
Article Type:	Research
Date Submitted by the Author:	07-Oct-2016
Complete List of Authors:	Schuez-Havupalo, Linnea; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research Toivonen, Laura; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research Karppinen, Sinikka; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research Kaljonen, Anne; Turun Yliopisto, Turku Institute for Child and Youth Research Peltola, Ville; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research
Primary Subject Heading:	Paediatrics
Secondary Subject Heading:	Infectious diseases
Keywords:	Respiratory infections < THORACIC MEDICINE, Paediatric infectious disease & immunisation < PAEDIATRICALS, Community child health < PAEDIATRICALS

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Daycare attendance and respiratory tract infections: a prospective birth cohort study

Linnea Schuez-Havupalo, MD,^{1,2} Laura Toivonen, MD,^{1,2} Sinikka Karppinen, MD,^{1,2} Anne Kaljonen, BSc,² and Ville Peltola, MD, PhD^{1,2}

Affiliations: ¹Department of Paediatrics and Adolescent Medicine, Turku University Hospital; and ²Turku Institute for Child and Youth Research, University of Turku, Turku, Finland

Correspondence to: Ville Peltola, MD, PhD, Department of Paediatrics and Adolescent Medicine, Turku University Hospital, 20521 Turku, Finland; ville.peltola@utu.fi; tel. +358-2-3130000; fax: +358-2-3131460.

Keywords: daycare, respiratory infections, children, the STEPS Study

Word count: 2499

Abstract

Objective. We explored the burden of respiratory tract infections in young children with regard to daycare initiation.

Design. Longitudinal prospective birth cohort study.

Setting and method. We recruited 1827 children for follow-up until the age of 24 months collecting diary data on respiratory tract infections and daycare. Children with continuous daycare type and complete data were divided into groups of centre-based daycare (n=299), family daycare (n=245), and home care (n=350). Using repeated measures variance analyses, we analysed days per month with symptoms of respiratory tract infection, antibiotic treatments, and parental absence from work for a period of 6 months prior to and 9 months after the start of daycare.

Results. We documented a significant effect of time and type of daycare, as well as a significant interaction between them for all outcome measures. There was a rise in mean days with symptoms from 3.79 (95% CI 3.04-4.53) during the month preceding centre-based daycare to 10.57 (95% CI 9.35-11.79) at 2 months after the start of centre-based daycare, with a subsequent decrease within the following 9 months. Similar patterns with a rise and decline were observed in the use of antibiotics and parental absences. The start of family daycare had weaker effects. Our findings were not changed when taking into account confounding factors.

Conclusions. Our study shows the rapid increase in respiratory infections after start of daycare and a relatively fast decline in the course of time with continued daycare. It is important to support families around the beginning of daycare.

Article summary

Strength and limitations of this study

- We prospectively collected detailed day-to-day diary data on respiratory tract infections and related outcomes from 0 to 2 years of age in a birth cohort population, and documented daycare type and starting date.
- The study design allowed us to analyze the time-dependent effects of centre-based daycare and family daycare on the days per month with symptoms of a respiratory infection in comparison with children of same age in home care.
- Different confounders were taken into account, but they did not change our findings of a peak in the rate of respiratory tract infections shortly after daycare initiation, and a clear decline thereafter.
- A limitation is that, due to strict requirements regarding detail in follow up and careful exclusion of missing data, we lost a substantial proportion of cases.

Introduction

In modern society a considerable proportion of children attend daycare from the age of less than two years. Respiratory tract infections (RTIs) in this age group constitute an important health problem. In addition to causing stress to children and families, they also affect transmission of pathogens to other age groups, parental absences from work with ensuing economic consequences, and rates of anti-microbial medication use with subsequent impact on resistance patterns.[1-3] Previous studies suggest that increased rates of infections may be a transient problem related to the beginning of daycare,[4-7] but results are conflicting. Few prospective cohort studies have examined the topic, and lose frequency of follow-up has limited chronological conclusions. Some studies have found a protective effect of starting daycare early with regard to RTIs later in life.[4, 8]

Families, child health care professionals, and daycare providers would benefit from more detailed knowledge about the burden of disease as a function of time before and after the start of daycare in the above mentioned age group. We examined this in a prospectively followed birth cohort.

Methods

Study population and conduct

We used the cohort of the observational 'Steps to the Healthy Development and Well-being of Children' Study (The STEPS Study), which consists of 1827 children from 1797 families.[9] Recruitment occurred in two stages from women with a live birth between 1 January 2008 and 31 March 2010 in the Southwest Finland Hospital District (n=9936). During the first stage, 1387 families were recruited through community midwifery services during pregnancy, and a further 410 families joined the study soon after the birth of their child. In this study we collected questionnaire-based data on family related factors and

1
2
3 daycare arrangements at gestational week 20, and at ages 13 months, 18 months, and 24
4
5 months, as applicable. Families kept study diaries that recorded health related factors, the
6
7 precise starting date of daycare, and data from physician visits. Parents documented the
8
9 child's symptoms, antibiotic treatments, and parental absence from work on a day-to-day
10
11 basis in these diaries. Parents were instructed to also mark symptom-free days in the diary.
12
13 We differentiated missing from negative data by excluding follow-up with no diary markings.
14
15

16 Families attended our nurse-led study clinic when the child was 2 and 13 months of
17
18 age. During RTIs part of the cohort (n=982) were also followed up by our physician-led study
19
20 clinic. Recruitment of children to our study is shown in figure 1. Exclusion criteria for this
21
22 study constituted discontinuous daycare during our follow-up, or insufficient information on
23
24 daycare arrangements including missing knowledge on the exact time of start of daycare. The
25
26 Ethics Committee of the Hospital District of Southwest Finland approved the study protocol.
27
28 The parents of the participating children gave their written, informed consent.
29
30
31
32
33

34 **Daycare attendance**

35
36 Children were categorized into three different groups according to daycare arrangements:
37
38 children cared for by parents or relatives (comparison group), children attending family
39
40 daycare (FDC), and children attending daycare centres (DCC). In this study family daycare
41
42 was defined as daycare provided by a trained carer in her or his home, or in the children's
43
44 homes using a rotational system. According to local regulations, FDC group sizes included
45
46 no more than 5 children. In rare cases, carers worked together forming a common nursery.
47
48 Since group sizes tended to be relatively small in those nurseries, these children were
49
50 included in the FDC group. Daycare centres were defined by larger-group, centre-based care
51
52 provided by several professional carers. FDC and care in DCCs were provided either by the
53
54 municipality or on a private basis.
55
56
57
58
59
60

Outcomes

The outcomes of the study were symptoms of RTIs (cough, rhinorrhea, fever, or wheeze), consumption of antibiotic medication, and parental absence from work as recorded on a day-to-day basis in the study diaries. Antibiotic medications prescribed for any reason were included, but the main indication was acute otitis media. Any parental absence from work due to the child's illness was recorded, and reasons were not limited to RTIs. Final outcome measures constituted days per month with RTI symptoms (sick days), antibiotic medication, and parental absence from work.

Confounding factors

The following potential confounders were taken into consideration: sex, siblings, season of year at start of daycare, asthma in the parents, pets (cats or dogs at home), and maternal post-secondary education.[10-12] There was only a small proportion of families with known smoking in parents (127 out of 1827), and since our results strongly indicated reporting bias, this variable was excluded from the analyses.

Statistical analysis

All statistical analyses were carried out using SAS version 9.4 (SAS, Cary, NC, USA). In all our analyses p values less than 0.05 were regarded as statistically significant. In the first step, outcome variables were calculated as days per month for each child and aligned in a time sequence relating to the beginning of daycare. Day-specific chronological follow-up included a time of 6 months previous to and 9 months after the beginning of daycare. Children commenced daycare at different ages, so that not all above defined data related to the fixed age frame of 0-2 years. Data from outside the agreed age frame were excluded from the

1
2
3 analysis using the SAS MIXED procedure. Chronological data for the comparison group
4
5 were obtained from children at equivalent ages not attending daycare. The equivalent age
6
7 range was calculated using the mean age of starting daycare in the FDC and DCC groups.
8
9 According to this mean age, follow-up of children within the comparison group included the
10
11 age range of 9 to 24 months (6 months prior to and 9 months after the mean start of daycare
12
13 at 15 months).

14
15
16 Data were analyzed by repeated measures variance analysis comparing all three
17
18 groups for type of daycare and time, as well as their interaction. In a second step,
19
20 chronological thresholds of significance were obtained by comparing all p values for the
21
22 difference of least square means between all pairs of analyzed months. We determined the
23
24 shortest time for any given trend with a $p < 0.05$. Although the distributions of monthly
25
26 outcome variables (sick days, days with antibiotic treatment, and parental absence from
27
28 work) were skewed in our analyses (coefficient of skewness up to 2.4 and kurtosis 7.6), the
29
30 pairwise difference variables followed approximately a normal distribution.
31
32

33
34 In a third step, repeated measures variance analysis was repeated according to the
35
36 previous model, but stratifying according to individual confounding factors.
37
38

39 40 41 **Results**

42
43 Figure 1 shows numbers of children within follow-up. Within our cohort of 1570 children in
44
45 active follow-up, 276 (21.8%) of 1264 children with data on daycare arrangements at age 13
46
47 months attended daycare (11.7% in DCC, 10.1% in FDC) and 593 (55.0%) of 1079 children
48
49 attended daycare at the age of 24 months (29.5% in DCC and 25.0% in FDC). At all ages the
50
51 vast majority of children in daycare spent over 5 hours per day at daycare (88.5% at age 13
52
53 months and 91.6% at age 24 months). Group sizes were clearly smaller for children attending
54
55 FDC as compared to DCC, with 88.3% (at 13 months) to 82.7% (at 24 months) of families
56
57
58
59
60

1
2
3 reporting a group size of less than 5 children per group. In DCC group sizes were variable
4
5 with the following results for the ages of 13 months (and 24 months): less than 5 children for
6
7 2.0%, 5-15 children for 86.8%, over 15 children for 11.1% (less than 5 children for 1.4%, 5-
8
9 15 children for 87.8%, over 15 children for 10.9%). Mean age at the beginning of daycare
10
11 was 15 months (SD 4.1). Sex was evenly distributed throughout the daycare groups with
12
13 49.1% of boys in the comparison group, 50.6% in FDC, and 54.2% in DCC. Start of daycare
14
15 (FDC or DCC) occurred in 38.2% of cases during fall-winter (October to March) and in
16
17 61.8% during spring-summer (April to September). Within the comparison, FDC, and DCC
18
19 groups, 97.5% of children were born full-term (gestational age equal to, or over 37 weeks),
20
21 with only 5/894 children (0.6%) having been born at a gestational age of less than 35 weeks.
22
23
24

25
26 There was a peak of mean sick days per month at 2 months after the start of daycare
27
28 for both FDC and DCC (figure 2). The rise in sick days was stronger for the DCC group than
29
30 for the FDC group. We observed increases from 3.53 mean sick days (95% CI 2.83-4.24)
31
32 during the month prior to the start of daycare to 8.34 mean sick days per month (95% CI
33
34 7.25-9.43) for the FDC group, and from 3.79 (95% CI 3.04-4.53) mean sick days prior to the
35
36 start of daycare to 10.57 (95% CI 9.35-11.79) mean sick days per month for the DCC group.
37
38 For both of these groups, the rise of outcome measures was a transient phenomenon; sick
39
40 days were comparable to the baseline group at 5 months after the start of daycare. Antibiotic
41
42 use (figure 3) and parental absences from work because of children's illnesses (figure 4) rose
43
44 and declined according to a pattern similar to that observed in sick days in relation to the start
45
46 of daycare, although this decline was less pronounced compared to sick days.
47
48

49
50 The shortest time for a significant rise of outcome measures was determined from the
51
52 last month prior to starting daycare (month -1) and for a significant decline from the peak of
53
54 each outcome variable. The rise in sick days (figure 2) and antibiotic use (figure 3) was more
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

rapid among children starting DCC than in those starting FDC. Within the comparison group there were no significant findings in pairwise comparisons between months.

In repeated measures variance analyses there was a significant overall effect for time and type of daycare and their interaction for all outcome measures (table 1).

For peer review only

Table 1. P values* for the effects of daycare type and time on the rates of RTI symptoms (sick days), antibiotic treatments, and parental absence from work because of a child's illness

	Sick days per month, p	Days with antibiotic treatment per month, p	Parental absence from work, days per month, p
Time (month)	<0.0001	0.004	0.003
Daycare type	<0.0001	<0.001	0.02
Contrast: comparison - FDC groups	0.55	0.18	ND
Contrast: comparison - DCC groups	<0.0001	<0.001	ND
Contrast: FDC - DCC groups	<0.001	0.04	ND
Interaction: time and daycare type	<0.0001	<0.0001	0.003
Time (month) for comparison group	0.005	0.64	ND
Time (month) for FDC group	<0.0001	0.002	<0.0001
Time (month) for DCC group	<0.0001	<0.0001	<0.0001

ND, not determined; RTI, respiratory tract infection; FDC, family daycare; DCC, daycare centre.

*By repeated measures variance analysis.

1
2
3 This interaction showed the above described pattern of rise and decline, which was strongest
4
5 for the DCC group and weaker for the FDC group. Within the comparison group there was no
6
7 such pattern, but significant variation was observed in the frequency of sick days per month
8
9 over time. When analyzing contrasts in sick days and antibiotic use between different types
10
11 of daycare without the aspect of time, only the DCC group reached significance in
12
13 comparison to the baseline group and FDC group.
14
15

16 The presence of older siblings in the family and higher post-secondary education in
17
18 mothers were associated with a higher burden of disease (table 2).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Effects of confounding factors on the rates of RTI symptoms (sick days) according to daycare type.

	Type of daycare						P*	P for interaction with mode of daycare
	Comparison group		Family daycare		Daycare centre			
	n/N (%)	Sick days per month, mean (SD)	n/N (%)	Sick days per month, mean (SD)	n/N (%)	Sick days per month, mean (SD)		
Older siblings							0.02	<0.0001
Yes	162/ 350 (46.3)	5.55 (3.72)	99 /245 (40.4)	5.26 (3.33)	132/ 299 (44.1)	5.10 (3.72)		
No	188/ 350 (53.7)	4.16 (2.95)	146/ 245 (59.6)	4.58 (3.59)	167/ 299 (55.9)	5.88 (4.31)		
Higher post-secondary education in mother							0.009	0.27
Yes	211/ 341 (61.9)	5.12 (3.53)	161/ 239 (67.4)	5.12 (3.72)	203/ 293 (69.3)	5.66 (3.75)		
No	130/ 341 (38.1)	4.40 (3.17)	78 / 239 (32.6)	4.23 (2.97)	90/ 293 (30.7)	5.45 (4.79)		
Start of daycare during fall-winter (October-March)							0.39	0.61
Yes	ND	ND	100/ 245 (40.8)	4.90 (3.15)	108/ 299 (36.1)	5.69 (4.21)		
No	ND	ND	145/ 245 (59.2)	4.82 (3.73)	191/ 299 (63.9)	5.45 (4.0)		
Asthma in parents							0.95	0.72
Yes	35/ 340 (10.3)	4.65 (3.26)	26/ 234 (11.1)	5.54 (4.0)	30/ 287 (10.5)	5.83 (4.25)		
No	305/ 340 (89.7)	4.86 (3.43)	208/ 234 (88.9)	4.76 (3.45)	257/ 287 (89.5)	5.56 (4.08)		
Cat or dog at home							0.31	0.23
Yes	94/ 295 (31.9)	4.43 (3.07)	80/ 196 (40.8)	4.83 (3.07)	57/ 219 (26.0)	5.80 (4.28)		
No	201/ 295 (68.1)	5.14 (3.48)	116/ 196 (59.2)	4.92 (3.14)	162/ 219 (74.0)	5.68 (3.66)		

n/N, number of children per number of those with data available; ND, not determined; RTI, respiratory tract infection.

*By univariate analysis, regardless of daycare type.

1 In repeated measures variance analyses stratified according to confounding factors,
2
3 associations remained ($p < 0.0001$ for the effect of time and $p < 0.001$ for daycare type for all
4
5 analyses, except for those with parental asthma, where $p = 0.004$ for daycare type). Online
6
7 supplementary figures S1 and S2 illustrate the association of start of daycare with respiratory
8
9 tract infections in children stratified according to the presence of siblings and maternal post-
10
11 secondary education. For children without older siblings, the rise of sick days per month after
12
13 start of daycare was more pronounced than for those with older siblings.
14
15
16
17
18
19

20 Discussion

21
22 In this study there was a strong but temporally limited effect of daycare initiation on the rate
23
24 of RTIs. This was strongest for the DCC group and weaker for the FDC group. An effect of
25
26 overall time in daycare,[4-7] as well as daycare type, on infections has been documented in
27
28 previous studies.[5, 13-15] Our study specifically focused on RTI dynamics around the start
29
30 of daycare.
31
32

33 We analyzed three main outcome measures. After showing a peak, the rate of sick
34
35 days returned near to baseline within our 9-month follow-up period from the start of daycare;
36
37 for antibiotic use there was more scatter with a less pronounced decline, and parental absence
38
39 from work did not return to baseline, which was to be expected due to the fact that both
40
41 parents tend to be at work after the start of daycare. Consideration of confounders did not
42
43 change our findings. The effect of older siblings on infections in relation to daycare has been
44
45 previously described,[7] and our findings are in line with previous results.
46
47
48

49 We obtained a small, but significant effect for overall time in our analysis for sick
50
51 days also within the comparison group of children in home care. There was no consistent
52
53 trend over time, but rather a random variation between different months in the comparison
54
55 group. The effects of virus epidemics and season of year were minimized by the way the
56
57 baseline group was formed, but they cannot be completely excluded.
58
59
60

1
2 We consider the main strength of our study the detailed follow-up on a day-to-day
3 basis for a relatively large number of cases and in chronological relation to the start of
4 daycare. Previous cohort studies mostly assessed data from follow-up with intervals of
5 months or longer periods and without information on the exact starting point of daycare, such
6 that detailed chronological analyses for the time around the start of daycare could not be
7 carried out.[5, 6, 8, 12] Retrospective collection of data relating to symptoms poses problems
8 in some other studies, since it strongly relies on parents' memories.
9

10
11
12
13
14
15
16
17 There are some limitations to our study. Follow-up time was relatively short albeit
18 sufficient to demonstrate the above described effects. Outcome measures were based on self-
19 reported data and are thus subject to bias. Daily diary-keeping demanded regular participation
20 of families, although diaries were concise and easily completed. There were several indicators
21 that, for the small number of families with known smoking, diaries were not completed as
22 comprehensively as in the rest of the cohort, so that we had to exclude parental smoking as a
23 confounder from our study. The group with more detailed follow-up during infections had
24 access to our medical services free of charge, and thus may have been more motivated to
25 accurately document all experienced events. Inclusion into this group was offered to all
26 families without any selection criteria.
27
28
29
30
31
32
33
34
35
36
37
38

39
40 Due to strict requirements for comprehensive follow-up and careful exclusion of
41 missing data, we lost a considerable proportion of cases. In order to screen for bias, a
42 comparison of background variables was performed for non-responders and responders at 13
43 months of age, and there were only minor differences between groups.[9]
44
45
46
47

48
49 Our cohort children had less often older siblings than all eligible children, and their
50 mothers had higher education and occupational status.[9] Otherwise our cohort represents the
51 Southwest Finnish population well. Although lower socioeconomic class has traditionally
52 been linked to an increased rate of infectious diseases,[16] a Swedish study found an
53 association between low social status, smaller likelihood to be cared for in out-of-home care,
54
55
56
57
58
59
60

1 and lesser consumption of medical services.[17] Within our cohort, social status, as indicated
2
3
4 by maternal education, was also reflected by a higher rate of RTIs, but this did not affect our
5
6 findings relating to daycare.
7
8
9

10 **Conclusion**

11
12 In this longitudinal cohort study, the respiratory infectious disease burden was clearly related
13
14 to out-of-home care, but it decreased already within a short follow-up time of 9 months after
15
16 the start of daycare. These findings have implications on a correctly focused approach when
17
18 supporting families with small children. Families, daycare providers and pediatricians may be
19
20 reassured of the transient nature of increased RTIs after the start of daycare.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole STEPS Study team, which includes our research physicians and study nurses.

Footnotes

Competing interests. The authors have no competing interests.

Funding. All phases of this study were supported by the University of Turku; the Åbo Akademi University; the Turku University Hospital; the Academy of Finland, Grants no. 123571, 140251, and 277535; the Foundation for Pediatric Research, Finland; and Research Funds from Specified Government Transfers, Hospital District of Southwest Finland. Dr. Schuez-Havupalo was also supported by the Turku University Foundation.

Data sharing statement: Enquiries regarding extra data can be emailed to Ville Peltola.

Contributors. LSH contributed to conception and design, the clinical follow-up and data collection, carried out data analyses, drafted the initial manuscript, and approved the final manuscript as submitted. LT and SK contributed to conception and design, the clinical follow-up and data collection, reviewed and revised the manuscript, and approved the final manuscript as submitted. AK contributed to conception and design, coordinated data collection, carried out data analyses, reviewed and revised the manuscript, and approved the final manuscript as submitted. VP conceptualized and designed the study, coordinated and supervised data collection and analyses, reviewed and revised the manuscript, and approved the final manuscript as submitted.

References

- 1 Nurmi T, Salminen E, Pönkä A. Infections and other illnesses of children in day-care centers in Helsinki. II: The economic losses. *Infection* 1991;19:331-5.
- 2 Holmes SJ, Morrow AL, Pickering LK. Child-care practices: effects of social change on the epidemiology of infectious diseases and antibiotic resistance. *Epidemiol Rev* 1996;18:10-28.
- 3 Thrane N, Olesen C, Mortensen JT, et al. Influence of day care attendance on the use of systemic antibiotics in 0- to 2-year-old children. *Pediatrics* 2001;107:e76.
- 4 Hurwitz ES, Gunn WJ, Pinsky PF, et al. Risk of respiratory illness associated with day-care attendance: a nationwide study. *Pediatrics* 1991;87:62-9.
- 5 The National Institute of Child Health and Human Development Early Child Care Research Network. Child care and common communicable illnesses: results from the National Institute of Child Health and Human Development Study of Early Child Care. *Arch Pediatr Adolesc Med* 2001;155:481-8.
- 6 Zutavern A, Rzehak P, Brockow I, et al. Day care in relation to respiratory-tract and gastrointestinal infections in a German birth cohort study. *Acta Paediatr* 2007;96:1494-9.
- 7 Kamper-Jørgensen M, Wohlfahrt J, Simonsen J, et al. Population-based study of the impact of childcare attendance on hospitalizations for acute respiratory infections. *Pediatrics* 2006;118:1439-46.
- 8 Cote S, Petitclerc A, Raynault M-F, et al. Short- and long-term risk of infections as a function of group child care attendance. *Arch Pediatr Adolesc Med* 2010;164:1132-7.
- 9 Lagström H, Rautava P, Kaljonen A, et al. Cohort profile: Steps to the healthy development and well-being of children (the STEPS study). *Int J Epidemiol* 2013;42:1273-84.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 10 Hatakka K, Piirainen L, Pohjavuori S, et al. Factors associated with acute respiratory illness in day care children. *Scand J Infect Dis* 2010;42:704-11.
- 11 Koopman LP, Smit HA, Heijnen ML, et al. Respiratory infections in infants: interaction of parental allergy, child care, and siblings - the PIAMA study. *Pediatrics* 2001;108:943-8.
- 12 Celedon J, Litonjua A, Weiss S, et al. Day care attendance in the first year of life and illness of the upper and lower respiratory tract in children with a familial history of atopy. *Pediatrics* 1999;104:495-500.
- 13 Hardy AM, Fowler MG. Child care arrangements and repeated ear infections in young children. *Am J Public Health* 1993;83:1321-5.
- 14 Marx J, Osguthorpe J, Parsons G. Day care and the incidence of otitis media in young children. *Otolaryngol Head Neck Surg* 1995;112:695-9.
- 15 Louhiala P, Jaakkola N, Ruotsalainen R, et al. Form of day care and respiratory infections among Finnish children. *Am J Public Health* 1995;85:1109-12.
- 16 Cohen S. Social status and susceptibility to respiratory infections. *Ann N Y Acad Sci* 1999;896:246-53.
- 17 Hjern A, Haglund B, Rasmussen F, et al. Socio-economic differences in daycare arrangements and use of medical care and antibiotics in Swedish preschool children. *Acta Paediatr* 2000;89:1250-6.

Figure legends

Figure 1. Flow-diagram of the recruitment and follow-up of study children.

Figure 2. Means and 95% CIs (dashed lines) of sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month).

Figure 3. Means and 95% CIs (dashed lines) of days with antibiotic treatment per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month). Dashed lines: transiently below the significance level.

Figure 4. Means and 95% CIs (dashed lines) of days with parental absence from work per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month). Dashed lines: transiently below the significance level.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1




Figure 2

Peer review only




Figure 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4

Peer review only

Supplemental material: figures S1 and S2.





Figure S1. Mean sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Data are shown for children with older siblings (A) and for children without older siblings (B).

A



B




Figure S2. Mean sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Data are shown for children of mothers with lower (A) and higher (B) post-secondary education.

BMJ Open

Daycare attendance and respiratory tract infections: a prospective birth cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-014635.R1
Article Type:	Research
Date Submitted by the Author:	06-May-2017
Complete List of Authors:	Schuez-Havupalo, Linnea; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research Toivonen, Laura; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research Karppinen, Sinikka; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research Kaljonen, Anne; Turun Yliopisto, Turku Institute for Child and Youth Research Peltola, Ville; Turku University Hospital, Department of Paediatrics and Adolescent Medicine; Turun Yliopisto, Turku Institute for Child and Youth Research
Primary Subject Heading:	Paediatrics
Secondary Subject Heading:	Infectious diseases
Keywords:	Respiratory infections < THORACIC MEDICINE, Paediatric infectious disease & immunisation < PAEDIATRICALS, Community child health < PAEDIATRICALS

SCHOLARONE™
Manuscripts

1
2
3 Manuscript ID bmjopen-2016-014635, revision 1
4

5
6 **Daycare attendance and respiratory tract infections: a**
7
8
9 **prospective birth cohort study**
10

11 Linnea Schuez-Havupalo, MD,^{1,2} Laura Toivonen, MD,^{1,2} Sinikka Karppinen, MD,^{1,2} Anne
12 Kaljonen, BSc,² and Ville Peltola, MD, PhD^{1,2}
13
14
15
16

17
18 **Affiliations:** ¹Department of Paediatrics and Adolescent Medicine, Turku University
19 Hospital; and ²Turku Institute for Child and Youth Research, University of Turku, Turku,
20
21 Finland
22
23
24

25
26
27 **Correspondence to:** Ville Peltola, MD, PhD, Department of Paediatrics and Adolescent
28 Medicine, Turku University Hospital, 20521 Turku, Finland; ville.peltola@utu.fi; tel. +358-
29
30 2-3130000; fax: +358-2-3131460.
31
32
33

34
35
36 **Keywords:** daycare, respiratory infections, children, the STEPS Study
37
38
39

40 **Word count:** 2951
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective. We explored the burden of respiratory tract infections in young children with regard to daycare initiation.

Design. Longitudinal prospective birth cohort study.

Setting and method. We recruited 1827 children for follow-up until the age of 24 months collecting diary data on respiratory tract infections and daycare. Children with continuous daycare type and complete data were divided into groups of centre-based daycare (n=299), family daycare (n=245), and home care (n=350). Using repeated measures variance analyses, we analysed days per month with symptoms of respiratory tract infection, antibiotic treatments, and parental absence from work for a period of 6 months prior to and 9 months after the start of daycare.

Results. We documented a significant effect of time and type of daycare, as well as a significant interaction between them for all outcome measures. There was a rise in mean days with symptoms from 3.79 (95% CI 3.04-4.53) during the month preceding centre-based daycare to 10.57 (95% CI 9.35-11.79) at 2 months after the start of centre-based daycare, with a subsequent decrease within the following 9 months. Similar patterns with a rise and decline were observed in the use of antibiotics and parental absences. The start of family daycare had weaker effects. Our findings were not changed when taking into account confounding factors.

Conclusions. Our study shows the rapid increase in respiratory infections after start of daycare and a relatively fast decline in the course of time with continued daycare. It is important to support families around the beginning of daycare.

Article summary

Strength and limitations of this study

- We prospectively collected detailed day-to-day diary data on respiratory tract infections and related outcomes from 0 to 2 years of age in a birth cohort population, and documented daycare type and starting date.
- The study design allowed us to analyze the time-dependent effects of centre-based daycare and family daycare on the days per month with symptoms of a respiratory infection in comparison with children of same age in home care.
- Different confounders were taken into account, but they did not change our findings of a peak in the rate of respiratory tract infections shortly after daycare initiation, and a clear decline thereafter.
- A limitation is that, due to strict requirements regarding detail in follow up, we lost a substantial proportion of cases.

Introduction

Daycare has been known to be a major risk factor for respiratory tract infections (RTIs) in children for over 30 years.[1-4] A considerable proportion of children in modern society attend daycare from the age of less than two years. RTIs in this age group constitute an important health problem. In addition to causing stress to children and families, they also affect transmission of pathogens to other age groups, parental absences from work with ensuing economic consequences, and rates of anti-microbial medication use with subsequent impact on resistance patterns.[5-7]

Within the daycare setting, factors reflecting contact rates with other children have consistently been identified as important determinants of infection risk. These factors include group sizes, or size and type of the daycare facility,[8-12] as well as weekly exposure time.[13] A number of studies have shown children of young age to be particularly vulnerable to daycare-related effects regarding transmission of respiratory virus infections,[8, 13-17] and early daycare has also been linked to more severe or long-term health problems, such as recurrent acute otitis media (AOM), high numbers of antibiotic medications during early childhood, an increased lifetime risk of asthma, and an increased risk of invasive pneumococcal infections.[18-20] In their cross-sectional study, Hurwitz et al. were the first to specifically demonstrate a lower daycare-attributable risk of RTIs with increased time after daycare initiation, and thus provided data to suggest that daycare-related infection risks were not exclusively linked to the absolute age of a child.[17] A large register-based study assessing RTI-related hospitalizations in under-school aged children confirmed a decrease of hospitalizations with an increased time in daycare of over 6 months.[14] Only a few larger-size longitudinal studies have assessed the effects of overall exposure time to daycare in relation to RTIs,[15, 16, 21] and lose frequency of follow-up has limited the conclusions of

1
2
3 chronological variations. Some studies have found a protective effect of starting daycare
4
5 early with regard to RTIs later in life.[16, 21]
6

7
8 Given the discrepancy between potential risks and benefits of early daycare in
9
10 children under the age of 2 years, detailed longitudinal studies assessing the burden of disease
11
12 as a function of time before and after the start of daycare are needed. Families, child health
13
14 care professionals, and daycare providers would benefit from more detailed knowledge about
15
16 the daycare-related impact of RTIs on this vulnerable age-group over time. We examined our
17
18 hypothesis of a time-limited effect of daycare on RTIs in a prospectively followed birth
19
20 cohort.
21

22 23 24 25 **Methods**

26 27 **Study population and conduct**

28
29 We used the cohort of the observational 'Steps to the Healthy Development and Well-being
30
31 of Children' Study (The STEPS Study), which consists of 1827 children from 1797
32
33 families.[22] Recruitment occurred in two stages from women with a live birth between 1
34
35 January 2008 and 31 March 2010 in the Southwest Finland Hospital District (n=9936).
36
37

38
39 During the first stage, 1387 families were recruited through community midwifery services
40
41 during pregnancy, and a further 410 families joined the study soon after the birth of their
42
43 child. In this study, follow-up included an age frame of 0-2 years, until March 2012. We
44
45 collected questionnaire-based data on family related factors and daycare arrangements at
46
47 gestational week 20, and at ages 13 months, 18 months, and 24 months, as applicable.
48

49
50 Families kept study diaries that recorded health related factors, the precise starting date of
51
52 daycare, and data from physician visits. Parents documented the child's symptoms, antibiotic
53
54 treatments, and parental absence from work on a day-to-day basis in these diaries. Parents
55
56
57
58
59
60

1
2
3 were instructed to also mark symptom-free days in the diary. We differentiated missing from
4
5 negative data by excluding follow-up with no diary markings.
6

7 Families attended our nurse-led study clinic when the child was 2 and 13 months of
8
9 age. During RTIs part of the cohort (n=982) were also followed up by our physician-led study
10
11 clinic.[23] Recruitment of children to our study is shown in figure 1. Exclusion criteria for
12
13 this study constituted discontinued follow-up before daycare initiation, insufficient
14
15 information on daycare arrangements, or a lack of information on home care. The Ethics
16
17 Committee of the Hospital District of Southwest Finland approved the study protocol. The
18
19 parents of the participating children gave their written, informed consent.
20
21
22
23
24

25 **Daycare attendance**

26
27 Children were categorized into three different groups according to daycare arrangements:
28
29 children cared for by parents or relatives (home care group), children attending family
30
31 daycare (FDC), and children attending daycare centres (DCC). In this study family daycare
32
33 was defined as daycare provided by a trained carer in her or his home, or in the children's
34
35 homes using a rotational system. According to local regulations, FDC group sizes included
36
37 no more than 5 children. In rare cases, carers worked together forming a common nursery.
38
39 Since group sizes tended to be relatively small in those nurseries, these children were
40
41 included in the FDC group. Daycare centres were defined by larger-group, centre-based care
42
43 provided by several professional carers. FDC and care in DCCs were provided either by the
44
45 municipality or on a private basis.
46
47
48
49
50

51 **Outcomes**

52
53 The outcomes of the study were symptoms of RTIs (cough, rhinorrhea, fever, or wheeze),
54
55 consumption of antibiotic medication, and parental absence from work as recorded on a day-
56
57
58
59
60

1
2
3 to-day basis in the study diaries. Antibiotic medications prescribed for any reason were
4 included, but the main indication was AOM. Any parental absence from work due to the
5 child's illness was recorded, and reasons were not limited to RTIs. Final outcome measures
6 constituted days per month with RTI symptoms (sick days), antibiotic medication, and
7 parental absence from work.
8
9
10
11
12

13 14 15 16 **Confounding factors**

17 The following potential confounders were taken into consideration: sex, siblings, season of
18 year at start of daycare, asthma in the parents, pets (cats or dogs at home), maternal post-
19 secondary education, and family monthly net income.[24-26] There was only a small
20 proportion of families with known smoking in either one of the parents (127 within the home
21 care, FDC and DCC groups), and since our results indicated reporting bias, this variable was
22 excluded from the analyses.
23
24
25
26
27
28
29
30
31
32
33

34 **Statistical analysis**

35 All statistical analyses were carried out using SAS version 9.4 (SAS, Cary, NC, USA). In all
36 our analyses P-values less than 0.05 were regarded as statistically significant. In the first step,
37 outcome variables were calculated as days per month for each child and aligned in a time
38 sequence relating to the beginning of daycare. Follow-up included a time of 6 months
39 previous to and 9 months after the beginning of daycare. Since children commenced daycare
40 at different ages, this meant that not all data of individual follow-up related to the fixed age
41 frame of 0-2 years. Data from outside this age range were excluded from the analysis using
42 the SAS MIXED procedure. For the home care group the equivalent age range of follow-up
43 was calculated using the mean age of starting daycare in the FDC and DCC groups (mean
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 starting age at 15 months), and resulted in a follow-up period from 9 to 24 months (6 months
4 prior to and 9 months after the mean start of daycare at 15 months).
5
6

7 If a child finished daycare during follow-up for any reason, data after discontinuation
8 of daycare were selectively excluded by the SAS MIXED procedure.
9

10 After comparisons using independent sample t-tests and variance analyses, data were
11 analyzed by repeated measures variance analysis comparing all three groups for type of
12 daycare and time, as well as their interaction. In a second step, we compared the difference of
13 least square means between all pairs of analyzed months and determined the shortest time
14 intervals during which statistically significant differences in our outcome measures could be
15 observed. The shortest time for a rise of outcome measures was determined from the last
16 month prior to starting daycare and for a decline from the peak of each outcome variable.
17 Although the distributions of monthly outcome variables (sick days, days with antibiotic
18 treatment, and parental absence from work) were skewed in our analyses (coefficient of
19 skewness up to 2.4 and kurtosis 7.6), the pairwise difference variables followed
20 approximately a normal distribution.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 In a third step, repeated measures variance analysis was repeated according to the
37 previous model, but stratifying according to individual confounding factors.
38
39

40 Sensitivity testing was performed by additional variance analyses excluding all data
41 from children who finished daycare during follow-up.
42
43
44
45
46

47 **Results**

48 Figure 1 shows numbers of children within follow-up. Within our cohort of 1570 children in
49 active follow-up, 276 (21.8%) of 1264 children with data on daycare arrangements at age 13
50 months attended daycare (11.7% in DCC, 10.1% in FDC) and 593 (55.0%) of 1079 children
51 attended daycare at the age of 24 months (29.5% in DCC and 25.0 % in FDC). At all ages the
52
53
54
55
56
57
58
59
60

1
2
3 vast majority of children in daycare spent over 5 hours per day at daycare (88.5% at age 13
4 months and 91.6% at age 24 months). Group sizes were clearly smaller for children attending
5 FDC as compared to DCC, with 88.3% (at 13 months) to 82.7% (at 24 months) of families
6 reporting a group size of less than 5 children per group. In DCC group sizes were variable
7 with the following results for the ages of 13 months (and 24 months): less than 5 children for
8 2.0%, 5-15 children for 86.8%, over 15 children for 11.1% (less than 5 children for 1.4%, 5-
9 15 children for 87.8%, over 15 children for 10.9%).

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Missing knowledge regarding the exact time of start of daycare, or earlier discontinuation of the study, led to a drop-out of a large part of cases, but these drop-outs occurred mostly before the start of daycare (figure 1). Baseline characteristics and mean (SD) values of outcome measures of children included in the analysis are shown in table 1. Mean age at the beginning of daycare was 15 months (SD 4.1). Start of daycare (FDC or DCC) occurred in 38.2% of cases during fall-winter (October to March) and in 61.8% during spring-summer (April to September). Over the whole follow-up period during daycare, children in DCC had only slightly more sick days per month (mean 5.54, SD 4.07) compared to children in FDC (mean 4.85, SD 3.49) or home care (mean 4.80, SD 3.39). There was no significant difference in days with antibiotic treatment. Parental absences from work could not be compared to the home care group, since the stay-at-home parent was not employed, and could therefore not be absent from work. Discontinuous daycare was observed in only a minority of children (11 children in FDC and 13 children in DCC). The results remained consistent in analyses performed without data from children who finished daycare during follow-up.

Table 1. Comparisons of baseline variables and outcome measures according to daycare type

	Type of daycare			P
	Home care (n = 350)	Family daycare (n = 245)	Daycare centre (n = 299)	
Age at the beginning of daycare, mean (SD)	NA	1.24 (0.37)	1.28 (0.35)	0.25*
Sex				
Males, number (%)	172 (49.1%)	124 (50.6%)	162 (54.1%)	0.4†
Born preterm (<37 gestational weeks), number (%)	18 (5.0%)	7 (2.9%)	14 (4.7%)	0.38†
Sick days per month, mean (SD)	4.80 (3.39)	4.85 (3.49)	5.54 (4.07)	0.03‡
Days with antibiotic treatment per month, mean (SD)	0.74 (1.41)	0.82 (1.36)	0.93 (1.14)	0.21‡
Days with parental absences from work per month, mean (SD)	NA	0.31 (0.38)	0.36 (0.45)	0.17*

NA, not applicable.

*Independent sample t-test

†Chi-square test

‡Unadjusted variance analysis

1
2
3 There was a peak of mean sick days per month at 2 months after the start of daycare
4 for both FDC and DCC (figure 2). The rise in sick days was stronger for the DCC group than
5 for the FDC group. We observed increases from 3.53 mean sick days (95% CI 2.83-4.24)
6 during the month prior to the start of daycare to 8.34 mean sick days per month (95% CI
7 7.25-9.43) for the FDC group, and from 3.79 (95% CI 3.04-4.53) mean sick days prior to the
8 start of daycare to 10.57 (95% CI 9.35-11.79) mean sick days per month for the DCC group.
9 For both of these groups, the rise of outcome measures was a transient phenomenon; sick
10 days were comparable to the home care group at 5 months after the start of daycare.
11 Antibiotic use (figure 3) and parental absences from work because of children's illnesses
12 (figure 4) rose and declined according to a pattern similar to that observed in sick days in
13 relation to the start of daycare, although this decline was less pronounced compared to sick
14 days.
15

16
17
18
19
20
21 The rise in sick days (figure 2) and antibiotic use (figure 3) was more rapid among
22 children starting DCC than in those starting FDC. Within the home care group there were no
23 significant findings in pairwise comparisons between months.
24
25

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

In repeated measures variance analyses there was a significant overall effect for time and type of daycare and their interaction for all outcome measures. This interaction showed the above described pattern of rise and decline, which was strongest for the DCC group and weaker for the FDC group. Within the home care group there was no such pattern, but variation was observed in the frequency of sick days per month over time. When analyzing sick days and antibiotic use between different types of daycare without the aspect of time, levels of these outcome measures were higher in the DCC group in comparison to the home care group ($P < 0.001$ for both measures) and FDC group ($P < 0.001$ and $P = 0.04$, respectively). There were no significant differences between the FDC and home care group.

1
2
3 The presence of older siblings in the family, higher post-secondary education in
4 mothers, and higher family income were all associated with a higher burden of disease (table
5
6
7 2).
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 2. Effects of confounding factors on the rates of RTI symptoms (sick days) according to daycare type

	Type of daycare						P*	P for interaction with mode of daycare
	Home care		Family daycare		Daycare centre			
	n/N (%)	Sick days per month, mean (SD)	n/N (%)	Sick days per month, mean (SD)	n/N (%)	Sick days per month, mean (SD)		
Older siblings							0.02	<0.001
Yes	162/ 350 (46.3)	5.55 (3.72)	99 /245 (40.4)	5.26 (3.33)	132/ 299 (44.1)	5.10 (3.72)		
No	188/ 350 (53.7)	4.16 (2.95)	146/ 245 (59.6)	4.58 (3.59)	167/ 299 (55.9)	5.88 (4.31)		
Higher post-secondary education in mother							0.009	0.27
Yes	211/ 341 (61.9)	5.12 (3.53)	161/ 239 (67.4)	5.12 (3.72)	203/ 293 (69.3)	5.66 (3.75)		
No	130/ 341 (38.1)	4.40 (3.17)	78 / 239 (32.6)	4.23 (2.97)	90/ 293 (30.7)	5.45 (4.79)		
Family net income under 2000 euros per month							0.005	0.37
Yes	83/ 343 (24.2)	4.15 (3.28)	33/ 239 (13.8)	3.43 (2.75)	56/ 294 (19.0)	5.29 (3.39)		
No	260/343 (75.8)	5.03 (3.43)	206/ 239 (86.1)	5.04 (3.57)	238/ 294 (81.0)	5.64 (4.23)		
Start of daycare during fall-winter (October-March)							0.39	0.61
Yes	NA	NA	100/ 245 (40.8)	4.90 (3.15)	108/ 299 (36.1)	5.69 (4.21)		
No	NA	NA	145/ 245 (59.2)	4.82 (3.73)	191/ 299 (63.9)	5.45 (4.0)		
Asthma in parents							0.95	0.72
Yes	35/ 340 (10.3)	4.65 (3.26)	26/ 234 (11.1)	5.54 (4.0)	30/ 287 (10.5)	5.83 (4.25)		
No	305/ 340 (89.7)	4.86 (3.43)	208/ 234 (88.9)	4.76 (3.45)	257/ 287 (89.5)	5.56 (4.08)		
Cat or dog at home							0.31	0.23
Yes	94/ 295 (31.9)	4.43 (3.07)	80/ 196 (40.8)	4.83 (3.07)	57/ 219 (26.0)	5.80 (4.28)		
No	201/ 295 (68.1)	5.14 (3.48)	116/ 196 (59.2)	4.92 (3.14)	162/ 219 (74.0)	5.68 (3.66)		

1
2
3
4
5 n/N, number of children per number of those with data available; NA, not applicable; RTI, respiratory tract infection.
6
7

8 *By univariate analysis, regardless of daycare type.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

For peer review only

1
2
3 In repeated measures variance analyses stratified according to confounding factors,
4
5 associations remained ($P < 0.001$ for the effect of time and $P < 0.001$ for daycare type for all
6
7 analyses regarding sick days, except for those with parental asthma, where $P = 0.004$ for
8
9 daycare type). Online supplementary figures S1 and S2 illustrate the association of start of
10
11 daycare with respiratory tract infections in children stratified according to the presence of
12
13 siblings and maternal post-secondary education. For children without older siblings, the rise
14
15 of sick days per month after start of daycare was more pronounced than for those with older
16
17 siblings.
18
19

20 21 22 23 **Discussion**

24
25 In this study there was a strong, but temporally limited effect of daycare initiation on the rate
26
27 of RTIs. This was strongest for the DCC group and weaker for the FDC group. An effect of
28
29 overall time in daycare,[14, 15, 17] as well as daycare type, on infections has been
30
31 documented in previous studies.[8, 10, 11, 13] Our study specifically focused on RTI
32
33 dynamics around the start of daycare.
34
35

36
37 We analyzed three main outcome measures. After showing a peak, the rate of sick
38
39 days returned near to baseline within our 9-month follow-up period from the start of daycare;
40
41 for antibiotic use there was more scatter with a less pronounced decline, and parental absence
42
43 from work did not return to baseline, which was to be expected due to the fact that both
44
45 parents tend to be at work after the start of daycare. Consideration of confounders did not
46
47 change our findings. The effect of older siblings on infections in relation to daycare has been
48
49 previously described,[17] and our findings are in line with previous results. The effects of
50
51 virus epidemics and season of year were minimized in our study setting, but they cannot be
52
53 completely excluded.
54
55
56
57
58
59
60

1
2
3 We consider the main strength of our study the detailed follow-up on a day-to-day
4 basis for a relatively large number of cases and in chronological relation to the start of
5 daycare. Previous cohort studies mostly assessed data from follow-up with intervals of
6 months or longer periods and without information on the exact starting point of daycare, such
7 that detailed chronological analyses for the time around daycare initiation could not be
8 carried out.[13, 15, 16, 26] Retrospective collection of data relating to symptoms poses
9 problems in some other studies, since it strongly relies on parents' memories.
10
11
12
13
14
15
16
17

18 There are some limitations to our study. Exposure to daycare was documented as
19 hours per day only. However, irregular daycare was rare and high daily exposure can be
20 extrapolated to high weekly or monthly exposure. Follow-up time was relatively short albeit
21 sufficient to demonstrate the above described effects. Outcome measures were based on self-
22 reported data and are thus subject to bias. Daily diary-keeping demanded regular participation
23 of families, although diaries were concise and easily completed. There were several
24 indicators that, for the small number of families with known smoking, diaries were not
25 completed as comprehensively as in the rest of the cohort, so that we had to exclude parental
26 smoking as a confounder from our study. The group with more detailed follow-up during
27 infections had access to our medical services free of charge, and thus may have been more
28 motivated to accurately document all experienced events. Inclusion into this group was
29 offered to all families without any selection criteria.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Due to strict requirements for comprehensive follow-up we lost a considerable
46 proportion of cases. For all cases lost, missing data were collected before or at the time of
47 daycare initiation, which means that the majority of drop-outs occurred before the critical
48 follow-up after daycare initiation. This decreases the potential effect of exclusion-related bias
49 on our results. In order to screen for bias, a comparison of background variables was
50
51
52
53
54
55
56
57
58
59
60

1
2
3 performed for non-responders and responders at 13 months of age, and there were only minor
4
5 differences between groups.[22]
6

7 Our cohort children had less often older siblings than all eligible children, and their
8
9 mothers had higher education and occupational status.[22] Otherwise our cohort represents
10
11 the Southwest Finnish population well. Although lower socioeconomic class has traditionally
12
13 been linked to an increased rate of infectious diseases,[27] a Swedish study found an
14
15 association between low social status, smaller likelihood to be cared for in out-of-home care,
16
17 and lesser consumption of medical services.[28] Within our cohort, social status, as indicated
18
19 by maternal education and family income, was also reflected by a higher rate of RTIs, but this
20
21 did not affect our findings relating to daycare.
22
23
24
25
26

27 **Conclusion**

28
29
30 In this longitudinal cohort study, the respiratory infectious disease burden was clearly related
31
32 to out-of-home care, but it decreased already within a short follow-up time of 9 months after
33
34 the start of daycare. Our findings demonstrate a clearly limited temporal impact of daycare-
35
36 related RTIs in early childhood, which has implications on a correctly focused approach
37
38 when supporting families with small children. Families, daycare providers and pediatricians
39
40 may be reassured of the transient nature of increased RTIs after the start of daycare.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole STEPS Study team, which includes our research physicians and study nurses.

Footnotes

Competing interests. The authors have no competing interests.

Funding. All phases of this study were supported by the University of Turku; the Åbo Akademi University; the Turku University Hospital; the Academy of Finland, Grants no. 123571, 140251, and 277535; the Foundation for Pediatric Research, Finland; and Research Funds from Specified Government Transfers, Hospital District of Southwest Finland. Dr. Schuez-Havupalo was also supported by the Turku University Foundation.

Data sharing statement: Enquiries regarding extra data can be emailed to Ville Peltola.

Contributors. LSH contributed to conception and design, the clinical follow-up and data collection, carried out data analyses, drafted the initial manuscript, and approved the final manuscript as submitted. LT and SK contributed to conception and design, the clinical follow-up and data collection, reviewed and revised the manuscript, and approved the final manuscript as submitted. AK contributed to conception and design, coordinated data collection, carried out data analyses, reviewed and revised the manuscript, and approved the final manuscript as submitted. VP conceptualized and designed the study, coordinated and supervised data collection and analyses, reviewed and revised the manuscript, and approved the final manuscript as submitted.

References

- 1 Doyle AB. Incidence of illness in early group and family day-care. *Pediatrics* 1976;58:607-13.
- 2 Strangert K. Respiratory illness in preschool children with different forms of day care. *Pediatrics* 1976;57:191-6.
- 3 Wald ER, Dashefsky B, Byers C, et al. Frequency and severity of infections in day care. *J Pediatr* 1988;112:540-6.
- 4 Fleming DW, Cochi SL, Hightower AW, et al. Childhood upper respiratory tract infections: to what degree is incidence affected by day-care attendance? *Pediatrics* 1987;79:55-60.
- 5 Nurmi T, Salminen E, Pönkä A. Infections and other illnesses of children in day-care centers in Helsinki. II: The economic losses. *Infection* 1991;19:331-5.
- 6 Holmes SJ, Morrow AL, Pickering LK. Child-care practices: effects of social change on the epidemiology of infectious diseases and antibiotic resistance. *Epidemiol Rev* 1996;18:10-28.
- 7 Thrane N, Olesen C, Mortensen J, et al. Influence of Day Care Attendance on the Use of Systemic Antibiotics in 0- to 2- Year- Old Children. *Pediatrics* 2001;107:e76.
- 8 Louhiala P, Jaakkola N, Ruotsalainen R, et al. Form of Day Care and Respiratory Infections among Finnish Children. *Am J Public Health* 1995;85:1109-1112.
- 9 Rovers MM, Zielhuis GA, Ingels K, et al. Day-care and otitis media in young children: a critical overview. *Eur J Pediatr* 1999;158:1-6.
- 10 Marx J, Osguthorpe J, Parsons G. Day care and the incidence of otitis media in young children. *Otolaryngol Head Neck Surg* 1995;112:695-9.
- 11 Hardy A, Fowler M. Child care arrangements and repeated ear infections in young children. *Am J Public Health* 1993;83:1321-1325.
- 12 Collet JP, Burtin P, Gillet J, et al. Risk of infectious diseases in children attending different types of day-care setting. Epicreche Research Group. *Respiration* 1994;61:16-9.
- 13 The National Institute of Child Health and Human Development Early Child Care Research Network. Child Care and Common Communicable Illnesses. Results From the National Institute of Child Health and Human Development Study of Early Child Care. *Arch Pediatr Adolesc Med* 2001;155:481-488.
- 14 Kamper-Jørgensen M, Wohlfahrt J, Simonsen J, et al. Population-Based Study of the Impact of Childcare Attendance on Hospitalizations for Acute Respiratory Infections. *Pediatrics* 2006;118:1439-46.
- 15 Zutavern A, Rzehak P, Brockow I, et al. Day care in relation to respiratory-tract and gastrointestinal infections in a German birth cohort study. *Acta Paediatr* 2007;96:1494-9.

1
2
3 16 Cote S, Petitcherc A, Raynault M-F., et al. Short- and Long-term Risk of Infections as a
4 Function of Group Child Care Attendance. *Arch Pediatr Adolesc Med* 2010;164:1132-7.
5

6 17 Hurwitz ES, Gunn WJ, Pinsky PF, et al. Risk of respiratory illness associated with day-
7 care attendance: a nationwide study. *Pediatrics* 1991;87:62-9.
8

9 18 Nafstad P, Hagen J, Øie L, et al. Day Care Centers and Respiratory Health. *Pediatrics*
10 1999;103:753-8.
11

12 19 de Hoog ML, Venekamp RP, van der Ent CK, et al. Impact of early daycare on healthcare
13 resource use related to upper respiratory tract infections during childhood: prospective
14 WHISTLER cohort study. *BMC Med* 2014;12:107.
15

16 20 Takala AK, Jero J, Kela E, et al. Risk factors for primary invasive pneumococcal disease
17 among children in Finland. *JAMA* 1995;273:859-64.
18

19 21 Ball TM, Holberg CJ, Aldous MB, et al. Influence of Attendance at Day Care on the
20 Common Cold From Birth Through 13 Years of Age. *Arch Pediatr Adolesc Med*
21 2002;156:121-126.
22

23 22 Lagström H, Rautava P, Kaljonen A, et al. Cohort profile: Steps to the healthy
24 development and well-being of children (the STEPS study). *Int J Epidemiol* 2013;42:1273-
25 84.
26

27 23 Toivonen L, Schuez-Havupalo L, Karppinen S, et al. Rhinovirus infections in the first 2
28 years of life. *Pediatrics* 2016;138:20161309.
29

30 24 Hatakka K, Piirainen L, Pohjavuori S, et al. Factors associated with acute respiratory
31 illness in day care children. *Scand J Infect Dis* 2010;42:704-11.
32

33 25 Koopman LP, Smit HA, Heijnen ML, et al. Respiratory infections in infants: interaction of
34 parental allergy, child care, and siblings - The PIAMA study. *Pediatrics* 2001;108:943-8.
35

36 26 Celedon J, Litonjua A, Weiss S, et al. Day Care Attendance in the First Year of Life and
37 Illness of the Upper and Lower Respiratory Tract in Children With a Familial History of
38 Atopy. *Pediatrics* 1999;104:495-500.
39

40 27 Cohen S. Social status and susceptibility to respiratory infections. *Ann N Y Acad Sci*
41 1999;896:246-53.
42

43 28 Hjern A, Haglund B, Rasmussen F, et al. Socio-economic differences in daycare
44 arrangements and use of medical care and antibiotics in Swedish preschool children. *Acta*
45 *Paediatr* 2000;89:1250-6.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Figure legends

Figure 1. Flow-diagram of the recruitment and follow-up of study children.

Figure 2. Means and 95% CIs (dashed lines) of sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month).

Figure 3. Means and 95% CIs (dashed lines) of days with antibiotic treatment per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month). Dashed lines: transiently below the significance level.

Figure 4. Means and 95% CIs (dashed lines) of days with parental absence from work per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month). Dashed lines: transiently below the significance level.




Flow-diagram of the recruitment and follow-up of study children.

66x50mm (600 x 600 DPI)

ew only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Means and 95% CIs (dashed lines) of sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month).

173x86mm (300 x 300 DPI)

review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Means and 95% CIs (dashed lines) of days with antibiotic treatment per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month). Dashed lines: transiently below the significance level.

173x87mm (300 x 300 DPI)

Review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Means and 95% CIs (dashed lines) of days with parental absence from work per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Horizontal lines indicate the shortest time for a significant rise (decline) from the last month prior to starting daycare (from the peak month). Dashed lines: transiently below the significance level.

173x87mm (300 x 300 DPI)

Review only

Supplemental material: figures S1 and S2.




Figure S1. Mean sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Data are shown for children with older siblings (A) and for children without older siblings (B).




Figure S2. Mean sick days per month according to daycare groups in relation to start of daycare. Negative months denote the period prior to start of daycare. Data are shown for children of mothers with lower (A) and higher (B) post-secondary education.

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of cohort studies

Manuscript ID bmjopen-2016-014635, revision 1. Schuez-Havupalo L et al. Daycare attendance and respiratory tract infections: a prospective birth cohort study

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	p 1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	p 2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	p 4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	p 5
Methods			
Study design	4	Present key elements of study design early in the paper	p 5
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	p 5-6
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	p 5-6
		(b) For matched studies, give matching criteria and number of exposed and unexposed	NA
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	p 6-7
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	p 5-7
Bias	9	Describe any efforts to address potential sources of bias	p 8 (cf. results/discussion)
Study size	10	Explain how the study size was arrived at	p 5-6
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	p 7-8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	p 7-8
		(b) Describe any methods used to examine subgroups and interactions	p 8
		(c) Explain how missing data were addressed	p 7-8 (cf. results/discussion)

		(d) If applicable, explain how loss to follow-up was addressed	p 8, 9
		(e) Describe any sensitivity analyses	p 8
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	p 8-9, fig 1
		(b) Give reasons for non-participation at each stage	fig 1
		(c) Consider use of a flow diagram	fig 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	p 8-10
		(b) Indicate number of participants with missing data for each variable of interest	p 8-9, fig 1
		(c) Summarise follow-up time (eg, average and total amount)	p 5, 7-8
Outcome data	15*	Report numbers of outcome events or summary measures over time	p 9, table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	p 11-15, table 2
		(b) Report category boundaries when continuous variables were categorized	tables 1 and 2
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	p 8-9
Discussion			
Key results	18	Summarise key results with reference to study objectives	p 15
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	p 15-17
Generalisability	21	Discuss the generalisability (external validity) of the study results	p 17
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	p 18

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

1
2
3
4 **Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE
5 checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at
6 <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

For peer review only