





80-

60

40-20-3 T 2-1-0

Neutrophil Lympocyte

800

ø

Eosinophil Basophil

8

85

Monocyte

Cell Composition (%)









С

150-

100

50-

0

B

÷

Cell number (x10⁶)











22°C HFD

30°C HFD









С

	b
abdus	
ed	
teriaceae	
ed	Bacteroidet
ales	Sphingobacterial
ed acteriales	
cus	
ccaceae	Eirmicut
llus	Thinked
xcaceae	
illales	1.
im_sensu_stricto	Lo
ed ceae 1	Charles and the
rax	Clostridial
les_Incertae_Sedis_XIII	Eŋ
im_XIVa	
m_XIVD	Erysi
ed	CI
iraceae	
m_XI find	Peptostr
reptococcaceae	Erysipelotrichace
runcus	Str
lum_IV	La
ractor	CI
fied	Clostridiun
coccaceae	C
fied	
ium_XVIII	Fr
cillus	-
otrichaceae_incertae_sedis	
fied	c
otrichaceae	Coriobacteriace
otrichales	conobacteriace
laceae	C
ansia	
microbiaceae	Clastridialas, Inc.
micropiales	Clostridiales_Ince
	, in the second s
	Clostridiaceae
	Peptostreptococcace
	Erysipelotrichaceae
	Bacteroidal
	Ruminococcace
	Lachnospirace
	Verruc
	v
	Verr
	Ve
	Rur









Supp Fig 7

6







100-

0

0

20

-D- WT - IL-17RA---

40 Minutes

HFD

60

90

🖶 IL-17A-/-

5000

n

ž IL17RA^{-/-} IL-17A^{-/-}

HFD











b



Table 1a Ts vs Tn PBMCs in Chow Fed Mice Supplement to Figure 1c

A = Decreased inflammatory response

B = Increased susceptibility to infection

C = Decreased acute inflamamtion

Gene Name	Fold Change	Log Fold Chang	p-value	Associated Pathway
Ccr2	-2.900268	-1.5361862	0.0179735	А, В, С
S100a9	-3.9386005	-1.9776831	0.0010768	A, C
Mmp19	-3.5310318	-1.8200898	0.01617527	A, C
Sele	-2.1671202	-1.1157792	0.1969843	А, В, С
Selp	-5.3711905	-2.425242	0.27060795	А, В, С
Fpr2	-3.1194942	-1.6413121	0.0179063	А, В, С
Clec4d	-4.0024166	-2.0008714	0.1282988	A, C
Mmp2	-2.9483821	-1.5599235	0.5034918	A
Spp1	-2.0498734	-1.0355347	0.0785838	А, В, С
Cysltr1	-2.6501172	-1.4060562	0.01161632	A, C
Tnfrsf11a	-2.299639	-1.2014074	0.05569378	A, C
Cd14	-2.3607728	-1.2392592	0.12847532	А, В, С
Cebpd	-2.9350784	-1.553399	0.03184895	A
Ccr1	-6.9453616	-2.7960498	0.05346816	A, C
ll1b	-2.1949959	-1.1342182	0.00162273	А, С
ll1r1	-2.256751	-1.1742473	4.52E-04	A, C
116	-3.610195	-1.8520766	0.12287884	А, В, С
Nlrp12	-2.782047	-1.4761468	0.0777145	A, C
Ptafr	-2.3166058	-1.2120125	0.11318901	A
Ptgdr	-2.2381516	-1.1623077	0.28921354	А, С
Ptgs1	-4.4043546	-2.1389306	0.29489788	A
Clec4e	-5.236093	-2.3884907	3.11E-04	A, C
Cers6	-2.0422053	-1.0301279	0.01177411	A
Lbp	-15.296995	-3.9351764	0.01348366	А, В, С
Ecm1	-3.1597195	-1.6597965	0.2930402	A, C
S1pr3	-3.8994565	-1.963273	0.16772528	A, C
Ednrb	-4.345838	-2.1196344	0.3942885	A, C
Ltc4s	-3.3450212	-1.7420154	0.52104604	A, C
Aicda	-6.6319103	-2.7294245	0.2532547	В
Bpifa1	-34.406723	-5.1046185	0.18959704	В
Mmp8	-5.7030334	-2.5117295	0.02418746	В
Fpr1	-3.0620682	-1.6145064	0.05993914	A
Msr1	-3.5590317	-1.8314848	0.08937913	В
Slc16a2	-2.064818	-1.0460145	0.41816664	В
Slpi	-5.138603	-2.3613763	0.2985094	В
Marco	-2.9889517	-1.5796396	0.5811841	В
Gbp3	-1.2843655	-0.36105585	0.01599239	В
Gpc3	-6.992956	-2.8059025	0.02658532	В
C1qa	-2.7704086	-1.4700987	0.50132746	В
C3	-2.7789478	-1.4745387	0.0182416	В
C4a	-4.980624	-2.3163266	0.17424007	В
C5ar1	-2.3440626	-1.229011	0.13113089	В
Cd5I	-4.0239587	-2.0086155	0.34872347	В
Tlr5	-5.115588	-2.3549001	0.01195425	В
Cfh	-2.7807312	-1.4754642	0.0563215	В
Padi4	-2.2842767	-1.1917374	0.3320089	В
Vsig4	-2.0757298	-1.0536187	0.62382525	В
Vwf	-2.5577865	-1.3548958	0.090023	В
Rbpj	-1.2285191	-0.29692027	0.1118677	В
Cxcr2	-2.7580562	-1.4636518	0.04135978	В
Dlg2	-3.9399412	-1.9781741	0.27481955	В
Jam3	-7.921922	-2.9858506	0.02596029	В
Dusp3	-2.2197046	-1.1503677	0.00344865	В
Lcn2	-3.122755	-1.6428194	0.07775574	В

b

Table 1b Ts vs Tn Liver in Chow Fed Mice

Supplement to Supplementary Figure 3f

A = Metabolism of lipids and lipoprotein

B = Fatty acid metabolic pathway C = Positive regulation of fatty acid oxidation

Gene Name	Fold Change	Log Fold Chan	p-value	Associated Pathway
Mgll	2.263733	1.1787038	0.00213764	А, В
Chat	2.1188455	1.0832783	0.01085947	A
Pex11a	2.6956837	1.4306512	0.00267294	A
Fabp1	2.518476	1.332551	0.01452876	А, В, С
Lpin1	9.916864	3.309884	0.03805904	А, В
Gpd2	2.9046128	1.5383458	0.04214943	A
Pnpla2	2.1270788	1.0888735	0.00152263	A
G0s2	2.284056	1.1915979	0.02547845	A
Ppara	2.0650632	1.046186	0.04076514	А, В, С
Cpt1a	2.1424978	1.0992937	0.00830894	А, В, С
Ppargc1b	3.6117263	1.8526886	0.00190269	A
Trib3	3.6351936	1.8620322	0.00677629	А, В
Prkd1	3.4745681	1.7968336	0.00992504	A
Acot4	2.8300936	1.5008497	0.00610927	В
Acot2	5.1152916	2.3548164	0.01852117	В
Por	2.4115312	1.2699494	0.01191301	В
Acot1	4.9764776	2.315125	0.02481512	В
Ehhadh	3.8523538	1.9457402	0.04431824	В

С

Table 1c Ts vs Tn Liver in HFD Fed Mice

Supplement to Figure 20

A = Collagen Formation

B = Positive regulation of apoptosis signaling pathway C = Top 200 marker games up regulated in the largelifered

C = Top 200 marker genes up-regulated in the 'proliferation' subclass of HCC				
Gene Name	Fold Change	Log Fold Chan	p-value	Associated Pathway
Col4a5	2.595481	1.3760018	0.02289232	A
Col9a2	3.503366	1.8087417	0.02282644	A
Adamts14	2.2215497	1.1515664	0.00680651	A
Agtr2	2.0226696	1.0162606	0.0129051	В
Ngfr	2.0322323	1.0230653	0.020088	В
Srpx	2.1963255	1.1350919	0.02551001	В
Nck2	2.0141392	1.0101633	0.03370601	В
Osm	2.8813698	1.5267549	0.02944854	В
Ррр3сс	2.104037	1.07316	0.00375977	В
Sox4	2.7958655	1.483295	0.01094856	С
Cdkn3	3.391282	1.7618308	0.03137689	С
Hk2	2.7739832	1.471959	0.01099649	С
Ddr1	2.311051	1.208549	0.02099115	С
Ect2	2.9456124	1.5585676	0.03136202	С
Cdk1	2.443331	1.2888494	0.01801695	С
Kif18b	2.1213937	1.0850123	0.01129002	С

d Table 1d Ts vs Tn PBMCs in Chow Fed Mice

Supplement to Figure 4a and 5a

- A = Cytokines and inflammatory response
- B = Positive regulation of TLR signaling
- C = Cellular Response to LPS
- D = Positive regulation of TLR4 signaling

E - Incleased IL-17 secretion				
Gene Name	Fold Change	Log Fold Char	p-value	Associated Pathway
115	7.01585	2.810618	0.03999559	A
II10	4.8075986	2.2653165	0.09900788	A, C
II15	2.1603541	1.1112678	0.23251489	A
Lbp	2.7008667	1.4334224	0.59380835	B, C, D
Tlr5	2.5381804	1.3437946	0.31909248	В, С
Ly96	2.7357821	1.4519533	0.07607134	С
Nr1d1	-1.623939	-0.6994975	0.37599245	С
Cyp17a1	12.413165	3.633799	0.02596169	С
Ltf	2.0185776	1.013339	0.36580375	В, С, В
ll1rn	2.6822443	1.4234406	0.03462122	D
Nfil3	2.4986777	1.3211648	0.03562596	D
Rag1	3.612676	1.8530678	0.28290012	D
ll17rc	1.9096935	0.9333411	0.42264974	D

Supp Table 1

Supplementary Figure Legends

Supplementary Figure 1. Thermoneutral housing does not alter basal immune cell numbers and composition. (**a-d**) 6 week old WT C57BL/6 male mice were housed at $T_S (22^{\circ}C)$ or $T_N (30^{\circ}C)$ conditions for a minimum of 3 weeks. (**a**) Total cell number, percent of hematopoietic stem and progenitor cells, and percent of lymphoid and myeloid cells in the bone marrow (n = 3 per group). (**b**) Total white blood cell number and levels of circulating immune cells (n = 7 per group). (**c**) Total cell number and cellular composition in the spleen (n = 6 per group at T_S , 5 per group at T_N). (**d**) Total cell number and cellular composition in the thymus (n = 6 per group at T_S , 5 per group at T_N). Blue denotes $T_S (22^{\circ}C)$; Red denotes $T_N (30^{\circ}C)$. (**a**, **c** and **d**) A single experiment. (**b**) A representative of 2 individual experiments. (**a-d**) Student's *t*-test.

Supplementary Figure 2. Thermoneutral housing's effect on adipose tissue, glucose metabolism, cholesterol and triglyceride levels. (**a**-**i**) 6 week-old WT C57BL/6 male mice were housed at T_S (22°C) or T_N (30°C) conditions for 24 weeks and fed a chow or HFD. (**a**) Representative epididymal white adipose tissue histology (n = 3 mice per group) by H&E staining at 20x magnification. Scale bars, 100µm. (**b**) Immune cell (CD45⁺) infiltration to epididymal white adipose tissue (n = 6 per group at T_S , 5 per group at T_N). (**c**) Mean fluorescent intensity (MFI) of Ly6c expression by CD11b+F4/80+ macrophages in white adipose tissue (n = 4 per group at T_S , 5 per group at T_N). (**d**) Glucose tolerance test and area under the curve after 14 weeks of diet (n = 8 per group). (**e**)

Glucose tolerance test and area under the curve after 20 weeks of diet (n = 8 per group). (**f**) Insulin tolerance test after 22 weeks of diet (n = 8 per group). (**g**) Hepatic cholesterol levels (n = 4 per group). (**h**) Serum triglyceride levels (n = 4 per group). (**i**) Serum cholesterol levels (n = 8 per group). Blue denotes T_s (22°C); Red denotes T_N (30°C). A representative of 2 individual experiments. Throughout *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. (**b**-i) One way ANOVA with *post hoc* Tukey's test. (**c**) Student's *t*-test.

Supplementary Figure 3. Thermoneutral housing alters hepatic gene expression. (a-d) 6 week-old WT C57BL/6 male mice were housed at T_S (22°C) or T_N (30°C) conditions for 24 weeks and fed a chow or HFD. (a) Expression of the indicated lipid mediator genes in the liver (n = 4 per group) of chow fed mice. (b) Expression of the indicated chemokine genes in the liver (n = 4 per group) of chow fed mice. (c) Expression of the indicated fibrosis marker genes in the liver (n = 4 per group) of chow fed mice. (d) Trichrome staining in liver of HFD fed mice (n = 4 mice per group) at 10x magnification. Scale bars, 250µm. (e) Heatmap of differential hepatic gene expression in mice housed at T_S or T_N and fed a chow or HFD for 8 weeks, as determined by RNAseq (n = 2 per group). (f) Upregulated gene expression pathways determined by RNASeq analysis comparing mice fed chow diet and housed at T_S or T_N (n = 2 per group). (g) Cytomap of relevant altered ontologies in livers of HFD fed T_N -housed or T_S housed mice. Blue denotes T_S (22°C); Red denotes T_N (30°C). (a-c) Data represents mean + s.e.m. (**a-e**) A single experiment. Throughout **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *****P* < 0.0001. (**b-d**) Student's *t*-test.

Supplementary Figure 4. Thermoneutral housing exacerbates obesity and NAFLD progression in the AKR strain of inbred mice. (a-k) 6 week-old AKR male mice were housed at T_S (22°C) or T_N (30°C) and fed HFD for 18 weeks (n = 3 per group). (a) Weight gain. (b) Epididymal, inguinal and perirenal white adipose tissue weights. (c) Liver weight. (d) Hepatic triglyceride levels. (e) Representative liver histology (n = 3 mice per group) by H&E staining at 20x and 40x magnification Scale bars, 100µm (left) and 50µm (right). (f) NAFLD activity score. (g) Expression of the indicated chemokine genes in the liver (h) Percent of hepatic CD11b⁺F4/80⁺ immune cells. (i) Serum ALT levels. (j) Expression of the indicated fibrosis marker genes in the liver. (k) Trichrome staining in liver of HFD fed mice (n = 3 mice per group) at 10x magnification. Scale bars, 250µm. Percent Trichrome positive pixels. Blue denotes T_S (22°C); Red denotes T_N (30°C). For bar graph, data represents mean + s.e.m. (**a**-**k**) a single experiment. Throughout *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. (**a-d** and **f-k**) Student's t-test.

Supplementary Figure 5. Adrenalectomy removes the inhibition of NAFLD progression associated with thermostressed housing. (**a-m**) 6 week-old WT C57BL/6 male mice with surgically removed adrenal glands were housed at T_s (22°C) or T_N (30°C) and fed chow (*n* = 4 per group) or HFD (*n* = 4 per group at T_s

and *n* = 5 per group at T_N) for 16 weeks. (a) Weight gain. (b) Epididymal white adipose tissue weight. (c) Inguinal white adipose tissue weight. (d) Perirenal white adipose tissue weight. (e) Liver weight. (f) Hepatic triglyceride levels. (g) Representative liver histology (*n* = 4 mice per group) by H&E staining at 20x and 40x magnification. Scale bars, 100µm (left and middle) and 50µm (right). (h) Expression of the indicated lipid mediator genes in the liver. (i) Expression of the indicated chemokine genes in the liver. (j) Percent of hepatic CD11b⁺F4/80⁺ immune cells. (k) Expression of the indicated chemokine genes in the liver. (I) Serum ALT levels. (m) Fold change in ALT levels as compared to nonadrenalectomized C57BL/6 mice fed HFD for 16 weeks at the same temperature. Blue denotes T_S (22°C); Red denotes T_N (30°C). For bar graphs, data represents mean + s.e.m. (a-m) A single experiment. Throughout **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *****P* < 0.0001. (a-f, k and I). One way ANOVA with *post hoc* Tukey's test. (h-j and m) Student's *t*-test.

Supplementary Figure 6. Thermoneutral housing in intestinal epithelial resistance and kinetic modulation of intestinal microbiome composition. (**a**,**b**) 6 week-old WT C57BL/6 male mice (n = 8 per group) were housed at T_s (22°C) or T_N (30°C) conditions for 24 weeks and fed a chow or HFD. (**a**) Representative intestinal histology (n = 4 mice per group) by H&E staining at20x magnification. Scale bar, 100µm. (**b**) Trans-epithelial resistance (n = 4 per group). (**c**-**e**) Kinetic phyla level analysis of T_s and T_N mice fed chow or HFD. Mice were separated into T_s or T_N rooms, housed for 2 weeks and fed a chow, prior to being separated

into a chow or HFD fed groups (c) Microbiome upon arrival, before room separation. (d) Microbiome after two-week acclimation. (e) Microbiome after 12 weeks of chow or HFD feeding. (f-k) Relative abundance of all microbiome phyla after 24 weeks (n = 8 mice per group). (f) Percent composition of actinobacteria. (g) Percent composition of bacteriodetes. (h) Percent composition of firmicutes. (i) Percent composition of proteobacteria. (j) Percent composition of tenericutes. (k) Percent composition of verrucomicrobia. Blue denotes T_S (22°C); Red denotes T_N (30°C). (a-b) A representative of 2 experiments. (c-k) A single experiment. (b and f-k) **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *****P* < 0.0001. One way ANOVA with *post hoc* Tukey's test.

Supplementary Figure 7. Thermoneutral housing alteration of the intestinal microbiome extends past the phyla level. (**a-c**) 6 week-old WT C57BL/6 male mice were housed at T_S (22°C) or T_N (30°C) conditions for 24 weeks and fed a chow or HFD (n = 8 per group). (**a**) LEfSe analysis generated cladogram displaying most differentially abundant taxa in T_S and T_N mice after 24 weeks of chow or HFD feeding. (**b**) LDA scores of differentially abundant taxa. (**c**) LEfSe analysis generated cladogram displaying most differential displaying most differentially abundant taxa in T_S and T_N mice after 24 weeks of T_S and T_N mice after 24 weeks of LEfSe analysis generated cladogram displaying most differentially abundant taxa. (**c**) LEfSe analysis generated cladogram displaying most differentially abundant taxa in T_S and T_N mice after 24 weeks of HFD only. (**a-c**) A single experiment.

Supplementary Figure 8. Hepatic phenotype of mice treated with gram-negative antibiotics. (**a-g**) 6 week-old WT C57BL/6 male mice (n = 8 per group) were housed at T_s (22°C) or T_N (30°C) and fed a HFD. After 8 weeks of HFD, mice

were mock-treated (Ctrl) or treated with antibiotics (Abx Tx) supplied in drinking water in addition to HFD exposure for additional 16 weeks. (**a**) Weight gain. (**b**) Epididymal white adipose tissue weight. (**c**) Inguinal white adipose tissue weight. (**d**) Perirenal white adipose tissue weight. (**e**) Liver weight. (**f**) Hepatic triglyceride levels. (**g**) Glucose tolerance test and area under the curve. Blue denotes T_s (22°C); Red denotes T_N (30°C). (**a**-**g**) A single experiment. Throughout **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *****P* < 0.0001. (**a**-**g**) One way ANOVA with *post hoc* Tukey's test.

Supplementary Figure 9. Hepatic phenotype of TLR4^{fl/fl}Vav1-Cre⁺ mice. (**a-g**) 6 week-old male TLR4^{fl/fl}Vav1-Cre⁻ mice housed at 22°C (n = 9 per group) and 30°C (n = 8 per group), and TLR4^{fl/fl}Vav1-Cre⁺ mice housed at 22°C (n = 7 per group) and 30°C (n = 5 per group) were fed HFD for 24 weeks. (**a**) Weight gain. (**b**) Epididymal white adipose tissue weight. (**c**) Inguinal white adipose tissue weight. (**d**) Perirenal white adipose tissue weight. (**e**) Liver weight. (**f**) Hepatic triglyceride levels. (**g**) Glucose tolerance test and area under the curve. Blue denotes T_S (22°C); Red denotes T_N (30°C). (**a-g**) 2 combined experiments. Throughout **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *****P* < 0.0001. (**a-g**) One way ANOVA with *post hoc* Tukey's test.

Supplementary Figure 10. LPS induction of IL-17 promoting cytokines and hepatic phenotype of the IL-17 axis deficient mice. (**a**-**c**) 6 week-old WT C57BL/6 male mice were housed at T_s (22°C) or T_N (30°C) conditions for 24 weeks and

fed a HFD (n = 3 per group). BMDCs derived from these mice were then stimulated with LPS and cytokine levels were quantified, (**a**) IL-6 and (**b**) IL-1 β . (**c**) Indicated gene expression in BMDCs. (**d**-**j**) 6 week old WT (n = 5 per group), IL-17RA^{-/-} (n = 4 per group) and IL-17A^{-/-} (n = 4 per group) mice were housed at T_N (30°C) for 24 weeks and fed a HFD. (**d**) Weight gain. (**e**) Epididymal white adipose tissue weight. (**f**) Inguinal white adipose tissue weight. (**g**) Perirenal white adipose tissue weight. (**h**) Hepatic triglyceride levels. (**i**) Glucose tolerance test and area under the curve. (**j**) Liver weight. Blue bars denote T_S (22°C); Red bars denote T_N (30°C). (**a-c**) A representative of 3 individual experiments. (**d**-**j**) A representative of 2 individual experiments. Throughout *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. (**a-c**) Student's *t*-test. (**d**-**j**) One way ANOVA with *post hoc* Tukey's test.

Supplementary Figure 11. Thermoneutrality-driven augmentation of hepatic SAA1 expression is linked to TLR4 and IL-17 axes. (**a**) Indicated gene expression in liver of 6 week-old WT C57BL/6 male mice housed at T_S (22°C; n = 4 per group) or T_N (30°C; n = 5 per group) for 24 weeks, fed a HFD. (**b**) Indicated gene expression in liver of 6 week old TLR4^{fl/fl}Vav1-Cre⁺ (n = 3 per group) and control TLR4^{fl/fl}Vav1-Cre⁻ (n = 3 per group) mice housed at T_N (30°C) conditions for 24 weeks, on a HFD. (**c**) Indicated gene expression in liver of 6 week old TLR4^{fl/fl}Vav1-Cre⁻ (n = 3 per group) and IL-17A^{-/-} (n = 3 per group) mice housed at T_N (30°C) conditions for 24 weeks, on a HFD. (**c**) Indicated gene expression in liver of 6 week old WT (n = 5 per group), IL-17RA^{-/-} (n = 4 per group) and IL-17A^{-/-} (n = 3 per group) mice housed at T_N (30°C) for 24 weeks, on a HFD. Blue denotes T_S (22°C); Red denotes T_N (30°C). A single experiment. Throughout *P < 0.05, **P < 0.01, ***P <

0.001, *****P* < 0.0001. (**a-b**) Student's *t*-test. (**c**) One way ANOVA with *post hoc* Tukey's test.

Supplementary Figure 12. Thermoneutral housing removes the barrier to modeling obesity in female mice. (**a-c**) 6 week-old WT C57BL/6 female mice were housed at $T_s (22^{\circ}C)$ fed chow (n = 4 per group) or HFD (n = 6 per group) or housed at $T_N (30^{\circ}C)$ and fed chow (n = 4 per group) or HFD (n = 6 per group) for 24 weeks. (**a**) Glucose tolerance test and area under the curve from glucose tolerance test. (**b**) Trichrome staining in liver of HFD fed mice (n = 4 mice per group) at 10x magnification. Scale bars, 250µm. (**c**) Indicated gene expression in liver of mice fed HFD. Blue denotes $T_S (22^{\circ}C)$; Red denotes $T_N (30^{\circ}C)$. Data represents mean + s.e.m. (**a-c**) A representative of 3 individual experiments. Throughout *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. (**a-b**) One way ANOVA with *post hoc* Tukey's test. (**c**) Student's *t*-test.

Supplementary Table 1. Genes associated with pathways upregulated by temperature and diet changes. (**a**-**d**) Genes significantly upregulated in gene associated pathways depicted in figures (n = 2 mice per group). (**a**) Genes associated with pathways depicted in Figure 1c. Genes are differentially expressed at least two-fold in PBMCs from T_N vs T_S chow fed mice. Negative fold change represents increased expression in T_S PBMCs. (**b**) Genes associated with pathways depicted in Supplemental Figure 3f. (**c**) Genes associated with pathways depicted in Figure 2o. Genes are differentially expressed at least two-

fold in the liver of T_N vs T_S HFD fed mice. Positive fold change represents increased expression in T_N liver. Genes are differentially expressed at least two-fold in the liver of T_N vs T_S chow fed mice. Positive fold change represents increased expression in T_N liver. (d) Genes associated with pathways depicted in Figures 4a and 5a. Genes are differentially expressed at least two-fold in PBMCs from T_N vs T_S HFD fed mice. Positive fold change represents increased expression in T_N BMCs.