

	Weighted %	aOR	CI	p
Pain				
No	8.1	(ref)		
Yes	21.3	2.39	1.96-2.91	<.0001
Age				
4-6	7.7	(ref)		
7-11	10.2	0.96	0.75-1.24	0.8
12-17	14.6	1.32	1.01-1.73	0.04
Sex				
Male	9.8	(ref)		
Female	13.5	1.5	1.27-1.78	<.0001
Race/Ethnicity				
White	15	(ref)		
Black	5.4	0.58	0.41-0.81	0.001
Hispanic	6.4	0.63	0.48-0.81	<0.0001
Other	14	0.98	0.73-1.32	0.89
Income				
\$0 - \$34,999	6.5	(ref)		
\$35,000 - \$74,999	10.6	1.28	0.99-1.66	0.06
\$75,000 - \$99,999	13.8	1.39	1.02-1.89	0.037
\$100,000 and over	18	1.59	1.19-2.15	0.002
Insurance Status				
Uninsured	7.8	(ref)		
Any private	14.8	0.99	0.66-1.48	0.97
Public only	7.5	0.86	0.57-1.29	0.47
Parental education				
Bachelor's or higher	17.5	(ref)		
Some college	11	0.62	0.49-0.77	<0.0001
HS or less	5.1	0.36	0.25-0.52	<0.0001
Region				
Northeast	10.9	(ref)		
Midwest	14.2	1.31	0.93-1.84	0.12
South	7.9	0.79	0.57-1.09	0.14
West	15.4	1.88	1.35-2.64	<.0001
Comorbidities				
0	8.2	(ref)		
1	12.4	1.3	1.06-1.60	0.014
2-3	16	1.75	1.37-2.25	<0.0001
4+	24	2.82	2.00-3.97	<0.0001
Has a personal healthcare provider				
Yes	8.3	(ref)		
No	12.4	1.2	0.91-1.60	0.2

aOR: adjusted Odds Ratio; CI: 95% Confidence Interval.

Supplemental Table A. Multivariate logistic regression analysis (adjusting for predisposing, enabling, and need factors) estimating the association between CAM use and sample sociodemographic and clinical characteristics.