

S1 Table. Median proportion of total foodborne cases attributed to exposure to specified foods, by subregions for selected parasites.

	AFR D	AFR E	AMR A	AMR B	AMR D	EMR B	EMR D	EUR A	EUR B	EUR C	SEAR B	SEAR D	WPR A	WPR B
<i>Number of Experts</i>	9	11	12	10	5	2	5	20	5	2	3	12	4	3
<i>Ascaris</i> spp.														
Percent of total disease that is foodborne*	0.38 (0.10- 0.66)	0.38 (0.07- 0.67)	0.83 (0.43- 0.97)	0.55 (0.17- 0.75)	0.37 (0.07- 0.68)	0.55 (0.15- 0.77)	0.55 (0.18- 0.75)	0.85 (0.47- 0.97)	0.55 (0.13- 0.76)	0.55 (0.14- 0.76)	0.54 (0.18- 0.75)	0.39 (0.11- 0.68)	0.85 (0.47- 0.97)	0.54 (0.16- 0.77)
Percent of foodborne disease attributed to:														
Fruits	0.21 (0.00- 0.48)	0.21 (0.00- 0.48)	0.06 (0.00- 0.36)	0.06 (0.00- 0.39)	0.21 (0.00- 0.47)	0.06 (0.00- 0.36)	0.22 (0.00- 0.48)	0.06 (0.00- 0.36)	0.06 (0.00- 0.37)	0.21 (0.02- 0.46)	0.06 (0.00- 0.40)	0.21 (0.00- 0.48)	0.06 (0.00- 0.39)	0.06 (0.00- 0.35)
Vegetables	0.77 (0.50- 0.98)	0.77 (0.50- 1.00)	0.94 (0.62- 1.00)	0.93 (0.60- 1.00)	0.77 (0.51- 0.98)	0.93 (0.62- 1.00)	0.77 (0.50- 0.99)	0.93 (0.63- 1.00)	0.93 (0.62- 1.00)	0.78 (0.52- 0.97)	0.92 (0.58- 1.00)	0.77 (0.50- 0.99)	0.93 (0.59- 1.00)	0.93 (0.62- 1.00)
Other	0.00 (0.00- 0.14)	0.00 (0.00- 0.15)	0.00 (0.00- 0.12)	0.00 (0.00- 0.12)	0.00 (0.00- 0.13)	0.00 (0.00- 0.13)	0.00 (0.00- 0.14)	0.00 (0.00- 0.13)	0.00 (0.00- 0.13)	0.00 (0.00- 0.14)	0.00 (0.00- 0.12)	0.00 (0.00- 0.14)	0.00 (0.00- 0.13)	0.00 (0.00- 0.13)
<i>Cryptosporidium</i> spp.														
Percent of total disease that is foodborne*	0.15 (0.00- 0.44)	0.15 (0.00- 0.47)	0.16 (0.01- 0.44)	0.11 (0.01- 0.38)	0.16 (0.01- 0.44)	0.09 (0.00- 0.41)	0.08 (0.00- 0.36)	0.10 (0.00- 0.39)	0.11 (0.00- 0.39)	0.09 (0.00- 0.40)	0.10 (0.00- 0.37)	0.10 (0.00- 0.42)	0.10 (0.00- 0.40)	0.10 (0.00- 0.45)
Percent of foodborne disease attributed to:														
Dairy	0.03 (0.00- 0.41)	0.04 (0.00- 0.46)	0.05 (0.00- 0.29)	0.08 (0.00- 0.35)	0.08 (0.00- 0.36)	0.04 (0.00- 0.49)	0.02 (0.00- 0.46)	0.04 (0.00- 0.40)	0.04 (0.00- 0.44)	0.05 (0.00- 0.49)	0.02 (0.00- 0.38)	0.02 (0.00- 0.40)	0.02 (0.00- 0.33)	0.02 (0.00- 0.44)
Fruits	0.31 (0.00- 0.78)	0.29 (0.00- 0.78)	0.31 (0.02- 0.71)	0.27 (0.01- 0.65)	0.24 (0.02- 0.63)	0.24 (0.02- 0.66)	0.23 (0.02- 0.63)	0.26 (0.01- 0.66)	0.25 (0.00- 0.67)	0.23 (0.01- 0.65)	0.25 (0.00- 0.63)	0.26 (0.02- 0.69)	0.25 (0.00- 0.61)	0.24 (0.02- 0.65)
Vegetables	0.56 (0.08- 0.91)	0.56 (0.07- 0.92)	0.61 (0.14- 0.88)	0.62 (0.18- 0.88)	0.65 (0.19- 0.89)	0.60 (0.13- 0.90)	0.62 (0.15- 0.91)	0.61 (0.19- 0.91)	0.60 (0.18- 0.91)	0.61 (0.18- 0.91)	0.64 (0.21- 0.91)	0.63 (0.21- 0.92)	0.64 (0.21- 0.92)	0.64 (0.21- 0.92)

Other	0.00 (0.00- 0.21)	0.00 (0.00- 0.21)	0.00 (0.00- 0.09)	0.00 (0.00- 0.10)	0.00 (0.00- 0.13)	0.02 (0.00- 0.21)	0.01 (0.00- 0.21)	0.02 (0.00- 0.13)	0.02 (0.00- 0.13)	0.02 (0.00- 0.14)	0.01 (0.00- 0.25)	0.01 (0.00- 0.20)	0.01 (0.00- 0.23)	0.01 (0.00- 0.16)
-------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------

	AFR D	AFR E	AMR A	AMR B	AMR D	EMR B	EMR D	EUR A	EUR B	EUR C	SEAR B	SEAR D	WPR A	WPR B
Number of Experts	9	11	12	10	5	2	5	20	5	2	3	12	4	3

Echinococcus granulosus

Percent of total disease that is foodborne*	0.21 (0.07- 0.42)	0.20 (0.05- 0.40)	0.20 (0.03- 0.40)	0.20 (0.02- 0.43)	0.21 (0.05- 0.41)	0.21 (0.05- 0.43)	0.21 (0.06- 0.41)	0.21 (0.04- 0.40)	0.21 (0.06- 0.40)	0.21 (0.04- 0.40)	0.21 (0.03- 0.44)	0.20 (0.06- 0.40)	0.20 (0.01- 0.39)	0.21 (0.05- 0.43)
---	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------

Percent of foodborne disease attributed to:

Fruits	0.22 (0.03- 0.53)	0.22 (0.03- 0.45)	0.22 (0.02- 0.53)	0.23 (0.04- 0.52)	0.22 (0.03- 0.52)	0.21 (0.02- 0.48)	0.23 (0.02- 0.49)	0.22 (0.02- 0.54)	0.21 (0.03- 0.49)	0.23 (0.02- 0.51)	0.22 (0.02- 0.55)	0.21 (0.01- 0.49)	0.21 (0.02- 0.48)	0.21 (0.01- 0.46)
Vegetables	0.77 (0.45- 0.96)	0.77 (0.49- 0.96)	0.77 (0.45- 0.98)	0.76 (0.44- 0.95)	0.77 (0.43- 0.96)	0.78 (0.50- 0.97)	0.77 (0.49- 0.98)	0.77 (0.40- 0.97)	0.78 (0.48- 0.96)	0.76 (0.47- 0.97)	0.77 (0.42- 0.97)	0.78 (0.50- 0.98)	0.78 (0.49- 0.98)	0.78 (0.51- 0.98)
Other	0.00 (0.00- 0.14)	0.00 (0.00- 0.25)	0.00 (0.00- 0.13)	0.00 (0.00- 0.22)	0.00 (0.00- 0.24)	0.00 (0.00- 0.13)	0.00 (0.00- 0.14)	0.00 (0.00- 0.31)	0.00 (0.00- 0.15)	0.00 (0.00- 0.13)	0.00 (0.00- 0.16)	0.00 (0.00- 0.14)	0.00 (0.00- 0.13)	0.00 (0.00- 0.17)

Echinococcus multilocularis

Percent of total disease that is foodborne*	0.51 (0.13- 0.79)		0.43 (0.09- 0.73)	0.48 (0.00- 0.77)	0.52 (0.15- 0.79)	0.45 (0.12- 0.72)	0.44 (0.12- 0.72)			0.58 (0.00- 0.88)	0.51 (0.09- 0.81)	0.48 (0.00- 0.78)		
---	-------------------------	--	-------------------------	-------------------------	-------------------------	-------------------------	-------------------------	--	--	-------------------------	-------------------------	-------------------------	--	--

Fruits	0.46 (0.06- 0.89)		0.44 (0.07- 0.86)	0.41 (0.00- 1.00)	0.46 (0.13- 0.89)	0.45 (0.11- 0.79)	0.43 (0.09- 0.76)			0.41 (0.00- 1.00)	0.45 (0.07- 0.89)	0.41 (0.00- 1.00)		
Vegetables	0.54 (0.10- 0.93)		0.54 (0.08- 0.89)	0.59 (0.00- 1.00)	0.53 (0.09- 0.86)	0.55 (0.20- 0.88)	0.54 (0.21- 0.87)			0.59 (0.00- 1.00)	0.55 (0.11- 0.93)	0.58 (0.00- 1.00)		

Other	0.00 (0.00- 0.01)	0.00 (0.00- 0.10)	0.00 (0.00- 0.00)	0.00 (0.00- 0.06)	0.00 (0.00- 0.09)	0.00 (0.00- 0.11)	0.00 (0.00- 0.00)	0.00 (0.00- 0.01)	0.00 (0.00- 0.22)	
<i>Entamoeba histolytica</i>										
Percent of total disease that is foodborne*	0.30 (0.00- 0.68)	0.30 (0.00- 0.68)	0.25 (0.00- 0.70)	0.21 (0.00- 0.62)	0.17 (0.00- 0.58)	0.24 (0.00- 0.62)	0.28 (0.00- 0.66)	0.33 (0.00- 0.71)	0.30 (0.00- 0.66)	0.26 (0.00- 0.64)
Fruits	0.32 (0.05- 0.66)	0.31 (0.05- 0.66)	0.25 (0.05- 0.59)	0.22 (0.05- 0.57)	0.17 (0.01- 0.57)	0.33 (0.04- 0.67)	0.33 (0.04- 0.68)	0.30 (0.04- 0.61)	0.33 (0.04- 0.67)	0.34 (0.05- 0.67)
Vegetables	0.64 (0.29- 0.90)	0.64 (0.29- 0.90)	0.71 (0.38- 0.91)	0.74 (0.39- 0.92)	0.79 (0.39- 0.96)	0.62 (0.28- 0.90)	0.62 (0.28- 0.89)	0.66 (0.35- 0.90)	0.62 (0.27- 0.90)	0.62 (0.28- 0.89)
Other	0.03 (0.00- 0.16)	0.03 (0.00- 0.17)	0.00 (0.00- 0.15)	0.00 (0.00- 0.15)	0.00 (0.00- 0.15)	0.03 (0.00- 0.16)	0.03 (0.00- 0.17)	0.03 (0.00- 0.15)	0.03 (0.00- 0.17)	0.03 (0.00- 0.15)

*Percent of total disease estimated to be foodborne is taken from Hald et al. (2016) and reported here to provide perspective. Numbers in parentheses are 5th and 95th percentile values.