

Supplementary material

Title

Complete reperfusion is required for maximal benefits of mechanical thrombectomy in stroke patients

Authors

Ángel Chamorro, MD, PhD¹; Jordi Blasco, MD²; Antonio López, MD²; Sergio Amaro MD, PhD¹; Luis San Román, MD, PhD²; Laura Llull MD, PhD¹; Arturo Renú MD¹; Salvatore Rudilosso, MD¹; Carlos Laredo PhD.¹; Victor Obach, MD.¹; Xabier Urrea, MD, PhD¹; Anna M. Planas PhD.³; Enrique C. Leira, MD, MS⁴; Juan Macho, MD, PhD²

Supplemental Table I Main baseline traits of the patients excluded from the study

	Patients included	Patients excluded	P value
N	125	222	-
Baseline traits			
Age yr, median	70 (61-80)	72 (63-79)	0.995
Males, %	53	48	0.366
Hypertension,%	51	57	0.290
Diabetes, %	16	25	0.050
Dyslipemia,%	38	39	0.815
CAD, %	15	19	0.341
Atrial fibrillation, %	28	34	0.304
Smoking, %	18	22	0.422
Pretreatment NIHSS, median (IQR)	16 (13-20)	17 (13-21)	0.237
NCCT ASPECTS >6, , median (IQR)	9 (8-10)	9 (8-10)	0.153
ASPECTS=Alberta Stroke Program Early CT score; NCCT = Non contrast CT scan			