

Supplemental Table 2. Percentage of Patients with Adverse Events by Grade for Relevant Adverse Event Categories

Category*	STAD + RT (n=763)					LTAD + RT (n=756)				
	Grade					Grade				
	1	2	3	4	5	1	2	3	4	5
Acute RT										
Skin	26.9	12.8	0.3	0.1	0	25.4	14.9	0.4	0.1	0
Bladder	37.0	25.3	4.9	0.4	0	37.8	24.2	2.9	0.4	0
Bowel	26.9	34.6	1.3	0	0	24.7	35.8	0.8	0	0
Other GI	9.0	5.4	0.1	0	0	6.6	7.4	0.1	0.1	0
Other GU	6.4	2.6	0.5	0.1	0	6.4	3.0	0.1	0.1	0
Other	30.1	42.1	3.9	0.8	0	29.2	42.7	3.8	0.9	0
Maximum (all acute RT)	30.0	55.0	8.5	1.31	0	27.8	54.0	6.6	1.7	0
Late RT										
Skin	5.5	0.9	0	0	0	7.5	0.7	0	0	0
Bladder	25.4	12.1	4.3	0.1	0	21.4	15.9	3.6	1.3	0
Bowel	28.8	13.5	0.8	0.3	0.1	28.0	13.1	1.9	0.3	0.3
Other GI	9.8	2.9	0.3	0	0	13.2	3.8	0.5	0.1	0
Other GU	15.1	5.2	1.2	0.1	0	15.9	6.4	0.8	0.4	0
Other	11.4	3.3	0.8	0	0	10.3	4.9	1.2	0.1	0
Maximum (all late RT)	36.4	25.8	6.6	0.5	0.1	31.3	26.9	6.9	2.0	0.3
Toxicities associated With hormone use										
Nausea	0	0.4	0	0	0	0	0	0.3	0	0
Vomiting	0.1	0.3	0	0	0	0	0.1	0.1	0	0
Diarrhea	1.6	4.2	0.9	0.1	0	0	2.8	0.9	0.1	0
Headache	0	0	0.1	0	0	0	0	0	0	0
Fluid Retention	0	0.1	0	0	0	0	0	0	0	0
Gynecomastia	0	0	0	0	0	0.1	0	0	0	0
Skin Rash	0.1	0.1	0	0	0	0.1	0.1	0.1	0	0
Infection	0.1	0.1	0	0	0	0	0.1	0	0	0
Elevated SGOT	6.7	2.6	1.0	0.1	0	12.6	2.9	1.5	0.1	0.1
Thromboph/Emboli	0.1	0	0	0	0	0	0	0	0	0
Cardiac	0	0	0.1	0	0	0	0	0.26	0	0
Hot Flashes	0	0	0	0	0	0.1	0	0	0	0
Impotence	0	0	0.1	0	0	0	0	0	0	0
Other	5.2	2.1	2.0	0.1	0	8.5	1.9	1.7	0.4	0.1
Maximum (all hormone associated)	9.4	7.3	3.8	0.3	0	15.7	56.0	4.5	0.7	0.1

*Toxicities were graded with NCI Common Toxicity Criteria (CTC) version 2.0.