

Supplementary Table 7. Studies reporting harms and adverse events associated with percutaneous renal mass sampling

Author, year	N	Radiation exposure, n (%)	Hemorrhage, n (%)	Hematoma, n (%)	Pain, n (%)	Pneumothorax, n (%)	Tumor Seeding, n (%)	Hematuria, n (%)	Other
Campbell, 1997 ¹²	25	25(100)	0	10(40)	NR	0	0	NR	
Chyhrai, 2010 ¹³	25	0	1(4)	NR	1(4)	0	0	NR	
Harisinghani, 2003 ¹⁵	28	28(100)	NR	NR	NR	NR	NR	NR	
Leveridge, 2011 ¹⁶	227*	151 (of 336) (45)	0	22(10)	NR	2(1)	0(0)	3(1)*	1 syncope
Millet, 2012 ¹⁹	187	187(100)	NR	NR	NR	NR	NR	NR	
Neuzillet, 2003 ²⁰	88	88(100)	0	0	0	NR	0	NR	
Park, 2013 ²¹	59	0	0	9(15)	3(5)	0	NR	0	
Prince, 2015 ³⁰	565	48(9)	3(0.5)**	NR	NR	NR	NR	NR	7 hospital admissions
Reichelt, 2007 ²⁸	30	0	1(3)	1(3)	1(3)	NR	NR	NR	
Richard, 2015 ⁵	492	115 (of 503) (23)	NR	24(5)	3(0.6)	NR	0	5(1)	12 coaxial sheath venous bleeding, 2 dizziness, 2 nausea/vomiting, 1 skin hematoma
Salem, 2012 ²²	145	145(100)	0	2(1)	0	NR	0	NR	1 flank ecchymosis
Schmidbauer, 2008 ²⁹	78	78(100)	0	4(5)	NR	1(1)	NR	NR	
Shannon, 2008 ²³	221	NR	1(0.4)	1(0.4)	NR	NR	0	NR	
Sofikerim, 2009 ²⁴	42	42(100)	0	0	0	NR	0	0	
Volpe, 2008 ²⁰	100	55(55)	0	NR	1(1)	1(1)	0	NR	1 syncope
Wang, 2009 ²⁷	110	66(60)	NR	2(2)	4(4)	0	NR	NR	1 wound infection, 1 hypotension requiring fluid resuscitation
Summary		1028(44)	6(0.4)	75(5)	13(1.2)	4(0.6)	0(0)	8(1)	

n= sample size; NR=Not Reported

*67 patients without complication data available; Clavien 1 complications in 10.1% of patients; one Clavien 3a complication (0.3%) noted involving gross hematuria leading to urinary retention due to formation of clots

**One Clavien 3a complication (selective renal artery embolization for bleeding leading to hemodynamic instability)

***One Clavien 3b complication (required percutaneous embolization)