

Supplemental Table I. Endovascular vs Open Preoperative Variables for Unmatched Entire Cohort

Parameter	Endovascular	Open	p-value
Patients	2935 (36.4%)	5131 (63.6%)	
Age	70 ± 12	68 ± 11	<0.0001
Sex (male)	1656 (56.4%)	3182 (62.0%)	<0.0001
Race (white)	2287 (78.3%)	3970 (77.4%)	0.015
Transfer Status	409 (14.0%)	614 (12.0%)	0.029
Inpatient Admission	1971 (67.2%)	5097 (99.3%)	<0.0001
ASA >2	2152 (73.3%)	4944 (96.4%)	<0.0001
Preoperative Antiplatelet Therapy	2406 (82.0%)	4107 (80.0%)	<0.0001
Preoperative Statin Therapy	1981 (67.5%)	3557 (69.3%)	<0.0001
Steroid Use	229 (7.8%)	277 (5.4%)	<0.0001
Bleeding Disorder	947 (32.3%)	1256 (24.3%)	<0.0001
Ascites	3 (0.1%)	8 (0.2%)	0.809
Congestive Heart Failure	134 (4.6%)	176 (3.6%)	0.002
Chronic Obstructive Pulmonary Disease	293 (10.0%)	661 (12.9%)	0.0003
Hypertension Medication	2525 (86.0%)	4313 (84.1%)	0.001
Tobacco Use	777 (26.5%)	2044 (39.9%)	<0.0001
Dialysis	426 (14.5%)	403 (7.9%)	<0.0001
Insulin Dependent Diabetes	1278 (43.5%)	1630 (31.7%)	<0.0001
Dependent Functional Status	457 (15.6%)	469 (9.1%)	<0.0001
Tissue Loss	1960 (66.8%)	2987 (58.2%)	<0.0001
Infrageniculate Revascularization	832 (28.4%)	2372 (46.2%)	<0.0001
Prior Bypass of Current Segment	461 (15.7%)	1145 (22.3%)	<0.0001
Prior Percutaneous Intervention of Current Segment	683 (23.3%)	884 (17.2%)	<0.0001