

Critical Role of ADAMTS-4 in the Development of Sporadic Aortic Aneurysm and Dissection in Mice

Pingping Ren^{1†}, Michael Hughes^{1†}, Swapna Krishnamoorthy¹, Sili Zou^{1,3}, Lin Zhang¹, Darrell Wu^{1,7}, Chen Zhang¹, John A. Curci⁴, Joseph S. Coselli^{1,2,6}, Dianna M. Milewicz⁵, Scott A. LeMaire^{1,2,6,7*}, & Ying H. Shen^{1,2,6,*}

¹Division of Cardiothoracic Surgery, Michael E. DeBakey Department of Surgery, Baylor College of Medicine, Houston, Texas, USA. ²Department of Cardiovascular Surgery, Texas Heart Institute, Houston, Texas, USA. ³Department of Vascular Surgery, Changzheng Hospital, Second Military Medical University, Shanghai, China. ⁴Division of Vascular Surgery, Vanderbilt University School of Medicine, Nashville, Tennessee, USA. ⁵Division of Medical Genetics, Department of Internal Medicine, The University of Texas Health Science Center at Houston, Houston, Texas, USA. ⁶Cardiovascular Research Institute, Baylor College of Medicine, Houston, Texas, USA. ⁷Department of Molecular Physiology and Biophysics, Baylor College of Medicine, Houston, Texas, USA.

[†]These authors contributed equally to the study.

*Correspondence and requests for materials should be addressed to Ying H. Shen, One Baylor Plaza, BCM 390, Houston, Texas 77030, USA. Phone 832-355-9952; fax 832-355-9951; e-mail hyshen@bcm.edu; and Scott LeMaire, MD, One Baylor Plaza, BCM 390, Houston, Texas 77030, USA. Phone 832-355-9942; fax 832-355-9948; e-mail slemaire@bcm.edu.

Figure S1




Figure S1. Representative western blot images of aortas from unchallenged WT mice (n=4) and challenged WT mice (n=5). Quantification studies showing the increased expression of ADAMTS-1 in the aortas of challenged WT mice versus unchallenged WT mice.

Figure S2




Figure S2. Free fatty acid levels in serum were significantly higher in challenged WT mice than in unchallenged WT mice.