

Suppression of adiponectin receptor 1 promotes memory dysfunction and Alzheimer's disease-like pathologies

Min Woo Kim^{1,‡}, Noman bin Abid^{1, ‡}, Myeong Hoon Jo¹, Min Gi Jo¹, Gwang Ho Yoon¹, and Myeong Ok Kim^{1, *}

¹ Division of Life Science and Applied Life Science (BK 21 plus), College of Natural Sciences, Gyeongsang National University, Jinju, Republic of Korea

‡ These authors contributed equally to this work.

*** Correspondence:**

Prof. Myeong Ok Kim, Ph.D.

Head of Neuroscience Pioneer Research Center,

Head of Brain-Metabolic Neurodegenerative Disease Center,

Division of Life Science and Applied Life Science, College of Natural Sciences,


Gyeongsang National University, Jinju, 660-701, South Korea

Tel.: +82-55-772-1345

Fax: +82-55-772-1349

To whom correspondence should be addressed: E-mail: mokim@gnu.ac.kr

Supplementary fig. 1



Supplementary figure. 1 AD-like pathologies in AKD mice (A). Oligomeric A β , GSK3 β and hyperphosphorylated tau expression in the AKD mice. (B) Neuroinflammatory marker including iba-1 and TNF- α in the AKD mice

Complete Immunoblots:

Figure 1a




Figure 1b




Figure 1c




Figure 5c




Figure 6a






Figure 6c



Supplementary fig. 1a



Supplementary fig. 1b

