

De Boer et al. DNA metabarcoding of orchid-derived products reveals widespread illegal orchid trade. Proceedings of the Royal Society B. DOI: 10.1098/rspb.2017-1182

Table S4. Plant species detected in more than 20 % of samples, excluding orchids (presence as % in total samples in parentheses).

| | |
|--|-----|
| <i>Aegilops lorentii</i> Hochst. | 47% |
| <i>Aegilops speltoides</i> Tausch | 27% |
| <i>Anthosachne multiflora</i> (Banks & Sol. ex Hook.f.) C.Yen & J.L.Yang | 20% |
| <i>Coriandrum sativum</i> L. | 23% |
| <i>Cucumis melo</i> L. | 20% |
| <i>Cyamopsis tetragonoloba</i> (L.) Taub. | 70% |
| <i>Ficaria verna</i> Huds. | 33% |
| <i>Hordeum vulgare</i> L. | 33% |
| <i>Ipomoea eriocarpa</i> R.Br. | 27% |
| <i>Ipomoea pes-tigridis</i> L. | 50% |
| <i>Medicago truncatula</i> Gaertn. | 30% |
| <i>Momordica charantia</i> L. | 20% |
| <i>Prunus mahaleb</i> L. | 33% |
| <i>Prunus pseudocerasus</i> Lindl. | 20% |
| <i>Secale cereale</i> L. | 43% |
| <i>Triticum aestivum</i> L. | 43% |
| <i>Triticum dicoccoides</i> (Körn. ex Asch. & Graebn.) Schweinf. | 60% |
| <i>Triticum durum</i> Desf. | 47% |
| <i>Triticum monococcum</i> L. | 20% |
| <i>Zea mays</i> L. | 20% |