

Supplementary information

Nuclear factor (erythroid derived 2)-like 2 activation increases exercise endurance capacity via redox modulation in skeletal muscles

Sechang Oh^{1,2,5,+}, Shoichi Komine^{2,+}, Eiji Warabi², Kentaro Akiyama^{2,5}, Akiko Ishii², Kazunori Ishige², Yuji Mizokami², Keisuke Kuga², Masaki Horie², Yoshihiro Miwa², Takao Iwawaki³, Masayuki Yamamoto⁴, and Junichi Shoda^{1,2,*}

¹The Center of Sports Medicine and Health Sciences, Tsukuba University Hospital, Tsukuba, Ibaraki 305-8576, Japan;



²Faculty of Medicine, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan;

³Department of Life Science, Medical Research Institute, Kanazawa Medical University, Uchinada, Kahoku, Ishikawa 920-0293, Japan;

⁴Department of Medical Biochemistry, Tohoku University Graduate School of Medicine, Sendai, Miyagi 980-8575, Japan;


⁵Japan Society for the Promotion of Science, Tokyo 102-0083, Japan

⁺These authors contributed equally to this work.



Supplementary figure S1.

Fig. S1. SFN-induced *Nrf2*-Luc activity detected using an In Vivo Imaging System and the results of genotyping in *Nrf2* and OKD48 transgenic mice in the supine position. Time course of the Luciferase assay is shown (a) ~ (e)



Supplementary figure S2. Full-length image of figure 3 (A)

The origin of these panels is identical with figure 3 (A) showing western blotting with phosphor-AMPK, AMPK, SIRT1, Actin, PGC1α, and Lamin A/C antibody