

Supplementary Table 1 : NFKB1 -94 ins/del ATTG and NFKB1A -826 C/T polymorphism with clinicopathological variables in HNSCC.

Variable	del/del	del/ins+ins/ins	OR (95% C.I.)	p value	AOR (95% C.I.)	p value
<b>NFKB1</b>						
<b>Clinical Stage</b>						
I+II	3(6.38%)	44(93.62%)	1 (ref)		1 (ref)	
III+IV	15(5.66%)	250(94.34%)	1.13(0.31-4.08)	0.845	1.28(0.34-4.75)	0.703
<b>Tumor Size</b>						
T1+T2	3(4.29%)	67(95.71%)	1 (ref)		1 (ref)	
T3+T4	15(6.20%)	227(93.80%)	0.67(0.19-2.41)	0.548	0.78(0.21-2.85)	0.712
<b>Lymph Node Metastasis</b>						
No	8(6.30%)	119(93.70%)	1 (ref)		1 (ref)	
Yes	10(5.41%)	175(94.59%)	1.17(0.45-3.06)	0.740	1.22(0.46-3.24)	0.687
NFKB1A Variable	CC	CT+TT	OR (95% C.I.)	p value	AOR (95% C.I.)	p value
<b>Clinical Stage</b>						
I+II	4(8.51%)	43(91.49%)	1 (ref)		1 (ref)	
III+IV	28(10.57%)	237(89.43%)	0.78(0.26-2.35)	0.669	0.83(0.27-2.55)	0.755
<b>Tumor Size</b>						
T1+T2	6(8.57%)	64(91.43%)	1 (ref)		1 (ref)	
T3+T4	26(10.74%)	216(89.26%)	0.77(0.30-1.97)	0.599	0.75(0.29-1.94)	0.555
<b>Lymph Node Metastasis</b>						
No	10(7.87%)	117(92.13%)	1 (ref)		1 (ref)	
Yes	22(11.89%)	163(88.11%)	0.63(0.28-1.38)	0.254	0.66(0.29-0.47)	0.316

- Abbreviations: OR: Odd Ratios, AOR; Adjusted Odd Ratios
- ORs with 95% CIs were estimated by logistic regression models.
- AORs with 95% CIs were estimated by multiple logistic regression models after controlling for sex, age, tobacco chewing, alcohol consumption, and smoking status.