

Human single chain-transbodies that bound to domain-I of non-structural protein 5A (NS5A) of hepatitis C virus


Kittirat Glab-ampai, Monrat Chulanetra, Aijaz Ahmad Malik, Thanate Juntadech, Jeeraphong

Thanongsaksrikul, Potjane Srimanote, Kanyarat Thueng-in, Nitat Sookrung, Pongsri



Tongtawe & Wanpen Chaicumpa

Supplementary Table 1. PCR primers used for amplification of DNA sequences coding for NS5A domains I, II, and III


Primer name	Sequence	Size of amplicon (bp)
D1 forward	5'-GAATTCCTCCCCTTCATCTCTTGTCAAAGG-3'	555
D1 reverse	5'-GTCGACTTAAGTCTCCGCCGTGATGTGGGG-3'	
D2 forward	5'-GAATTCTATGACGTGGACATGGTCGATGCC-3'	280
D2 reverse	5'-GTCGACTTAACAACCAGCAACGGTGGGCGG-3'	
D3 forward	5'-GAATTCAGGAGACGCCGGACAGTGGGT-3'	360
D3 reverse	5'-GTCGACTTAGCAGCACACGGTGGTATCGTC-3'	



Supplementary Figure S1. Unprocessed immunoblots that relate to Figure 2.



Supplementary Figure S2. Unprocessed immunoblots that relate to Figure 5.



Supplementary Figure S3. Unprocessed immunoblots that relate to Figure 10.