

Time points			
Groups	24h	48h	72h
<b><i>TLR-2</i></b>			
Control	- <sup>a</sup>	- <sup>a</sup>	- <sup>a</sup>
Septic	23.20±5.76	62.9±0.55 <sup>b</sup>	110.7±0.82 <sup>c, d</sup>
<b><i>TLR-3</i></b>			
Control	- <sup>a</sup>	- <sup>a</sup>	- <sup>a</sup>
Septic	2.85±0.36	5.08±0.77 <sup>b</sup>	17.58±0.37 <sup>c, d</sup>
<b><i>TLR-4</i></b>			
Control	- <sup>a</sup>	- <sup>a</sup>	- <sup>a</sup>
Septic	21.64±0.39	76.16±1.15 <sup>b</sup>	112.2±1.78 <sup>c, d</sup>
<b><i>TLR-7</i></b>			
Control	- <sup>a</sup>	- <sup>a</sup>	- <sup>a</sup>
Septic	5.64±0.54	45.93±0.82 <sup>b</sup>	50.71±0.5 <sup>c, d</sup>

---

**Supplemental table 3.** PCR showing changes of TLR 2, 3, 4 and 7 in lung tissues at the experimental time points. Values are expressed as the mean±SD. Statistical significances ( $p<0.05$ ) between the groups at the same time point are indicated as follows: a Control (C) vs Septic (S); b: 24S vs 48S; c: 48S vs 72S; d: 24S vs 72S.