

Time points			
Groups	24h	48h	72h
<b><i>TLR-2</i></b>			
Control	-	- <sup>a</sup>	- <sup>a</sup>
Septic	4.86±0.53	15.79±0.49 <sup>b</sup>	79.76±6.38 <sup>c, d</sup>
<b><i>TLR-3</i></b>			
Control	-	- <sup>a</sup>	- <sup>a</sup>
Septic	4.12±0.13	7.75±0.3	31±4.7 <sup>c, d</sup>
<b><i>TLR-4</i></b>			
Control	- <sup>a</sup>	- <sup>a</sup>	- <sup>a</sup>
Septic	7.16±0.86	15.06±0.58 <sup>b</sup>	75.19±2.31 <sup>c, d</sup>
<b><i>TLR-7</i></b>			
Control	- <sup>a</sup>	- <sup>a</sup>	- <sup>a</sup>
Septic	22.5±1.1	32.8±0.25 <sup>b</sup>	55.63±4.87 <sup>c, d</sup>

---

**Supplemental table 4.** PCR showing changes of TLR 2, 3, 4 and 7 in the intestine at the experimental time points. Values are expressed as the mean±SD. Statistical significances ( $p<0.05$ ) between the groups at the same time point are indicated as follows: a Control (C) vs Septic (S); b: 24S vs 48S; c: 48S vs 72S; d: 24S vs 72S.